

	Smoking cigarette	-0.068 (-0.087 to -0.050)	< 0.001	-0.036 (-0.051 to -0.021)	< 0.001	-0.010 (-0.022 to 0.003)	0.1296	0.027 (0.007 to 0.046)	0.0068	0.899 (0.868 to 0.931)	< 0.001
	Using e- cigarette	0.089 (0.031 to 0.147)	0.0025	0.239 (0.193 to 0.284)	< 0.001	-0.177 (-0.213 to -0.142)	< 0.001	-0.416 (-0.473 to -0.358)	< 0.001	0.885 (0.855 to 0.917)	0.5523
No	Total smoker	-0.101 (-0.106 to -0.096)	< 0.001	-0.048 (-0.052 to -0.045)	< 0.001	-0.024 (-0.028 to -0.021)	< 0.001	0.024 (0.019 to 0.029)	< 0.001	0.838 (0.830 to 0.846)	< 0.001
	Smoking cigarette	-0.109 (-0.114 to -0.104)	< 0.001	-0.051 (-0.055 to -0.047)	< 0.001	-0.027 (-0.030 to -0.024)	< 0.001	0.024 (0.019 to 0.030)	< 0.001	0.824 (0.816 to 0.832)	< 0.001
	Using e- cigarette	0.069 (0.053 to 0.086)	< 0.001	0.248 (0.235 to 0.261)	< 0.001	-0.205 (-0.215 to -0.195)	< 0.001	-0.453 (-0.469 to -0.437)	< 0.001	0.776 (0.752 to 0.802)	< 0.001

Note. BMI, body mass index; CI, confidence interval; OR, odds ratio. Data were estimated using a linear regression model described as β -coefficients with 95% confidence interval (CI) and a binary logistic regression model as odds ratio (OR) with 95% CI. Numbers in bold indicate a significant difference ($P < 0.05$).