

Association between Childhood Leukaemia and Exposure to Power-frequency Magnetic Fields in Middle Europe

Vitezslav JIRIK^{1,2}, Ludek PEKAREK³, Vladimir JANOUT¹, and Hana TOMASKOVA¹

1. Department of Epidemiology and Public Health, Faculty of Medicine, University of Ostrava, Czech Republic; 2. Department of Biophysics, Faculty of Science, Palacky University Olomouc, Czech Republic; 3. National Institute of Public Health, Prague, Czech Republic

Abstract

Objective Higher levels of exposure to extremely low-frequency magnetic fields (ELF-MF) are associated with a slightly increased risk of childhood leukaemia. Compared with more-developed Western countries, higher exposure levels are evident in the Czech Republic, probably because of the different types of housing. In light of this, we aimed to examine the association between ELF-MF exposure and childhood leukaemia in the Czech Republic.

Methods We conducted a paired case-control study. The cases (children with leukaemia) were age-sex- and permanent residence-matched to controls (children without leukaemia). Although this limited potential bias and confounding, it also limited our number of participants.

Results The matched analyses included 79 case-control pairs. No significant association between ELF-MF exposure and childhood leukaemia was observed for exposures over 0.2 μT (odds ratio [OR]=0.93, confidence interval [CI]=0.45-1.93), 0.3 μT (OR=0.77, CI=0.34-1.75), or 0.4 μT (OR=0.9, CI=0.37-2.22).

Conclusion Despite higher levels of exposure in Middle and Eastern Europe, no indication of an association between ELF-MF exposure and childhood leukaemia was determined. This in contrast to the findings of previous studies conducted in different countries.

Key words: Childhood leukaemia, Power-frequency magnetic field exposure, Extremely low-frequency magnetic field, Housing types

Biomed Environ Sci, 2012; 25(5):597-601

doi: 10.3967/0895-3988.2012.05.015

ISSN:0895-3988

www.besjournal.com(full text)

CN: 11-2816/Q

Copyright ©2012 by China CDC

INTRODUCTION

Childhood leukaemia is a rare disease, and according to data from the Institute of Health Information and Statistics, each year there are about 40-70 new cases per 1.5-1.7 million children (see^[1] for a detailed overview). In CZ during the period 1998-2008, the average number of new cases per year was about 34 per 1 million children, which is slightly lower than the number of new cases in the United Kingdom (approximately 42 cases per 1

million children^[2-3]), and Germany (approximately 45 cases per 1 million children^[3-4]).

Case-control epidemiological studies of the association between childhood leukaemia and power-frequency magnetic field exposure have been conducted in several developed countries (UK, Germany, USA, Canada, Japan, New Zealand, and others). Most of these studies^[5-16] failed to detect a significant association, possibly because of their limited statistical power. After the publication of some pooled analyses^[17-19] the International Agency

[#]Correspondence should be addressed to Vítězslav Jiřík. Tel: 420 597 091 791, E-mail: vitezslav.jirik@email.cz

Biographical note of the first author: Faculty of Medicine, University of Ostrava, Syllabova 19, 703 00 Ostrava - Zábřeh, Czech Republic.

Received: September 2, 2011;

Accepted: December 23, 2011

of Research on Cancer (IARC) classified extremely low frequency magnetic fields (ELF-MF) as being “possibly carcinogenic to humans” (Group 2B in 2002)^[20]. Slightly increased risks of childhood leukaemia have been estimated with an odds ratio (OR) of 2.0 (95% confidence interval [CI]=1.27-3.13) for exposure levels $>0.4 \mu\text{T}^{[18]}$, and of 1.7 (95% CI=1.2-2.3) for exposure levels $>0.3 \mu\text{T}^{[19]}$. Following this classification, other studies attempted to clarify the existence of an association between ELF-MF and childhood leukaemia^[21-25]. A pooled analysis^[26] of these recent studies found a smaller, non-significant OR of 1.44 (95% CI=0.88-2.36) for exposure levels $>0.3 \mu\text{T}$. No attempts have been made to examine ELF-MF and childhood cancer in the Middle or Eastern Europe, or the Czech Republic (CZ), even though the housing stock and related ELF-MF exposures in these areas differ from those in previously studied countries (because of the previous politico-economic system)^[27-28]. We previously identified that average ELF-MF exposure levels were up to three times higher in CZ in comparison with developed Western countries^[29]. For this reason, we conducted a case-control study to elucidate whether ELF-MF exposure in CZ was associated with an increased risk of childhood leukaemia.

METHODS

Equipment and Software

The assessment of exposure to power-frequency magnetic fields was made using purpose-built equipment described elsewhere^[29]. The epidemiological data were processed by the programs STATA (Stata Corp., Release 9, College Station, Texas USA) and OPENEP^[30].

Procedure

A paired case-control study and logistic regression were used to examine the association between ELF-MF and childhood leukaemia. The study was conducted over approximately three years, and the participants were recruited from a range of housing types in the cities, towns and villages of CZ (see^[29]).

Selection of Cases

The cases included children less than 15 years old, living permanently in CZ with leukaemia (predominantly acute lymphoblastic leukaemia) diagnosed by authorized physicians (diagnoses C91-C95^[1]). The time since diagnoses was from several

weeks up to 5 years for a few cases. Recent cases have been preferred. The quality of leukaemia diagnostics in CZ is very good; therefore, the identification of “cases” is reliable and specific. Our study cases ($n=82$) were identified from medical records of university hospitals. The number of cases was limited by the non-participation of some hospitals.

Selection of Controls

Our controls were also identified from the medical records of university hospitals. Just two (Ostrava and Olomouc) of the five university hospitals (Prague, Hradec Kralove, Brno, Ostrava, Olomouc) provided data. The controls had visited a university hospital for reasons other than cancer (for instance, injury, respiratory, or digestive illness). Controls (children without leukaemia) were age- sex- and permanent residence-matched to cases (children with leukaemia). Controls were selected from the same district (for village-based cases) or the same town as their matched case.

In total, 163 participants were recruited, 82 cases and 81 controls. No matches were made for 3 cases and 2 controls. Therefore, only 79 pairs, i.e. 79 cases and 79 controls, were included in the matched analyses.

Exposure Assessment

A detailed description of the ELF-MF exposure assessment method is provided in our previous publication^[29]. We divided all housing types into three categories (high-rise, multifamily and family housing) and measured ELF-MF in the participants’ houses, in the vicinity of the houses, and inside the participants’ schools in different seasons. Each measurement site corresponded with a case or control’s permanent residence; therefore, we were able to evaluate individual exposure. The exposure duration was based on the participant’s age (range 1-14 years).

Epidemiological Data Analysis

The association between ELF-MF exposure and childhood leukaemia occurrence was investigated using conditional logistic regression. We calculated ORs and CIs for various exposure cut points.

RESULTS

The distribution of ELF-MF exposure by case-control pairs, and by case and control groups is presented in Tables 1 and 2.

Table 1. Distribution of ELF-MF Exposure by Case-control Pairs ($n=79$)

Case-control Pairs		Controls					
Exposure	³ 0.2 T	<0.2 T	³ 0.3 T	<0.3 T	³ 0.4 T	<0.4 T	
Cases	³ 0.2 T	17	14	—	—	—	—
	<0.2 T	15	33	—	—	—	—
	³ 0.3 T	—	—	7	10	—	—
	<0.3 T	—	—	13	49	—	—
	³ 0.4 T	—	—	—	—	4	9
	<0.4 T	—	—	—	—	10	56

Table 2. Distribution of ELF-MF Exposure by Case ($n=82$) and Control ($n=81$) Groups

ELF-MF Exposure	Cases			Controls		
	0.2 T	0.3 T	0.4 T	0.2 T	0.3 T	0.4 T
Number of Participants	32	18	14	33	21	15
ELF-MF Exposure	0.2 T	0.3 T	0.4 T	0.2 T	0.3 T	0.4 T
Number of Participants	50	64	68	48	60	66

The logistic regression results are presented in Table 3. Odd ratios were calculated for the matched data (case-control pairs) as well as for the unmatched data. All of the 95% CIs contained 1, implying that there is no association between ELF-MF exposure at any of the examined cut points and childhood leukaemia.

Table 3. Summary of Logistic Regression Analysis Results

Number of Participants	Cut Point	Odds Ratio	95% Confidence Interval
79 Case-control Pairs	0.2 T	0.933	0.451-1.933
	0.3 T	0.769	0.337-1.754
	0.4 T	0.900	0.366-2.215
82 Cases and 81 Controls, Unmatched	0.2 T	0.931	0.474-1.828
	0.3 T	0.805	0.365-1.759
	0.4 T	0.906	0.373 – 2.191

DISCUSSION

Main Finding of this Study

Although higher ELF-MF exposure levels were found in CZ in comparison with developed Western countries^[29], we did not identify an association between exposure to power-frequency magnetic fields and childhood leukaemia. This finding is at odds with other studies^[18-19] that report a

significantly increased risk of childhood leukaemia with higher ELF-MF exposure. However, our results are in agreement with the findings of a recent pooled analysis^[26]. Considering the aetiology of childhood leukaemia^[34-41], and the very important role of infections^[42-45], any contribution of ELF-MF to the most frequent childhood malignant disease is probably minor.

What is Already Known on this Topic

IARC classified ELF-MF as “possibly carcinogenic to humans” in group 2B in 2002^[20] based on some pooled analyses. The true association may actually be weaker than those reported in the earlier pooled analyses^[18-19], and possibly not statistically significant, because the most recent pooled analysis of over ten thousand participants generated an OR of 1.44 and 95% CI of 0.88-2.36^[26].

What this Study Adds

Primarily, the CZ context provided the opportunity to execute a case-control study under higher exposure conditions (Tables 1 and 2, and^[29]). Many previous studies were limited by very low numbers of participants in relatively higher exposures^[12]. This study was carefully designed to decrease unwanted impacts, e.g. comparatively uneven distribution of disease in the population, variation in age and sex, and impacts of some

chemical agents. We successfully reduced possible bias and environmental confounding factors by our paired selection of cases and controls. In addition, this is the first study of ELF-MF and childhood leukaemia in Middle and Eastern Europe, where the characteristics of housing and related ELF-MF exposures are unique^[27-28].

Limitation of This Study

This study was limited by the inclusion of only a small number of participants. This occurred because the number of new diagnoses of childhood leukaemia is low in CZ and some data are not available. Unavailability of data was caused by the refusal of cooperation of several university hospitals. Our use of matched pairs increased the validity of our study, but we did not account for their socioeconomic status and related factors. Nevertheless, our results may be suitable for use in a pooled analysis.

CONCLUSIONS

A low delay between diagnosis and exposure assessment, and the use of matched case-control pairs ensured a higher validity of this study. Although power-frequency magnetic field exposures appear a little higher in CZ in comparison with some Western countries, no indication of any association between exposure levels and childhood leukaemia was found for exposure levels over 0.2 μ T, 0.3 μ T, or 0.4 μ T in our study.

ACKNOWLEDGEMENTS

The authors thank Jaroslav VOLF, Ph.D. (National Institute of Public Health in Prague) for his support; Zdenek SMERHOVSKY, Ph.D. (National Institute of Public Health in Prague) for his help with study design; Bohumir BILAZEK and Hana PTOSZKOVA (Clinic of Pediatrics in University Hospital Ostrava) for their kindness; Lukas JELINEK, Ph.D. (National Institute of Public Health in Prague) for his technical support; Ariana LAJCIKOVA, Ph.D., (National Institute of Public Health in Prague) for her advice; Andrea BABICKOVA, and Marek and Lukas JIRIK for their helpfulness.

REFERENCES

1. ÚZIS - Institute of Health Information and Statistics of the Czech Republic. Novotvary, Česká republika. Available from: <http://www.uzis.cz/katalog/zdravotnicka-statistika/novotvary> [on line, accessed June 30, 2011].
2. Graves M. Childhood leukaemia. *British Medical Journal*, 2002; 324(7332), 283-7.
3. ACCIS-Automated Childhood Cancer Information System. Available from <http://www-dep.iarc.fr/accis/data.htm>, [on line, accessed June 30, 2011].
4. Kaatch P, Spix C, Jung I, Blettner M. Childhood Leukemia in the Vicinity of Nuclear Power Plants in Germany. *Dtsch Arztebl Int*, 2008; 105(42), 725-32.
5. London SJ, Thomas DC, Bowman JD, Sobel E, Cheng TC, Peters JM, et al. Exposure to residential electric and magnetic fields and risk of childhood leukemia. *Am J Epidemiol*, 1991; 134, 923-37.
6. Olsen JH, Nielsen A. & Schulgen G. Residence near high voltage facilities and risk of cancer in children. *Br med J*, 1993; 307, 891-5.
7. Verkasalo PK, Pukkala E, Hongisto MY, et al. Risk of cancer in Finnish children living close to power lines. *Br med J*, 1993; 307, 895-9.
8. Feychting M, Ahlbom A. Magnetic fields and cancer in children residing near Swedish high-voltage power lines. *Am J Epidemiol*, 1993; 138, 467-81.
9. Tynes T, Haldorsen T. Electromagnetic fields and cancer in children residing near Norwegian high-voltage power lines. *Am J Epidemiol*, 1997; 145, 219-26.
10. Linet MS, Hatch EE, Kleinerman RA, et al. Residential exposure to magnetic fields and acute lymphoblastic leukemia in children. *N Engl J Med*, 1997; 337, 1-7.
11. Michaelis J, Schüz J, Meinert R, et al. Combined risk estimates for two German population-based case control studies on residential magnetic fields and childhood acute leukemia. *Epidemiology*, 1998; 9, 92-4.
12. Dockerty JD, Elwood JM, Skegg DC. Electromagnetic fields exposures childhood cancers in New Zealand. *Cancer Causes Control*, 1998; 9, 299-300.
13. UKCCSI - UK Childhood Cancer Study Investigators. Exposure to power-frequency magnetic fields and the risk of childhood cancer. *Lancet*, 1999; 354, 1925-31.
14. Mc Bride ML, Gallagher RP, Theriault G, et al. Power-frequency electric and magnetic fields and risk of childhood leukemia in Canada. *Am J Epidemiol*, 1999; 149, 831-42.
15. Green LM, Miller AB, Agnew DA, et al. Childhood leukemia and personal monitoring of residential exposures to electric and magnetic fields in Ontario, Canada. *Cancer Causes Control*, 1999; 10, 233-43.
16. Auvinen A, Linet MS, Hatch EE, et al. Extremely low-frequency magnetic fields and childhood acute lymphoblastic leukemia: an exploratory analysis of alternative exposure metrics. *Am J Epidemiol*, 2000; 152, 20-31.
17. Angelillo IF, Villari P. Residential exposure to electromagnetic fields and childhood leukaemia: a meta-analysis. *Bull World Health Organ*, 1999; 77, 906-15.
18. Ahlbom A, Day N, Feychting M, et al. A pooled analysis of magnetic fields and childhood leukemia. *Br J Cancer*, 2000; 83, 692-8.
19. Greenland S, Sheppard AR, Kaune WT, et al. A pooled analysis of magnetic fields, wire codes and childhood leukemia. *Epidemiology*, 2000; 11, 624-34.
20. IARC Monographs, Vol.80 (2002): Non-ionizing Radiation, Part 1: Static And Extremely Low-Frequency (ELF) Electric And Magnetic Fields, IARC Press, Lyon, France, ISBN 92 832 1280 0.
21. Li DK, Odouli R, Wi S, et al. A population-based prospective cohort study of personal exposure to magnetic fields during

- pregnancy and the risk of miscarriage. *Epidemiology*, 2002; 13(1), 9-20.
22. Draper G, Vincent T, Kroll ME, et al. Childhood cancer in relation to distance from high voltage power lines in England and Wales: a case-control study. *British Medical Journal*, 2005; 330, 1290.
 23. Kabuto M, Nitta H, Yamamoto S, et al. Childhood leukemia and magnetic fields in Japan: A case-control study of childhood leukemia and residential power-frequency magnetic fields in Japan. *Int J Cancer*, 2006; 119, 643-50.
 24. Mejia-Arangure JM, Fajardo-Gutierrez A, Perez-Saldivar ML, et al. Magnetic fields and acute leukemia in children with Down syndrome. *Epidemiology*, 2007; 18, 158-61.
 25. Kroll ME, Swanson J, Vincent TJ, Draper GJ. Childhood cancer and magnetic fields from high-voltage power lines in England and Wales: a case-control study. *British Journal of Cancer*, 2010; 103 (7), 1122-7.
 26. Kheifets L, Ahlbom A, Crespi CM. Pooled analysis of recent studies on magnetic fields and childhood leukaemia. *British Journal of Cancer*, 2010; 103, 1128-35.
 27. FIHF (Federcasa Italian Housing Federation), Ministry of Infrastructure of the Italian Republic: Housing Statistics in the European Union 2005/2006, pp 56, tab. 2.5. Available at: <http://www.mmr.cz/CMSPages/GetFile.aspx?guid=e99a614d-a fb2-49cf-b80e-f1eb1c5e48d5> [on-line, accessed January 20, 2007].
 28. FIHF (Federcasa Italian Housing Federation), Ministry of Infrastructure of the Italian Republic: Housing Statistics in the European Union 2010, pp 55, tab. 2.5. Available at: <http://www.mmr.cz/CMSPages/GetFile.aspx?guid=a5c4e9a4-4 19a-4e3d-91f4-ada9d2b22016> [on-line, accessed August 15, 2011].
 29. Jirik V, Pekarek L, Janout V. Assessment of Population Exposure to Extremely Low Frequency Magnetic Fields and Its Possible Childhood Health Risk in the Czech Republic. *Indoor Built Environ*, 2011; 20, 362-8.
 30. Dean AG, Sullivan KM, Soe MM. OpenEpi: Open Source Epidemiologic Statistics for Public Health, Version 2.3.1. Available at www.OpenEpi.com, updated 2011/23/06 [on line, accessed 2011/08/29].
 31. Calvente I, Fernandez MF, Villalba J, et al. Exposure to electromagnetic fields (non-ionizing radiation) and its relationship with childhood leukemia: A systematic review. *Science of the Total Environment*, 2010; 408, 3062-9.
 32. Wartenberg D. The Potential Impact of Bias in Studies of Residential Exposure to Magnetic Fields and Childhood Leukemia. *Bioelectromagnetics*, 2001; Supplement 5, S32-47.
 33. Otto M, von Muhlendahl KE. Electromagnetic fields (EMF): Do they play a role in children's environmental health (CEH)? *International Journal Of Hygiene And Environmental Health*, 2007; 210 (5), 635-44.
 34. American Cancer Society: Childhood leukemia. Revised 2011. Available from <http://www.cancer.org/acs/groups/cid/documents/webcontent/003095-pdf.pdf> [on line, accessed June 30, 2011].
 35. Graves M. Childhood leukaemia. *British Medical Journal*, 2002; 324(7332), 283-7.
 36. Stiller CA. Epidemiology and genetics of childhood cancer. *Oncogene*, 2004; 23, 6429-44.
 37. Stewart A, Webb J, Hewitt D. A survey of childhood malignancies. *British Medical Journal*, 1958; 1, 1495-508.
 38. Little J (1999). Epidemiology of childhood cancer. Lyons, International Agency for Research on Cancer, IARC Scientific Publication No 149.
 39. Greaves MF, Wiemels J. Origins of chromosome translocations in childhood leukaemia. *Nat Rev Cancer*, 2003; 3, 639-49.
 40. Hjalgrim LL, Rostgaard K, Hjalgrim H, et al. Birth weight and risk for childhood leukemia in Denmark, Sweden, Norway, and Iceland. *J Nat Cancer Inst*, 2004; 96, 1549-56.
 41. Dickinson HO. The causes of childhood leukaemia. *British Medical Journal*, 2005; 330, 1279-80.
 42. Kinlen LJ. Infection, childhood leukaemia and the Seascale cluster. *Radiol Prot Bull*, 2000; 226, 9-18.
 43. McNally RJQ, Eden TOB. An infectious aetiology for childhood leukaemia: a review of the evidence. *Br J Haematology*, 2004; 127, 243-63.
 44. Gilman C, Peto J, Simpson J, et al. Day care in infancy and risk of childhood acute lymphoblastic leukaemia (ALL): findings from a UK case-control study. *British Medical Journal*, 2005; 330, 1294-7.
 45. Roman E, Simpson J, Ansell P, et al. Childhood Acute Lymphoblastic Leukemia and Infections in the First Year of Life: A Report from the United Kingdom Childhood Cancer Study. *Am J Epidemiol*, 2007; 165, 496-504.