

Original Article



Changes on Stroke Burden Attributable to Ambient Fine Particulate Matter in China*

Jingyu Wang^{1,2,&}, Yan Wang^{1,2,&}, Xiaohua Liang^{3,&}, Keyong Huang¹, Fangchao Liu¹,
Shufeng Chen¹, Xiangfeng Lu^{1,2}, and Jianxin Li^{1,2,#}

1. Department of Epidemiology, Fuwai Hospital, National Center for Cardiovascular Diseases, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100037, China; 2. Key Laboratory of Cardiovascular Epidemiology, Chinese Academy of Medical Sciences, Beijing 100037, China; 3. Department of Clinical Epidemiology and Biostatistics, Children's Hospital of Chongqing Medical University, National Clinical Research Center for Child Health and Disorders, Ministry of Education Key Laboratory of Child Development and Disorders, Chongqing Key Laboratory of Pediatric Metabolism and Inflammatory Disease, Chongqing 400016, China

Abstract

Objective In recent decades, China has implemented a series of policies to address air pollution. We aimed to assess the health effects of these policies on stroke burden attributable to ambient fine particulate matter (PM_{2.5}).

Methods Joinpoint regression was applied to explore the temporal tendency of stroke burden based on data from the Global Burden of Disease 2019 study.

Results The age-standardized rates of disability-adjusted life year (DALY) for stroke attributable to ambient PM_{2.5} in China, increased dramatically during 1990–2012, subsequently decreased at an annual percentage change (APC) of –1.98 [95% confidence interval (CI): –2.26, –1.71] during 2012–2019. For ischemic stroke (IS), the age-standardized DALY rates doubled from 1990 to 2014, and decreased at an APC of –0.83 (95% CI: –1.33, –0.33) during 2014–2019. Intracerebral hemorrhage (ICH) showed a substantial increase in age-standardized DALY rates from 1990 to 2003, followed by declining trends, with APCs of –1.46 (95% CI: –2.74, –0.16) during 2003–2007 and –3.33 (95% CI: –3.61, –3.06) during 2011–2019, respectively. Conversely, the age-standardized DALY rates for subarachnoid hemorrhage (SAH) generally declined during 1990–2019.

Conclusion Our results clarified the dynamic changes of the ambient PM_{2.5}-attributable stroke burden in China during 1990–2019, highlighting the health effects of air quality improvement policies.

Key words: Ambient fine particulate matter; Stroke; Disease burden; Temporal trend; Policy

Biomed Environ Sci, 2024; 37(8): 823-833 doi: [10.3967/bes2024.083](https://doi.org/10.3967/bes2024.083) ISSN: 0895-3988

www.besjournal.com (full text) CN: 11-2816/Q Copyright ©2024 by China CDC

*This research was supported by National Key Research and Development Program of China (2018YFE0115300, 2022YFC3600800, 2017YFC0211706); Chinese Academy of Medical Sciences (CAMS) Innovation Fund for Medical Sciences (2021-I2M-1-010); National Natural Science Foundation of China (82073658, 82070473); National High Level Hospital Clinical Research Funding (2022-GSP-GG-1, 2022-GSP-GG-2); Research Unit of Prospective Cohort of Cardiovascular Diseases and Cancers, CAMS (2019RU038); National Clinical Research Center for Cardiovascular Diseases, Fuwai Hospital, CAMS (NCRC2020006).

[&]These authors contributed equally to this work.

[#]Correspondence should be addressed to Jianxin Li, E-mail: leeljx@126.com, Tel: 86-10-60866572.

Biographical notes of the first authors: Jingyu Wang, male, born in 1998, MD, majoring in cardiovascular epidemiology; Yan Wang, female, born in 1985, MD, majoring in cardiovascular epidemiology; Xiaohua Liang, female, born in 1982, PhD, majoring in etiology and prevention measures of cardiovascular diseases.

INTRODUCTION

Stroke is the third leading cause of disability-adjusted life years (DALYs), and is a major public health issue worldwide^[1]. According to the Global Burden of Disease 2019 (GBD 2019) study, the incident cases, deaths and DALYs of stroke have increased sharply in the past three decades, reaching 12.2 million, 6.55 million and 143 million in 2019, respectively^[2]. The burden of stroke is more pronounced in developing countries such as China^[3]. A remarkable increase of stroke burden occurred in China during 1990–2019. It is estimated that 3.94 million new stroke cases, 13 million patients, 2.19 million deaths, and 45.9 million DALYs occurred in China in 2019^[4-6].

Ambient fine particulate matter with an aerodynamic diameter $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$) has been proved to be one of the leading environmental risk factors for population health in many countries^[7-9], and it has presented a stronger effect on stroke compared to others pollutants, such as PM_{10} and NO_2 ^[10,11]. As the largest developing country, China has experienced severe air pollution in recent decades^[12]. Previous studies in China have showed that a $10 \mu\text{g}/\text{m}^3$ increment in $\text{PM}_{2.5}$ can lead to a 0.34% increase in hospital admission of ischemic stroke for short-term exposure^[13], and a 13% higher risk of incident stroke for long-term exposure^[14]. Air pollution has caused widespread concern in China, and a series of governance policies have been implemented in recent years^[15]. Subsequently, the national $\text{PM}_{2.5}$ concentration in China has gradually decreased since 2011^[16]. Improvements in air quality can lead to significant public health benefits. Understanding the temporal patterns of the $\text{PM}_{2.5}$ -attributable stroke burden is essential for developing tailored strategies for stroke prevention^[17]. Thus, previous studies explored the trends in stroke burden attributable to $\text{PM}_{2.5}$ using the GBD 2019 study^[18,19]. However, they simply focused on the general increasing trends since 1990, without a comprehensive analysis of the change patterns in different periods, especially the trends after the implementation of policies on air quality improvement in recent decades. Moreover, $\text{PM}_{2.5}$ has different effects on stroke subtypes^[14]. Therefore, the dynamic trends in the burden of stroke and its subtypes attributable to ambient $\text{PM}_{2.5}$ in China, remain unclear. There is an urgent need to further evaluate the temporal trends for stroke and its subtypes.

Based on the GBD 2019 study, we aimed to

comprehensively elaborate on the changing patterns in the disease burden of stroke and its subtypes attributable to ambient $\text{PM}_{2.5}$ at different stages across the past three decades in China.

MATERIALS AND METHODS

Data Sources

We obtained age-standardized rates, percents and numbers of DALY, death, year lived with disability (YLD), and year of life lost (YLL) for stroke and its subtypes attributable to ambient $\text{PM}_{2.5}$ in China during 1990–2019 from the GBD 2019 study. Age-standardized rates were computed using the 2019 Global Standard Population ([Supplementary Table S1](#), available in www.besjournal.com)^[20]. DALY indicates health loss from both non-fatal and fatal outcomes, and is calculated as the sum of YLL and YLD^[4]. YLL is the loss of life due to premature death, and is computed as the number of stroke deaths multiplied by the standard remaining life expectancy at the time of death^[4]. YLD is the loss of a healthy life caused by disability, and is calculated using stroke prevalence multiplied by the corresponding disability weights, representing the extent of health loss related to a particular health outcome^[4]. These metrics can help policymakers better understand the disease burden caused by $\text{PM}_{2.5}$, improve air quality, and further contribute to stroke prevention and control. The prevalence estimates of stroke in China were based on systematic reviews of current Chinese researches, using a Bayesian meta-regression tool^[4,21]. Stroke deaths in the GBD 2019 study were estimated using the Cause of Death Ensemble modelling method based on data mainly from surveillance systems, surveys, and the Center for Disease Control and Prevention in China^[4,21].

Definition of Stroke

In the GBD 2019 study, stroke was identified as a rapidly progressing clinical sign of disturbance of cerebral function lasting more than 24 h or resulting in death according to the World Health Organization clinical criteria^[21]. Stroke was classified as ischemic stroke (IS), intracerebral hemorrhage (ICH) and subarachnoid hemorrhage (SAH). IS was defined as an episode of neurological dysfunction triggered by focal cerebral, spinal, or retinal infarction; ICH was diagnosed as a non-traumatic hemorrhagic stroke with focal accumulation of blood in the brain; SAH was identified as a non-traumatic stroke caused by

bleeding into the subarachnoid space in the brain. The International Classification of Diseases codes for stroke and its subtypes are listed in [Supplementary Table S2](#) (available in www.besjournal.com)^[4].

Estimates of Stroke Burden Attributable to Ambient PM_{2.5}

The exposure levels of ambient PM_{2.5} in the GBD 2019 study were estimated by an improved rigorous modelling approach called the Data Integration Model for Air Quality, using multiple data sources, including satellite observations of aerosols in the atmosphere, PM_{2.5} ground measurements, population data and chemical transport model simulations^[7]. Notably, the GBD 2019 study made important changes to revise the relationship between PM_{2.5} and stroke, including recruiting recent Chinese studies on PM_{2.5}, and using flexible splines to fit the risk data^[7]. The ambient PM_{2.5}-attributable disease burden of stroke and its subtypes was evaluated by comparing the distribution of exposure to ambient PM_{2.5} with exposure-risk estimates at each exposure level^[21].

Statistical Analysis

Joinpoint regression was used to analyze the changing patterns of the burden of ambient PM_{2.5}-attributable stroke and its subtypes during 1990–2019. Grid search algorithm was selected to identify the optimal number and position of joinpoints, which were then verified using the Bayesian Information Criterion test. The maximum number of joinpoints in the *Joinpoint* regression model was set to five. The annual percentage change (APC) of each slope segment and the average annual percentage change (AAPC) from 1990 to 2019 showed the average degree of change in stroke burden over a specific period, and their 95% confidence intervals (CI) were calculated based on the t-distribution^[22]. To compare the stroke burden in China with that globally, a z-test was applied^[23]. All statistical tests were 2-sided and $P < 0.05$ was considered as statistically significant. Additionally, we calculated the contributions of YLLs and YLDs to DALYs for stroke attributable to PM_{2.5} in China during 1990–2019. We further described the sex- and age-specific changing patterns of PM_{2.5}-attributable stroke burden.

The GBD Data Tool repository (<http://ghdx.healthdata.org/gbd-results-tool>) was used to obtain the available stroke burden data, and all analyses were performed using *Joinpoint* software, version 4.9.0.1, National Cancer Institute.

RESULTS

Age-Standardized Rates of DALY

The global ambient PM_{2.5}-attributable age-standardized DALY rates for stroke have fluctuated slightly over the past three decades, with values of 348.06, 146.16, 174.82, and 27.08 per 100,000 in 2019 for stroke, IS, ICH, and SAH, respectively ([Supplementary Figure S1](#), available in www.besjournal.com). Notably, China had much higher rates than the global level for most years ([Supplementary Tables S3–S6](#), available in www.besjournal.com). As shown in [Figure 1A](#) and [Table 1](#), the age-standardized DALY rates per 100,000 for stroke in China increased dramatically from 490.54 in 1990 to 727.08 in 2012, thereafter it decreased consistently with an APC of -1.98 (95% CI: $-2.26, -1.71$) during 2012–2019. It is worth noting that a brief decline was observed during 2004–2007, showing an APC of -1.31 (95% CI: $-3.37, 0.79$). Distinct trends were observed for different stroke subtypes. The age-standardized DALY rates per 100,000 for IS climbed from 141.06 to 303.55 during 1990–2014, then persistently decreased with an APC of -0.83 (95% CI: $-1.33, -0.33$) during 2014–2019, demonstrating a noticeable decline in most age groups among people aged 45 years or above ([Figure 2A](#)). For ICH, the DALY rates per 100,000 increased substantially from 273.64 in 1990 to 426.36 in 2003, and presented downward trends subsequently, with APCs of -1.46 (95% CI: $-2.74, -0.16$) during 2003–2007 and -3.33 (95% CI: $-3.61, -3.06$) during 2011–2019, respectively. We observed significant decreases of DALY rates for ICH during 2003–2019 among population aged ≥ 40 years ([Figure 2B](#)). In contrast, the DALY rates for SAH showed a general downward trend over the past three decades, with an AAPC of -2.64 (95% CI: $-2.92, -2.37$). Among the stroke subtypes, ICH had the highest PM_{2.5}-attributable age-standardized DALY rates, followed by IS. However, the gaps between ICH and IS have attenuated in recent decades. The trends in age-standardized DALY rates for stroke and its subtypes related to ambient PM_{2.5} in males were similar to those in females, while males had much higher rates than females ([Figure 1B](#) and [Supplementary Table S7](#), available in www.besjournal.com).

Age-Standardized Percents of DALY Attributable to Ambient PM_{2.5}

As shown in [Supplementary Figure S2](#) and

Supplementary Table S8 (available in www.besjournal.com), the age-standardized percents of stroke burden attributable to ambient $PM_{2.5}$ increased with an AAPC of 2.86 (95% *CI*: 2.69, 3.03) during 1990–2019 in China, with a significant decline occurring after 2014. This phenomenon was also observed for ICH, IS and SAH. We found similar declining trends in both males and females, with males showing a greater decline than females.

Numbers of DALY

The numbers of DALY for stroke attributable to ambient $PM_{2.5}$ in China presented an upward trend over the past three decades, which increased consistently from 4.18 million in 1990 to 12.85 million in 2019, with an AAPC of 3.94 (95% *CI*: 3.61, 4.26) (Figure 1C and Table 1). The similar upward trend occurred in DALYs for IS, with an AAPC of 5.79 (95% *CI*: 5.50, 6.08). For ICH, the DALYs increased

rapidly from 2.38 million in 1990 to 6.48 million in 2011, and then presented an obvious downward trend during 2011–2019, with an APC of -0.48 (95% *CI*: -0.76 , -0.19). For SAH, DALYs fluctuated at a relatively low level over the entire period. Among the stroke subtypes, ICH dominated more than half of the stroke DALYs attributable to $PM_{2.5}$ in China each year from 1990 to 2019. By contrast, contribution of IS on stroke DALYs rose dramatically from 26.89% to 44.66% during 1990–2019, while contribution of SAH declined from 16.16% to 5.49%. We found that males and females shared similar trends in DALYs for stroke and its subtypes, whereas males had much higher DALYs than females (Figure 1D and Supplementary Table S9, available in www.besjournal.com).

Age-Standardized Rates of YLL and YLD

Considering that the DALY comprises YLL and




Figure 1. Age-standardized rates and numbers of DALY for stroke attributable to ambient $PM_{2.5}$ in China from 1990 to 2019. (A) Age-standardized DALY rates in total population; (B) Age-standardized DALY rates in males and females, respectively; (C) Numbers of DALY in total population; (D) Numbers of DALY in males and females, respectively. DALY, disability-adjusted life year; IS, ischemic stroke; ICH, intracerebral hemorrhage; SAH, subarachnoid hemorrhage.

YLD, we further elaborated on their trends, separately (Figure 3). The age-standardized YLL rates for stroke and its subtypes attributable to ambient PM_{2.5} in China were much higher than those in YLD. Age-standardized YLL rates showed similar trends to DALY regardless of stroke subtype and sex. In comparison, the age-standardized YLD rates for stroke and its subtypes showed upward trends,

especially for IS, which sharply increased from 22.34 to 68.92 per 10,000 during 1990–2019. The age-standardized rates of YLD for IS ranked first across the past 30 years, which was even 5-fold and 14-fold higher than those for ICH and SAH in 2019, respectively. Additionally, we found that the age-standardized YLD rates for stroke and its subtypes were higher in females than males except for ICH,

Table 1. The trends in age-standardized rates and numbers of DALY for stroke attributable to ambient PM_{2.5} in China during 1990–2019 using Join-Point regression

Variables	Age-standardized DALY Rates				Numbers of DALY		
	Segments	Period	APC (%; 95% CI)	P value	Period	APC (%; 95% CI)	P value
Stroke	1	1990–1995	2.18 (1.70, 2.66)	< 0.001	1990–1995	5.00 (4.52, 5.47)	< 0.001
	2	1995–2001	3.99 (3.50, 4.48)	< 0.001	1995–2001	7.03 (6.54, 7.51)	< 0.001
	3	2001–2004	1.88 (–0.25, 4.05)	0.079	2001–2004	4.67 (2.57, 6.81)	< 0.001
	4	2004–2007	–1.31 (–3.37, 0.79)	0.199	2004–2007	1.69 (–0.35, 3.77)	0.097
	5	2007–2012	1.08 (0.41, 1.76)	0.004	2007–2012	4.15 (3.48, 4.81)	< 0.001
	6	2012–2019	–1.98 (–2.26, –1.71)	< 0.001	2012–2019	1.12 (0.85, 1.39)	< 0.001
	AAPC (95% CI)	1990–2019	0.94 (0.61, 1.27)	< 0.001	1990–2019	3.94 (3.61, 4.26)	< 0.001
IS	1	1990–1995	3.15 (2.63, 3.67)	< 0.001	1990–1997	6.34 (6.09, 6.59)	< 0.001
	2	1995–2004	4.92 (4.66, 5.18)	< 0.001	1997–2000	9.30 (7.42, 11.22)	< 0.001
	3	2004–2007	–0.16 (–2.39, 2.12)	0.881	2000–2004	7.36 (6.43, 8.30)	< 0.001
	4	2007–2011	3.55 (2.39, 4.73)	< 0.001	2004–2007	3.31 (1.53, 5.12)	0.001
	5	2011–2014	1.58 (–0.69, 3.90)	0.158	2007–2013	6.49 (6.08, 6.91)	< 0.001
	6	2014–2019	–0.83 (–1.33, –0.33)	0.004	2013–2019	2.96 (2.66, 3.27)	< 0.001
	AAPC (95% CI)	1990–2019	2.54 (2.17, 2.91)	< 0.001	1990–2019	5.79 (5.50, 6.08)	< 0.001
ICH	1	1990–1995	1.77 (1.18, 2.37)	< 0.001	1990–1995	4.49 (3.88, 5.10)	< 0.001
	2	1995–2003	4.99 (4.62, 5.36)	< 0.001	1995–2001	8.38 (7.75, 9.02)	< 0.001
	3	2003–2007	–1.46 (–2.74, –0.16)	0.030	2001–2004	5.77 (3.03, 8.58)	< 0.001
	4	2007–2011	–0.13 (–1.43, 1.18)	0.832	2004–2007	1.04 (–1.58, 3.72)	0.410
	5	2011–2019	–3.33 (–3.61, –3.06)	< 0.001	2007–2011	3.04 (1.70, 4.40)	< 0.001
	6	—	—	—	2011–2019	–0.48 (–0.76, –0.19)	0.003
	AAPC (95% CI)	1990–2019	0.49 (0.21, 0.77)	0.001	1990–2019	3.45 (3.02, 3.88)	< 0.001
SAH	1	1990–1996	1.77 (1.42, 2.11)	< 0.001	1990–1996	4.63 (4.27, 4.99)	< 0.001
	2	1996–2000	–3.82 (–4.77, –2.86)	< 0.001	1996–2000	–0.81 (–1.81, 0.21)	0.109
	3	2000–2004	–13.68 (–14.54, –12.82)	< 0.001	2000–2004	–10.77 (–11.67, –9.86)	< 0.001
	4	2004–2007	–6.24 (–8.09, –4.36)	< 0.001	2004–2007	–3.34 (–5.29, –1.36)	0.003
	5	2007–2014	1.88 (1.54, 2.22)	< 0.001	2007–2014	4.56 (4.20, 4.92)	< 0.001
	6	2014–2019	–1.49 (–1.92, –1.05)	< 0.001	2014–2019	0.89 (0.43, 1.35)	0.001
	AAPC (95% CI)	1990–2019	–2.64 (–2.92, –2.37)	< 0.001	1990–2019	0.13 (–0.16, 0.42)	0.379

Note. DALY, disability-adjusted life year; IS, ischemic stroke; ICH, intracerebral hemorrhage; SAH, subarachnoid hemorrhage; APC, annual percent change; AAPC, average annual percent change; CI, confidence interval.

with the greatest difference observed for IS.

Numbers of YLL and YLD

We further evaluated the trends in the numbers of YLL and YLD (Figure 4). The DALYs of stroke arose mostly from the YLLs every year over the entire period. Therefore, the YLLs showed trends similar to those of the DALYs. The YLDs for stroke increased dramatically from 1990 to 2019, particularly for IS, which climbed from 0.20 million to 1.38 million. Moreover, we observed that females had noticeably higher YLDs for IS than males. As for the contribution of YLL to DALY, we noticed that the proportion obviously decreased from 81.89% in 2004 to 76.01% in 2019 for IS, and from 96.09% in 1990 to 86.12% in 2019 for SAH (Supplementary Figure S3, available in www.besjournal.com).

Age Standardized Rates and Numbers of Mortality

We also estimated trends in the age-standardized rates and numbers of mortality for stroke and its subtypes attributable to ambient PM_{2.5}. We found that the mortality of stroke and its subtypes shared similar trends with DALY (Supplementary Figure S4, Supplementary Table S10 and Supplementary Table S11, available in www.besjournal.com).

DISCUSSION

Based on the GBD 2019 study, we found that China has much higher age-standardized DALY rates

for stroke than the global level. A declining trend has been observed for age-standardized DALY rates of stroke in recent years, although there was a general upward trend in China from 1990 to 2019. However, we did not find a downward trend in the numbers of DALY. Among the stroke subtypes, the age-standardized DALY rates for ICH ranked first and then decreased from 2003, thereafter the numbers of DALY declined from 2011. Although the age-standardized DALY rates for IS have decreased since 2014, the downward trend in the ratio of YLL to DALY started as early as 2004. Conversely, the SAH burden showed a successive downward trend over the past three decades. We found that YLLs contributed most of the DALYs for stroke and subtypes attributable to PM_{2.5} throughout the study period. In addition, males had higher DALYs and YLLs for PM_{2.5}-attributable stroke and its subtypes than females. However, females had higher YLDs for stroke, IS and SAH than males did.

The age-standardized rates, percents and numbers of DALY for stroke attributable to PM_{2.5} in China all increased dramatically during 1990–2004, which could be explained by the increasing emission of ambient particulate matter due to rapid economic growth and growing energy consumption over the past decades^[12,24]. Since the early 2000s, China has been actively addressing air pollution and its adverse health effects, implementing policies aimed at improving air quality and reducing the disease burden. A slight decrease in the PM_{2.5}-attributable age-standardized rates of DALY for stroke existed




Figure 2. Average annual percentage change of DALY rates for stroke attributable to ambient PM_{2.5} in China. (A) Average annual percentage change of DALY rates for IS during 2014–2019; (B) Average annual percentage change of DALY rates for ICH during 2003–2019. DALY, disability-adjusted life year; IS, ischemic stroke; ICH, intracerebral hemorrhage.

during 2004–2007, especially for ICH during 2003–2007, which may be related to stricter emission standards of particulate matter for power plants in 2003 and the subsequent implementation of energy-conservation and emission-reduction policies^[15]. In addition, stringent emission controls on regional air quality in preparation for the 2008 Beijing Olympic Games could partially contribute to the slight decrease in stroke burden^[25]. Although no apparent decrease in PM_{2.5} occurred during this period, these policies might have led to changes in the chemical composition and sources of ambient PM_{2.5}, especially the decrease in fossil fuels and industrial sources, and further influenced its pathogenic and lethal effects on stroke^[16,26,27]. The influence of these policies was not strong enough to reduce the DALY rates for IS in the early stages, but it evidently decreased the proportion of YLL to DALY,

implying a weakened lethal effect of PM_{2.5}. Since 2011, China has paid more attention to PM_{2.5}, promulgated brand-new national ambient air quality standard, issued new air quality assessment indicators, optimized the national air monitoring system, promoted the enforcement of the first National Action Plan on Air Pollution Prevention and Control, and developed the air pollution and health effects monitoring system^[15,28–30]. Consequently, population-weighted mean geophysical PM_{2.5} concentrations in China have shown an apparent downward trend since 2011^[16]. This was consistent with the markedly declining trends in the age-standardized rates and percents of DALY for stroke attributable to PM_{2.5} after 2012 and 2014, respectively. It also reduced the numbers of DALY for ICH considerably at the same time, which indicates the effectiveness of the policies for improving air




Figure 3. Age-standardized rates of YLL and YLD for stroke attributable to ambient PM_{2.5} in China from 1990 to 2019. (A) Age-standardized YLL rates in total population; (B) Age-standardized YLL rates in males and females, respectively; (C) Age-standardized YLD rates in total population; (D) Age-standardized YLD rates in males and females, respectively. YLL, year of life lost; YLD, year lived with disability; IS, ischemic stroke; ICH, intracerebral hemorrhage; SAH, subarachnoid hemorrhage.

quality in public health. Additionally, in recent years, policies focusing on desulfurization and denitrification have greatly contributed to reducing the precursors of $PM_{2.5}$, playing an important role in the decline of stroke burden^[31,32]. In summary, our study suggests that remarkable public health benefits can be obtained through policies that control air pollution.

This study identified different changing patterns in the burden of stroke subtypes attributable to ambient $PM_{2.5}$. The declining trend in $PM_{2.5}$ -related age-standardized DALY rates for ICH occurred earlier and more substantially than that for IS. This suggests that the reduction in $PM_{2.5}$, with the implementation of the policies, could yield more health benefits for ICH. This may be explained by the fact that $PM_{2.5}$ is more relevant to ICH than IS^[33]. Additionally, it may be related to the fact that $PM_{2.5}$ could trigger an

increase of blood pressure which has twice the effect on ICH compared to IS, and the fatality rate of ICH is triple that of IS in China^[34-37]. In contrast, the SAH burden attributed to ambient $PM_{2.5}$ remained at a lower level and showed a general downtrend during 1990–2019. This may be more attributable to the substantial improvement in diagnosis and treatment, but less so for $PM_{2.5}$. For example, more non-fatal non-aneurysmal SAH cases have been identified due to improvements in imaging technology, whereas case fatality has decreased because of advances in surgical and medical management^[38].

Our study also showed sex differences in the stroke burden attributable to ambient $PM_{2.5}$. The $PM_{2.5}$ -related stroke burden was much higher in males than females, particularly the fatal stroke burden (YLL). This phenomenon may be explained by




Figure 4. Numbers of YLL and YLD for stroke attributable to ambient $PM_{2.5}$ in China from 1990 to 2019. (A) Numbers of YLL in total population; (B) Numbers of YLL in males and females, respectively; (C) Numbers of YLD in total population; (D) Numbers of YLD in males and females, respectively. YLL, year of life lost; YLD, year lived with disability; IS, ischemic stroke; ICH, intracerebral hemorrhage; SAH, subarachnoid hemorrhage.

the higher probability of ambient PM_{2.5} exposure in males due to a higher proportion of outdoor work^[39,40], and the higher prevalence of risk factors for stroke in males, including hypertension, diabetes, tobacco use, and alcohol consumption^[41-44]. In addition, it may be related to the different pathophysiological functions of cerebral circulation between males and females related to sex hormones^[45]. Therefore, strategies for reducing PM_{2.5}-related stroke burden should be given more attention to males, and provide more protection in high PM_{2.5}-pollution settings. However, it is worth noting that females had higher levels of YLD for stroke than males, particularly for IS, implying females have a greater non-fatal stroke burden. This may be explained by the protective effect of estrogen, which makes females less likely to die from stroke^[45].

This study has some limitations. First, the stroke burden attributed to ambient PM_{2.5} in the GBD 2019 study is not directly observed data, but estimated data generated through mathematical conversion. Therefore, caution should be exercised when interpreting these results. Second, our study is a population-based analysis of trends in stroke burden, which may be inapplicable at the individual level due to potential ecological fallacy and regression dilution bias^[46,47]. Third, the stroke burden at the provincial level in China was not analyzed, because of the lack of relevant data in the GBD 2019 study. Fourth, toxic chemicals bounded to PM_{2.5}, like heavy metals and organic matters, can accumulate in blood and organs, leading to inflammation and further increasing the risk of stroke^[48,49]. Moreover, other determinants, such as ambient temperature, relative humidity, and other pollutants (SO₂, NO₂, CO, O₃ and PM₁₀), also affect stroke^[11,50]. They could confound the association between PM_{2.5} and stroke, which requires further research in the future.

CONCLUSION

Our results illustrate different changing patterns of disease burden for stroke subtypes attributable to ambient PM_{2.5} in China from 1990 to 2019, reflecting the health effects of the policies on improving air quality. Moreover, our results suggest that effective policies should be implemented persistently, and more attention should be paid to males.

ACKNOWLEDGEMENTS

All authors contributed to the interpretation of

results and critically revised the draft. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

CONFLICT OF INTEREST

The authors have no competing interests to declare that are relevant to the content of this article.

AUTHOR CONTRIBUTION

Jingyu Wang: Methodology, Formal analysis, Visualization, Writing-original draft, Writing-review & editing. Yan Wang: Methodology, Software, Writing-original draft, Writing-review & editing. Xiaohua Liang: Methodology, Software, Writing-original draft, Writing-review & editing. Keyong Huang: Writing-review & editing. Fangchao Liu: Funding acquisition, Writing-review & editing. Shufeng Chen: Funding acquisition, Writing-review & editing. Xiangfeng Lu: Funding acquisition, Writing-review & editing. Jianxin Li: Conceptualization, Funding acquisition, Writing-review & editing.

Received: October 27, 2023;

Accepted: April 10, 2024

REFERENCES

1. GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*, 2020; 396, 1204-22.
2. Roth GA, Mensah GA, Johnson CO, et al. Global burden of cardiovascular diseases and risk factors, 1990-2019: update from the GBD 2019 study. *J Am Coll Cardiol*, 2020; 76, 2982-3021.
3. Kim J, Thayabaranathan T, Donnan GA, et al. Global stroke statistics 2019. *Int J Stroke*, 2020; 15, 819-38.
4. Ma QF, Li R, Wang LJ, et al. Temporal trend and attributable risk factors of stroke burden in China, 1990-2019: an analysis for the Global Burden of Disease Study 2019. *Lancet Public Health*, 2021; 6, e897-906.
5. The Writing Committee of the Report on Cardiovascular Health and Diseases in China. Report on cardiovascular health and diseases in China 2021: an updated summary. *Biomed Environ Sci*, 2022; 35, 573-603.
6. The Writing Committee of the Report on Cardiovascular Health and Diseases in China. Report on cardiovascular health and diseases in China 2022: an updated summary. *Biomed Environ Sci*, 2023; 36, 669-701.
7. GBD 2019 Risk Factors Collaborators. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*, 2020; 396, 1223-49.
8. Di Q, Wang Y, Zanobetti A, et al. Air pollution and mortality in

- the medicare population. *N Engl J Med*, 2017; 376, 2513–22.
9. Al Ahad MA, Demšar U, Sullivan F, et al. Long-term exposure to air pollution and mortality in Scotland: A register-based individual-level longitudinal study. *Environ Res*, 2023; 238, 117223.
 10. Tian F, Cai M, Li HT, et al. Air pollution associated with incident stroke, poststroke cardiovascular events, and death: a trajectory analysis of a prospective cohort. *Neurology*, 2022; 99, e2474–84.
 11. Shah ASV, Lee KK, McAllister DA, et al. Short term exposure to air pollution and stroke: systematic review and meta-analysis. *BMJ*, 2015; 350, h1295.
 12. Li FZ, Liu Y, Lü JJ, et al. Ambient air pollution in China poses a multifaceted health threat to outdoor physical activity. *J Epidemiol Community Health*, 2015; 69, 201–4.
 13. Tian YH, Liu H, Zhao ZL, et al. Association between ambient air pollution and daily hospital admissions for ischemic stroke: A nationwide time-series analysis. *PLoS Med*, 2018; 15, e1002668.
 14. Huang KY, Liang FC, Yang XL, et al. Long term exposure to ambient fine particulate matter and incidence of stroke: prospective cohort study from the China-PAR project. *BMJ*, 2019; 367, l6720.
 15. Jin YN, Andersson H, Zhang SQ. Air pollution control policies in China: a retrospective and prospects. *Int J Environ Res Public Health*, 2016; 13, 1219.
 16. Hammer MS, van Donkelaar A, Li C, et al. Global estimates and long-term trends of fine particulate matter concentrations (1998–2018). *Environ Sci Technol*, 2020; 54, 7879–90.
 17. Huang C, Moran AE, Coxson PG, et al. Potential cardiovascular and total mortality benefits of air pollution control in urban China. *Circulation*, 2017; 136, 1575–84.
 18. Sang SW, Chu C, Zhang TC, et al. The global burden of disease attributable to ambient fine particulate matter in 204 countries and territories, 1990–2019: A systematic analysis of the Global Burden of Disease Study 2019. *Ecotoxicol Environ Saf*, 2022; 238, 113588.
 19. Chen HJ, Zhou ZH, Li ZL, et al. Time trends in the burden of stroke and subtypes attributable to PM_{2.5} in China from 1990 to 2019. *Front Public Health*, 2022; 10, 1026870.
 20. GBD 2019 Demographics Collaborators. Global age-sex-specific fertility, mortality, healthy life expectancy (HALE), and population estimates in 204 countries and territories, 1950–2019: a comprehensive demographic analysis for the Global Burden of Disease Study 2019. *Lancet*, 2020; 396, 1160–203.
 21. GBD 2019 Stroke Collaborators. Global, regional, and national burden of stroke and its risk factors, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet Neurol*, 2021; 20, 795–820.
 22. Kim HJ, Fay MP, Feuer EJ, et al. Permutation tests for joinpoint regression with applications to cancer rates. *Stat Med*, 2000; 19, 335–51.
 23. Altman DG, Bland JM. Interaction revisited: the difference between two estimates. *BMJ*, 2003; 326, 219.
 24. Ding S, Wei ZW, He JH, et al. Estimates of PM_{2.5} concentrations spatiotemporal evolution across China considering aerosol components in the context of the Reform and Opening-up. *J Environ Manage*, 2022; 322, 115983.
 25. Chen YY, Jin GZ, Kumar N, et al. The promise of Beijing: Evaluating the impact of the 2008 Olympic Games on air quality. *J Environ Econ Manage*, 2013; 66, 424–43.
 26. Martinelli N, Olivieri O, Girelli D. Air particulate matter and cardiovascular disease: a narrative review. *Eur J Intern Med*, 2013; 24, 295–302.
 27. Zhu YH, Huang L, Li JY, et al. Sources of particulate matter in China: Insights from source apportionment studies published in 1987–2017. *Environ Int*, 2018; 115, 343–57.
 28. Liu J, Han YQ, Tang X, et al. Estimating adult mortality attributable to PM_{2.5} exposure in China with assimilated PM_{2.5} concentrations based on a ground monitoring network. *Sci Total Environ*, 2016; 568, 1253–62.
 29. Chen Z, Wang JN, Ma GX, et al. China tackles the health effects of air pollution. *Lancet*, 2013; 382, 1959–60.
 30. Guo H, Cheng TH, Gu XF, et al. Assessment of PM_{2.5} concentrations and exposure throughout China using ground observations. *Sci Total Environ*, 2017; 601–602, 1024–30.
 31. Wang SW, Zhang Q, Martin RV, et al. Satellite measurements oversee China's sulfur dioxide emission reductions from coal-fired power plants. *Environ Res Lett*, 2015; 10, 114015.
 32. Zhai SX, Jacob DJ, Wang X, et al. Fine particulate matter (PM_{2.5}) trends in China, 2013–2018: Separating contributions from anthropogenic emissions and meteorology. *Atmos Chem Phys*, 2019; 19, 11031–41.
 33. Lin HL, Tao J, Du YD, et al. Differentiating the effects of characteristics of PM pollution on mortality from ischemic and hemorrhagic strokes. *Int J Hyg Environ Health*, 2016; 219, 204–11.
 34. Lacey B, Lewington S, Clarke R, et al. Age-specific association between blood pressure and vascular and non-vascular chronic diseases in 0.5 million adults in China: a prospective cohort study. *Lancet Glob Health*, 2018; 6, e641–9.
 35. Feng SL, Gao D, Liao F, et al. The health effects of ambient PM_{2.5} and potential mechanisms. *Ecotoxicol Environ Saf*, 2016; 128, 67–74.
 36. Bai L, Chen H, Hatzopoulou M, et al. Exposure to ambient ultrafine particles and nitrogen dioxide and incident hypertension and diabetes. *Epidemiology*, 2018; 29, 323–32.
 37. Tu WJ, Chao BH, Ma L, et al. Case-fatality, disability and recurrence rates after first-ever stroke: A study from bigdata observatory platform for stroke of China. *Brain Res Bull*, 2021; 175, 130–5.
 38. Mackey J, Khoury JC, Alwell K, et al. Stable incidence but declining case-fatality rates of subarachnoid hemorrhage in a population. *Neurology*, 2016; 87, 2192–7.
 39. Shen Y, Zhang XD, Chen C, et al. The relationship between ambient temperature and acute respiratory and cardiovascular diseases in Shenyang, China. *Environ Sci Pollut Res Int*, 2021; 28, 20058–71.
 40. Yang DY, Xu CD, Wang JF, et al. Spatiotemporal epidemic characteristics and risk factor analysis of malaria in Yunnan Province, China. *BMC Public Health*, 2017; 17, 66.
 41. Wang ZW, Chen Z, Zhang LF, et al. Status of hypertension in China: results from the China hypertension survey, 2012–2015. *Circulation*, 2018; 137, 2344–56.
 42. Zhang M, Yang L, Wang LM, et al. Trends in smoking prevalence in urban and rural China, 2007 to 2018: Findings from 5 consecutive nationally representative cross-sectional surveys. *PLoS Med*, 2022; 19, e1004064.
 43. Millwood IY, Walters RG, Mei XW, et al. Conventional and genetic evidence on alcohol and vascular disease aetiology: a prospective study of 500 000 men and women in China. *Lancet*, 2019; 393, 1831–42.
 44. Wang YJ, Li ZX, Gu HQ, et al. China Stroke Statistics: an update on the 2019 report from the National Center for Healthcare Quality Management in Neurological Diseases, China National Clinical Research Center for Neurological Diseases, the Chinese Stroke Association, National Center for Chronic and Non-communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention and Institute for Global Neuroscience and Stroke Collaborations. *Stroke Vasc Neurol*, 2022; 7, 415–50.
 45. Haast RAM, Gustafson DR, Kiliaan AJ. Sex differences in stroke.

- [J Cereb Blood Flow Metab](#), 2012; 32, 2100–7.
46. Wang ZK, Hu SB, Sang SP, et al. Age-period-cohort analysis of stroke mortality in China: data from the global burden of disease study 2013. [Stroke](#), 2017; 48, 271–5.
47. Guo YM, Zeng HM, Zheng RS, et al. The association between lung cancer incidence and ambient air pollution in China: A spatiotemporal analysis. [Environ Res](#), 2016; 144, 60–5.
48. Li QZ, Liu HB, Alattar M, et al. The preferential accumulation of heavy metals in different tissues following frequent respiratory exposure to PM_{2.5} in rats. [Sci Rep](#), 2015; 5, 16936.
49. Lu H, Wang RH, Li JJH, et al. Long-term exposure to the components of fine particulate matters and disability after stroke: Findings from the China National Stroke Screening Surveys. [J Hazard Mater](#), 2023; 460, 132244.
50. Tian YH, Liu H, Si YQ, et al. Association between temperature variability and daily hospital admissions for cause-specific cardiovascular disease in urban China: A national time-series study. [PLoS Med](#), 2019; 16, e1002738.