

Letter



Associations of Dietary Behaviors and Physical Activity with Anxiety and Depressive Symptoms among Chinese Adults Aged 18–60 Years: A Population-based Cross-sectional Study

Jia Fu^{1,&}, Shige Qi^{2,&}, Xiaobing Liu³, Yanli Xue⁴, Yu Zhang^{1,#}, and Haoyan Guo^{5,#}

This population-based cross-sectional study aimed to examine the associations of unhealthy diet and insufficient physical activity (PA) with anxiety and depressive symptoms among Chinese adults aged 18–60 years. Data were derived from the 2022–2023 Psychology and Behavior Investigation of Chinese Residents (PBICR) Project, a large cross-sectional survey conducted in China. After exclusions, 14,358 adults were included in this study. Dietary risk factors were summed into a 0–5 score. PA was measured by the International Physical Activity Questionnaire (IPAQ-7, short form). Anxiety and depressive symptoms were defined as the Generalized Anxiety Disorder Scale (GAD-7) ≥ 10 and Patient Health Questionnaire-9 (PHQ-9) ≥ 10 , respectively. Adjusted odds ratios (aORs) and 95% confidence intervals (CIs) were estimated using ordered logistic regression; dose-response was examined with restricted cubic splines. Overall, 1,879 participants (13.09%) self-reported anxiety and 3,084 (21.48%) self-reported depressive symptoms. A monotonic dose-response relationship was observed between the number of unhealthy dietary behaviors and both outcomes. Compared with participants reporting no unhealthy dietary behaviors, those reporting 1, 2, 3, 4, and 5 behaviors had progressively higher odds of anxiety (aOR = 1.47, 2.03, 2.75, 4.59, and 8.37) and depression (aOR = 1.53, 2.14, 2.84, 5.59, and 16.46). Physical activity level was not significantly associated with either anxiety or depressive symptoms after adjustment for covariates. The strong relationship between cumulative unhealthy dietary behaviors and mental health symptoms highlights the potential importance

of dietary interventions for preventing anxiety and depression.

Key words: Dietary behaviour; Physical activity; Anxiety; Depression; Dose-response; Chinese adults

Mental health disorders, particularly anxiety and depression, have emerged as leading public health challenges worldwide owing to their high prevalence, disabling consequences, and association with an increased risk of dementia and suicide^[1]. Accumulating evidence highlights that unhealthy dietary patterns, such as frequent consumption of sugary drinks, skipping breakfast, and excessive intake of processed foods, are linked to metabolic disturbances and adverse psychological outcomes. By contrast, regular physical activity promotes endorphin and dopamine release, alleviates emotional distress, and improves mood regulation^[2]. Existing research in this area on Chinese populations has focused mainly on particular groups of people, such as adolescents^[3] and older adults^[4,5], and few studies have investigated the associations between dietary behavior, physical activity intensity, and anxiety or depressive symptoms in Chinese adults. Nonetheless, adults are better able to take charge of their own lives, exhibit more self-control, and are better suited to directly change their lifestyles, such as nutritional status and physical exercise level. A better understanding of the role of these lifestyle factors in the onset and progression of adult depressive symptoms is necessary, especially given their variable nature and their potential for use in the prevention and treatment of anxiety and depressive symptoms. Therefore, this study investigated the associations of dietary behaviors

doi: [10.3967/bes2026.054](https://doi.org/10.3967/bes2026.054)

1. National Institute for Viral Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 100050, China; 2. National Center for Chronic and Noncommunicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 100050, China; 3. National Institute for Nutrition and Health, Chinese Center for Disease Control and Prevention, Beijing 100050, China; 4. School of Biomedical Engineering, Capital Medical University, Beijing 100069, China; 5. Guangzhou Women and Children's Medical Center, Guangzhou 510623, Guangdong, China

and physical activity intensity with anxiety and depressive symptoms among Chinese adults aged 18–60, using data from the PBICR program.

This cross-sectional study used data from the Psychology and Behavior Investigation of Chinese Residents (PBICR) Project conducted from 2022–2023, using a multistage, stratified random sampling method. The survey covered 15 provinces, 148 cities, 202 districts and counties, 390 townships/towns/streets, and 780 communities/villages across China. From a starting national sample of 21,916 individuals, the analysis was restricted to participants aged 18–60 years. Following the exclusion of cases with age out of range, logical errors, or implausible physical activity data, the final sample consisted of 14,358 adults.

Anxiety and depressive symptoms were assessed using two validated self-report instruments: the Generalized Anxiety Disorder Scale (GAD-7) and Patient Health Questionnaire-9 (PHQ-9). Anxiety and depressive symptoms were dichotomized at the clinically significant threshold of ≥ 10 for prevalence estimation and as the dependent variable in binary logistic regression models. Additionally, to examine dose-response relationships across the full spectrum of symptom severity, the ordinal categories (minimal/mild/moderate/severe) were used as the outcome in ordered logistic regression models. Detailed scoring criteria are provided in Supplementary Table S1. Unhealthy dietary behaviors were identified based on the Dietary Guidelines for Chinese Residents (2022), encompassing five domains: frequent consumption of sugary drinks, alcohol use, skipping breakfast, takeaway consumption, and dieting/fasting practices (Supplementary Table S2). Physical activity intensity was assessed using the International Physical Activity Questionnaire (IPAQ-7, short form) and categorized into low, moderate, and high levels according to MET minutes per week.

Descriptive statistics are presented as means and standard deviations for continuous variables and as frequencies and percentages for categorical variables. Chi-squared tests were used in order to compare anxiety and depressive symptoms across lifestyle categories. Ordered logistic regression models were employed to estimate the adjusted odds ratios (aORs) and 95% confidence intervals (CIs) for the associations between dietary behaviors, physical activity intensity, and mental health outcomes. After adjusting for variables such as age, gender, marital status, urban and rural status, education status, career status, residential

status, insurance status and per capita monthly household income, two models were constructed: Model I: Included the total number of unhealthy dietary behaviors and physical activity intensity. Model II: Further incorporated specific types of unhealthy dietary behaviors and physical activity intensity. Restricted cubic spline analyses were applied in order to examine potential dose-response relationships between the number of unhealthy dietary behaviors and the prevalence of anxiety and depressive symptoms. All analyses were two-sided, with a significance level of $P < 0.05$. Statistical analyses were conducted using SAS 9.4 (SAS Institute Inc., Cary, NC, USA) and R 4.2.2 (drc package).

Among the 14,358 adults included, anxiety was self-reported by 1,879 participants (13.09%) and depressive symptoms were self-reported by 3,084 participants (21.48%). There were significant differences seen among the different groups in terms of age, sex, education status, region, career status, monthly household income per capita, marital status, living alone, and health insurance ($P < 0.05$) (Supplementary Tables S3–S4). We also examined the distribution of unhealthy dietary behaviors and physical activity intensity in patients with anxiety and depression. A total of 15.93%, 12.52%, and 13.08% of participants with low, moderate, and high physical activity levels, respectively, suffered from anxiety. A total of 19.41%, 15.14%, 23.31%, 16.94%, and 23.86% of people with preference for sugary beverages, being drunk, skipping breakfast, takeaways, and on a diet, suffer from anxiety. Individuals who engaged in low, moderate, and high levels of physical activity experienced depressive symptoms at a rate of 24.55%, 21.21%, and 21.04%, respectively. And individuals who preferred for sugary beverages, being drunk, skipping breakfast, takeaways, and on a diet exhibited depressive symptom rates of 30.47%, 25.17%, 34.77%, 26.49%, and 35.71%, respectively (Table 1).

In Model I, compared with those reporting no unhealthy dietary behaviors, individuals with a greater number of unhealthy dietary behaviors had higher odds of developing anxiety and depressive symptoms, suggesting a dose-response relationship between dietary risk accumulation and adverse mental health outcomes (Figure 1). In Model II, the types of unhealthy eating behaviors were positively correlated with anxiety and depression symptoms, indicating that specific dietary patterns may contribute differentially to symptom burden.

However, physical activity showed no significant association with either anxiety or depressive symptoms in both Model I and Model II, after adjusting for demographic and lifestyle covariates (Table 2). Subgroup analyses stratified according to sex and age confirmed the robustness of these associations (Table 3).

The prevalence of self-reported depressive symptoms (21.48%) and anxiety prevalence (13.09%) in our study is consistent with recent large-scale surveys among Chinese adults, which have reported rates ranging from 16.3% to 24.9% using comparable screening instruments^[6]. These concordances support the representativeness of our sample despite its cross-sectional design.

Diet has a significant effect on mental health. According to the current study, eating poorly increases anxiety and depressive symptoms. There is a noticeable detrimental effect on people's experiences of anxiety and depressive symptoms when the number of unhealthy eating habits reaches three or more. Furthermore, individuals who preferred for sugary beverages ($OR = 1.40$, 95% CI 1.29–1.52) and being drunk ($OR = 1.32$, 95% CI 1.22–1.43) had an impact on anxiety and depressive symptoms in relation to the type of diet. Similar to our results, Oliván et al. demonstrated that consuming multiple carbonated or sugar-filled beverages daily is indicative of increased depressive symptoms^[7]. Daily intake of carbonated or soft

Table 1. Distribution of poor dietary behaviors and physical activity intensity in patients with anxiety and depressive

Variables	Anxiety					Depressive				
	Overall	Mild (%)	Moderate (%)	Severe (%)	P value	Overall	Mild (%)	Moderate (%)	Severe (%)	P value
Poor dietary behaviors										
Preference for sugary beverages	19.41	36.34	14.82	4.59	< 0.0001	30.47	37.92	17.83	12.64	< 0.0001
Being drunk	15.14	36.59	11.74	3.39	< 0.0001	25.17	37.74	15.52	9.65	< 0.0001
Skipping breakfast	23.31	34.94	17.79	5.51	< 0.0001	34.77	34.68	18.96	15.81	< 0.0001
Preference for takeaway	16.94	36.74	11.82	5.13	< 0.0001	26.49	38.73	15.30	11.19	< 0.0001
On a diet	23.86	40.65	16.42	7.44	< 0.0001	35.71	40.50	18.41	17.30	< 0.0001
Physical activity level										
					0.005					< 0.0001
Low	15.93	31.86	11.45	4.48		24.55	33.72	13.72	10.83	
Mid	12.52	33.72	9.85	2.67		21.21	37.31	13.41	7.80	
High	13.08	33.25	10.11	2.97		21.04	35.49	13.52	7.52	




Figure 1. Dose-response relationship between the number of unhealthy dietary behaviors and the prevalence of anxiety and depressive.

Table 2. Results of the ordered logistic regression of poor dietary behaviors and physical activity intensity with anxiety and depressive

Variables	Anxiety		Depressive	
	Model I	Model II	Model I	Model II
No. of poor dietary behavior				
0	1.00		1.00	
1	1.47 (1.36–1.58)		1.53 (1.42–1.64)	
2	2.03 (1.85–2.24)		2.14 (1.95–2.35)	
3	2.75 (2.38–3.18)		2.84 (2.46–3.27)	
4	4.59 (3.42–6.17)		5.59 (4.19–7.47)	
5	8.37 (3.59–19.55)		16.46 (6.64–43.09)	
Poor dietary behaviors				
Preference for sugary beverages		1.40 (1.29–1.52)		1.52 (1.40–1.64)
Being drunk		1.32 (1.22–1.43)		1.34 (1.24–1.44)
Skipping breakfast		1.65 (1.51–1.80)		1.73 (1.59–1.89)
Preference for takeaway		1.15 (1.06–1.26)		1.13 (1.04–1.22)
On diet		1.97 (1.77–2.19)		2.02 (1.82–2.25)
Physical activity level				
Mid	1.00	1.00	1.00	1.00
Low	1.06 (0.95–1.19)	1.07 (0.95–1.19)	1.02 (0.92–1.13)	1.02 (0.92–1.14)
High	1.04 (0.97–1.11)	1.04 (0.98–1.12)	0.97 (0.90–1.03)	0.97 (0.91–1.04)

Note. Adjusted variables: age, gender, marital status, urban and rural status, education, career status, residential status, insurance status, and per capita monthly household income; Model I: effect of the number of poor eating behaviors versus physical activity on anxiety and depression; Model II: effect of poor eating behaviors versus physical activity on anxiety and depression.

Table 3. Associations of unhealthy dietary behavior and physical activity level with depressive and anxiety symptoms according to age and gender

Dependent variables	Independent variables	No. of unhealthy dietary behavior						Physical activity level			
		0	1	2	3	4	5	Mid	Low	High	
Anxiety	Age (years)	18–34	1.00 (1.34–1.66)	1.49 (1.78–2.28)	2.01 (2.21–3.10)	2.61 (3.21–6.15)	4.44 (4.68–29.16)	11.64 (0.23–15.14)	1.00 (0.90–1.21)	1.05 (0.90–1.08)	0.99 (0.90–1.08)
		35–44	1.00 (1.25–1.76)	1.48 (1.97–3.09)	2.47 (1.99–4.34)	2.94 (1.92–9.36)	4.24 (0.23–15.14)	2.02	1.00 (0.84–1.41)	1.09 (0.99–1.35)	1.16
		45–60	1.00 (1.27–1.67)	1.46 (1.46–2.26)	1.82 (2.50–6.12)	3.91 (1.72–51.26)	9.29	–	1.00 (0.86–1.37)	1.09 (0.94–1.22)	1.07
	Gender	Males	1.00 (1.30–1.63)	1.46 (1.81–2.36)	2.07 (2.36–3.46)	2.86 (2.93–6.37)	4.32 (2.68–19.94)	7.28	1.00 (0.89–1.26)	1.06 (1.00–1.22)	1.10
		Females	1.00 (1.36–1.68)	1.51 (1.75–2.32)	2.01 (2.05–3.22)	2.57 (3.14–7.85)	4.97 (2.22–55.90)	10.99	1.00 (0.92–1.23)	1.06 (0.88–1.07)	0.97
		18–34	1.00 (1.42–1.74)	1.57 (1.91–2.43)	2.16 (2.45–3.41)	2.89 (4.22–8.03)	5.82 (7.63–61.99)	20.72	1.00 (0.88–1.17)	1.01 (0.84–1.00)	0.92
Depressive	Age (years)	35–44	1.00 (1.25–1.74)	1.48 (1.92–2.96)	2.38 (1.61–3.43)	2.35 (1.99–8.80)	4.19 (0.60–68.27)	6.55	1.00 (0.78–1.28)	1.00 (1.00–1.34)	1.16
		45–60	1.00 (1.32–1.72)	1.51 (1.57–2.39)	1.94 (2.30–5.68)	3.62 (1.89–63.75)	10.71	–	1.00 (0.85–1.32)	1.06 (0.84–1.07)	0.95
		Males	1.00 (1.38–1.70)	1.53 (1.96–2.53)	2.23 (2.37–3.45)	2.86 (3.68–7.79)	5.35 (6.23–62.84)	18.72	1.00 (0.87–1.22)	1.03 (0.91–1.09)	0.99
	Gender	Females	1.00 (1.41–1.72)	1.56 (1.79–2.36)	2.06 (2.25–3.50)	2.81 (3.71–9.15)	5.81 (2.31–62.81)	11.42	1.00 (0.88–1.17)	1.02 (0.85–1.02)	0.93
		18–34	1.00 (1.42–1.74)	1.57 (1.91–2.43)	2.16 (2.45–3.41)	2.89 (4.22–8.03)	5.82 (7.63–61.99)	20.72	1.00 (0.88–1.17)	1.01 (0.84–1.00)	0.92
		35–44	1.00 (1.25–1.74)	1.48 (1.92–2.96)	2.38 (1.61–3.43)	2.35 (1.99–8.80)	4.19 (0.60–68.27)	6.55	1.00 (0.78–1.28)	1.00 (1.00–1.34)	1.16

drinks, including sugar, has been linked to an increased level of anxiety, according to a randomized controlled experiment. Additionally, daily consumption of carbonated or soft drinks containing sugar and non-nutritive sweeteners has been linked to increased depressive symptoms^[8]. Claims regarding alcohol consumption have been controversial; however, a 2018 study published in *The Lancet* showed that the safest amount of alcohol to consume is 0 mL, with no drops recommended^[9]. Although drinking alcohol and sugary drinks can stimulate hormone secretion in the body and produce excitement to a certain extent, as a dietary behavior, prolonged intake will still have an impact on mental health, and dependence on addiction will, in contrast, cause psychological stress. Furthermore, consuming excessive alcohol and sweet drinks can harm one's organs and increase the risk of heart disease, stroke, and other conditions that can worsen anxiety and depressive symptoms^[10].

The results of this study revealed that poor dietary behavior was a risk factor for anxiety and depressive symptoms. A cross-sectional study demonstrated that people who rarely ate breakfast were most likely to have a profile of psychological issues ($OR = 3.59$; 95% CI : 2.52–5.11)^[11]. Zhang et al. reported that the odds ratios and 95% confidence intervals for depressive symptoms were 1.35 (1.08–1.68) and 1.40 (1.07–1.83) when breakfast consumers were compared to occasional consumers and skippers, respectively^[12]. Yun et al. reported a correlation between depressive symptoms and the frequency of eating breakfast and ate out^[13]. Zhang et al. assessed 3337 commuters in total and discovered a strong correlation between eating out frequently and a greater risk of mental symptoms, particularly when dining out for lunch and supper. Individuals who ate at home were least likely to experience mental symptoms, followed by those who ate in the canteen at work^[14].

In contrast to the robust dietary effects, our study found no significant association between PA level and anxiety or depressive symptoms after adjustment for covariates. This null finding appears to differ from the prevailing consensus that regular PA protects against depression and anxiety. Numerous systematic reviews and meta-analyses have shown that regular physical activity protects against depression and anxiety. For instance, early cross-sectional studies demonstrated that individuals with moderate-to-high levels of physical activity exhibited substantially lower anxiety and depression symptom scores than their sedentary

counterparts^[15]. Subsequently, Baumeister et al. provided further evidence linking domain-specific physical activity (both exercise and leisure time) with reduced rates of anxiety and depressive symptoms^[16]. Notably, a prospective cohort study of adults identified a linear inverse relationship between 4–7 hours/week of moderate-to-vigorous physical activity and incident depression, whereas > 7 hours/week conferred no additional benefit, suggesting a ceiling effect or optimal dose range rather than unlimited protection^[17]. Guo et al. also showed a nonlinear dose-response relationship between leisure-time physical activity (LTPA) and depression risk; low and moderate doses were protective against incident depression, whereas high doses were associated with increased risk^[18].

Several explanations may account for the null findings in our study. First, dietary behaviors and physical activity are often correlated lifestyle factors; individuals with healthier diets may also engage in more physical activity. When both are included in the same model, the variance in mental health outcomes explained by these correlated behaviors may be predominantly captured by the dietary risk score, which showed a stronger and more graded association with anxiety and depression than PA level. This statistical collinearity may have reduced the independent contribution of PA to the model. Second, the cross-sectional design precludes determination of temporal sequence; any protective effect of regular PA may require longer exposure durations than captured by a single-point measurement to manifest detectable mental health benefits.

This study has several limitations, including its cross-sectional design, self-reported data, and semi-quantitative dietary assessment. Despite these limitations, our findings highlight the strong dose-response relationship between cumulative unhealthy dietary behaviors and mental health symptoms, underscoring the potential importance of dietary interventions. The null findings for physical activity warrant further longitudinal research to confirm causality and explore potential effect modifiers such as PA type, intensity, and context.

Competing Interests The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Ethics This study was approved by the Institutional Review Committee of Ji'nan University, Guangzhou, China (JNUKY-2021-018). All the participants fully

understood the study and voluntarily signed informed consent forms.

Authors' Contributions Data curation, Formal analysis, Visualization: Jia Fu; Writing – original draft: Shige Qi; Conceptualization, Writing – review & editing: Xiaobing Liu; Writing – review & editing: Yanli Xue, Yu Zhang and Haoyan Guo. These authors contributed equally to this work.

Acknowledgements We thank all the investigators of the PBICR program and all the subjects who participated in this survey.

Data Sharing The data that support the findings of this study are available from the corresponding author upon reasonable request. The supplementary materials will be available in www.besjournal.com.

[&]These authors contributed equally to this work.

[#]Correspondence should be addressed to Yu Zhang, PhD, E-mail: zhangyu@ninh.chinacdc.cn; Haoyan Guo, PhD, E-mail: guohy@chinacdc.cn

Biographical notes of the first authors: Jia Fu, MM, majoring in medicinal chemistry, E-mail: fujia0922@163.com; Shige Qi, MM, majoring in epidemiology and health statistics, E-mail: qishige@ncncd.chinacdc.cn

Received: December 30, 2025;

Accepted: April 29, 2026

REFERENCES

- Stafford J, Chung WT, Sommerlad A, et al. Psychiatric disorders and risk of subsequent dementia: systematic review and meta-analysis of longitudinal studies. *Int J Geriatr Psychiatry*, 2022; 37, 1-22.
- Liu LY, Guo CF, Lang FF, et al. Association of breakfast, total diet quality, and mental health in adolescents: a cross-sectional study of HBSC in Greece. *Eur J Pediatr*, 2023; 182, 5385-97.
- Cao RL, Gao TT, Ren H, et al. Unique and cumulative effects of lifestyle-related behaviors on depressive symptoms among Chinese adolescents. *Int J Soc Psychiatry*, 2022; 68, 354-64.
- Shao P, Xu Y, Pan CW. Factors associated with and prevalence of depressive features amongst older adults in an urban city in eastern China. *S Afr J Psychiatry*, 2017; 23, 1064.
- Tsai AC, Chi SH, Wang JY. Cross-sectional and longitudinal associations of lifestyle factors with depressive symptoms in ≥ 53-year old Taiwanese — results of an 8-year cohort study. *Prev Med*, 2013; 57, 92-7.
- Jiang BC, Zhang J, Yang M, et al. Prevalence and risk factors of depressive and anxiety symptoms and functional constipation among university students in Eastern China. *World J Psychiatry*, 2025; 15, 106451.
- Oliván-Blázquez B, Aguilar-Latorre A, Motrico E, et al. The relationship between adherence to the Mediterranean diet, intake of specific foods and depression in an adult population (45-75 years) in primary health care. A cross-sectional descriptive study. *Nutrients*, 2021; 13, 2724.
- Castro A, Gili M, Visser M, et al. Soft drinks and symptoms of depression and anxiety in overweight subjects: a longitudinal analysis of an European cohort. *Nutrients*, 2023; 15, 3865.
- GBD 2016 Alcohol Collaborators. Alcohol use and burden for 195 countries and territories, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*, 2018; 392, 1015-35.
- Millwood IY, Walters RG, Mei XW, et al. Conventional and genetic evidence on alcohol and vascular disease aetiology: a prospective study of 500000 men and women in China. *Lancet*, 2019; 393, 1831-42.
- Haghighatdoost F, Feizi A, Esmailzadeh A, et al. Breakfast skipping alone and in interaction with inflammatory based quality of diet increases the risk of higher scores of psychological problems profile in a large sample of Iranian adults. *J Nutr Sci*, 2021; 10, e10.
- Zhang LY, Yin JJ, Cai XF, et al. Association between eating behaviors and depressive symptoms in Chinese adults: a population-based cross-sectional study. *Psychol, Health Med*, 2022; 27, 1176-83.
- Yun H, Kim DW, Lee EJ, et al. Analysis of the effects of nutrient intake and dietary habits on depression in Korean adults. *Nutrients*, 2021; 13, 1360.
- Zhang L, Xie YY, Li BX, et al. Psychiatric symptoms and frequency of eating out among commuters in Beijing: a bidirectional association?. *Nutrients*, 2022; 14, 4221.
- Brunes A, Augestad LB, Gudmundsdottir SL. Personality, physical activity, and symptoms of anxiety and depression: the HUNT study. *Soc Psychiatry Psychiatr Epidemiol*, 2013; 48, 745-56.
- Baumeister SE, Leitzmann MF, Bahls M, et al. Associations of leisure-time and occupational physical activity and cardiorespiratory fitness with incident and recurrent major depressive disorder, depressive symptoms, and incident anxiety in a general population. *J Clin Psychiatry*, 2017; 78, e41-7.
- Harvey SB, Øverland S, Hatch SL, et al. Exercise and the prevention of depression: Results of the HUNT cohort study. *Am J Psychiatry*, 2018; 175, 28-36.
- Guo ZG, Li R, Lu ST. Leisure-time physical activity and risk of depression: a dose-response meta-analysis of prospective cohort studies. *Medicine*, 2022; 101, e29917.