Original Articles

Occupational Hazard Factors And The Trajectory Of Fasting Blood Glucose Changes In Chinese Male Steelworkers Based On Environmental Risk Scores: A Prospective Cohort Study



Mingxia Zou¹, Wei Du¹, Qin Kang¹, Yuhao Xia¹, Nuoyun Zhang¹, Liu Feng¹, Feiyue Li¹, Tiancheng Ma¹, Yajing Bao¹, and Hongmin Fan^{1,2,3,#}

1. Department of Epidemiology, School of Public Health, North China University of Science and Technology, Tangshan 063210, Hebei, China; 2. Hebei Key Laboratory of Occupational Health and Safety for Coal Industry, Tangshan 063210, Hebei, China; 3. Hebei Coordinated Innovation Centre of Occupational Health and Safety, Tangshan 063210, Hebei, China

Abstract

Objective We aimed to investigate the patterns of fasting blood glucose (FBG) trajectories and analyze the relationship between various occupational hazard factors and FBG trajectories in male steelworkers.

Methods The study cohort included 3,728 workers who met the selection criteria for the Tanggang Occupational Cohort between 2017 and 2022. A group-based trajectory model was used to identify the FBG trajectories. Environmental risk scores (ERS) were constructed using regression coefficients from the occupational hazard model as weights. Univariate and multivariate logistic regression analyses were performed to explore the effects of occupational hazard factors using the ERS on FBG trajectories.

Results FBG trajectories were categorized into three groups. An association was observed between high temperature, noise exposure, and FBG trajectory (P < 0.05). Using the first quartile group of ERS1 as a reference, the fourth quartile group of ERS1 had an increased risk of medium and high FBG by 1.90 and 2.21 times, respectively (odds ratio [OR] = 1.90, 95% confidence interval [CI]: 1.17–3.10; OR = 2.21, 95% CI: 1.09–4.45).

Conclusion An association was observed between occupational hazards based on ERS and FBG trajectories. The risk of FBG trajectory levels increase with an increase in ERS.

Key words: Fasting blood glucose; Occupational hazards; Group-based trajectory modeling; Environmental risk scores; Steelworkers

Biomed Environ Sci, 2025; 38(6): 673-684	doi: 10.3967/bes2025.058	ISSN: 0895-3988
www.besjournal.com (full text)	CN: 11-2816/Q	Copyright ©2025 by China CDC

INTRODUCTION

asting blood glucose (FBG) levels are widely used in clinical practice to detect type 2 diabetes mellitus (T2DM) and prediabetes. Recent studies have shown that FBG trajectories are associated with diseases such as hypertension, cardiovascular disease (CVD), and cancer^[1-3]. Factors influencing FBG levels are complex and are closely related to family history, lifestyle, and environmental factors. Previous studies mainly focused on lifestyle patterns^[4,5]. However, environmental factors, particularly occupational hazards, also affect FBG levels. Several studies have shown that occupational

[#]Correspondence should be addressed to Hongmin Fan, MD, Tel: 86-15030574016, E-mail: hongminfan1004 @ncst.edu.cn

Biographical note of the first author: Mingxia Zou, Master's Degree, majoring in public health and preventive medicine, E-mail: zoumingxia11@163.com

hazards, such as high temperature, noise, and dust exposure, are related to changes in FBG levels^[6-9]. However, studies exploring the relationships between occupational risk factors, combined occupational exposure, and changes in FBG levels remain limited.

Several methods have been applied to analyze combinations of occupational hazards, such as environmental risk scores (ERS), Bayesian kernel machine regression (BKMR), and weighted quartile sum regression (WQS)^[10,11]. Although BKMR and WQS can solve high-dimensionality and collinearity problems, the incorporation of high-dimensional data with high hardware requirements remains a constantly evolving topic in statistical research. In WQS, converting data into quartiles can result in data loss^[10,12]. In comparison, ERS is a simple tool for assessing the risk of exposure to multiple harmful factors and can integrate the disease risk of exposure to multiple pollutant mixtures; this easyto-operate tool supports new research results and addresses the need for better risk stratification from a single pollutant to a multiple pollutant framework^[11,13].

Hence, a cohort of occupational hazards and blood glucose changing trajectories from 2017 to 2022 in steelworkers was built to analyze the relationship between single exposure and a combination of occupational hazard factors and long-term changes in FBG levels. In addition, we evaluated the dose-response relationship between cumulative exposure to occupational hazard factors and the changing trajectory of FBG levels. Our study provides a basis for blood glucose level control and the prevention of adverse outcomes in Chinese male steelworkers.

METHODS

Study Design and Population

A cohort of occupational hazard factors and blood glucose levels was constructed using data from the Tang Gang Occupational Cohort (TGOC). The TGOC is a registered occupational population health effects longitudinal cohort (ChiCTR2400083306). A total of 7,636 workers aged 18–60 years who were eligible for occupational health examinations were recruited from 13 departments (hot rolling, iron making, cold rolling, hot rolling, and steelmaking) at Tangshan Iron and Steel Group Co., Ltd. from February 2017 to June 2017. The main occupational hazards in factories include high temperatures, noise, dust, and carbon monoxide (CO). The five-wave follow-up was completed between 2018 and 2022. The exclusion criteria were as follows: 1) FBG follow-up of less than four times; 2) missing baseline occupational hazard data; and 3) female sex. Female workers were excluded as they mainly engage in auxiliary work at Tanggang and are rarely exposed to occupational hazard factors. Finally, 3,728 male workers who underwent at least five physical examinations between 2017 and 2022 were included to analyze the relationship between occupational hazard factors and FBG trajectory. Based on the sample size calculation formula for the cohort study, each group (occupational hazard exposure and nonexposure) required 587 participants. Considering a 10% loss to follow-up rate, 646 participants were included in each group. In total, 3,728 workers met the sample size requirements.

Questionnaire

unified structured questionnaire Α was completed by well-trained investigators through a one-on-one survey. The questionnaire collected data on demographic information (age, nationality, monthly family income per capita, education level, and marital status), behavioral habits (smoking, drinking, dietary habits, and physical activity [PA]), personal and family disease history (diabetes and hypertension), medication history, occupational history (factory category, factory entry date, work categories, and shift work status), and use of protective equipment (usage of dustproof masks, noise-proof earplugs, and cooling drinks). PA and dietary habit data were collected using the International Physical Activity Questionnaire (IPAQ) and Food Frequency Questionnaire (FFQ), which included 15 items.

Physical Examination and Laboratory Test

Height and weight were measured and recorded by a professional nurse, using a height and weight measurement instrument (HGM301, China). Participants were required to remain standing while wearing light clothing and no shoes. The precision of height and weight was 0.1 cm and 0.1 kg respectively [body mass index (BMI) = weight (kg) / height² (m²)].

Fasting venous blood (5 mL) was collected after 6–8 h of overnight fasting and placed in anticoagulant tubes. FBG levels were measured using the glucose oxidase method in the same laboratory using an automatic blood chemical analyzer (Hitachi 7600-020, Japan).

On-site Hygiene Investigation

Data on occupational high temperatures, noise, dust, and CO exposure were obtained from annual monitoring reports of occupational hazard factors of companies^[14]. Temperature measurements were performed at multiple points in the workplace using a wet-bulb black globe temperature gauge (WBGT) procedures^[15]. according to standard Hightemperature exposure was defined according to the GBZ/T229.3-2010 Classification of Occupational Hazards at Workplaces Part 3: Occupational Exposure to Heat Stress^[16]. Noise was measured using the HS6288B Noise Spectrum Analyzer, and noise exposure was determined according to the GBZ/T229.4-2010 Classification of Occupational Hazards at Workplaces Part 4: Occupational Exposure to Noise^[17,18]. Dust was measured using a Dust Sampler with a sampling time of 45 min at each sampling point, and dust exposure was determined according to GBZ-T192.1-2007 Determination of Dust in the Air of Workplace Part 1: Total Dust Concentration^[19]. CO was measured using a portable CO infrared analyzer (Model GXH-3011A), and CO exposure was determined according to the GBZ2.1-2007 Occupational Exposure Limits for Hazardous Agents in the Workplace Part 1: Chemical Hazardous Agents^[20,21].

Cumulative exposure to occupational hazards was measured using a Job Exposure Matrix for steelworkers. Cumulative exposure measurements (CEM) for occupational hazards were calculated using the following formula:

$$CEM = \sum L = L_1 T_1 + L_2 T_2 + \dots + L_n T_n$$
 (1)

where: \boldsymbol{L}_n is the exposure of the hazard factor at \boldsymbol{T}_n time.

Definition and Grouping of Indicators

Nationality was divided into Han and other groups. Marital status included unmarried, married, and other (divorced, widowed, or remarried). Educational levels were divided into three groups: middle school and below; high school and secondary specialty schools; and college and above. Smoking and alcohol consumption were defined according to World Health Organization^[14]. According to the IPAQ classification standard^[22], the tensity of PA is classified as low, medium, and high. Product-line factories included hot rolling, cold rolling, and ironmaking, while the others were auxiliary

factories. Owing to the product process reforms in 1998 and 2007, the factory entry era was divided into three groups (before 1998, 1998–2007, and after 2007). Shift work was defined as a system of irregular working hours in which one or more teams perform tasks continuously for 24 h by working in shifts^[14]. The FFQ dietary score ranges from 15 to 60 points, based on the plant-based diet index (PDI)^[23]; higher PDI scores indicate healthier plant-based foods and better dietary quality. The diagnostic criteria for diabetes and hypertension were as follows: 1) pre-existing diagnosis of hypertension or diabetes by second-level hospital diagnosis; and 2) taking medication for hypertension or diabetes.

Statistical Analysis

Means and standard deviations (SD) were used to describe continuous data with a normal distribution, whereas numbers with percentages (%) were used for categorical data. Comparisons between groups were performed using the chisquare or Fisher's exact test. Univariate repeatedmeasures analysis of variance was performed to evaluate changes in FBG levels over time. Missing values for monthly family income per capita and marital status were determined by using multiple interpolations.

Group-based trajectory modeling (GBTM) was used to identify the different patterns of FBG trajectories in the TGOC from 2017 to 2022. The optimal number of patterns in the trajectories was determined based on the following criteria^[24,25]: (1) the absolute value of the Bayesian information criterion (BIC) of the model was closest to 0; (2) the average posterior probability (AvePP) of each group was \geq 0.7; and (3) the sample size of each group was > 5% of the total participants. The shape of the trajectory for each group was determined by a stepwise procedure that started with all groups set to a cubic order, and the change in the BIC and significance of the parameters were then compared.

The ERS assesses the risk of exposure to multiple hazard factors in epidemiologic studies^[11,26]. The ERS calculation procedure was as follows: (1) the relationships between cumulative occupational hazard exposure and FBG trajectory were analyzed using nonconditional logistic regression with adjusted age, nationality, education level, monthly family income per capita, work category, shift work, BMI, PA, smoking, drinking, PDI, usage of dustproof masks, usage of noise-proof earplugs, usage of cooling drinks, personal and family disease history, and medication history. (2) ERS was calculated by

summing up using a regression coefficient (weight) of 10 times and grouped according to ERS quintiles. ERS1 calculates the weight using 10 times the regression coefficient of each hazard factor as a single hazard factor, whereas ERS2 calculates the weight using 10 times the regression coefficient from a multiple-hazard-factor model that includes all the hazard factors simultaneously.

Multinomial logistic regression models were used to compare the effects of occupational hazard factors and ERS on FBG trajectories in 3,728 participants. Model 1 was unadjusted and Model 2 was adjusted for age, nationality, education level, monthly family income per capita, factory entry date, work categories, shift work, usage of dustproof masks, usage of cooling drinks, usage of noise-proof earplugs, personal and family disease history, medication history, BMI, PA, smoking, drinking, and PDI scores. Odds ratios (ORs) and 95% confidence intervals (CIs) were calculated to assess the risk associated with exposure to each occupational hazard and ERS for different FBG trajectories. The chi-square trend test was used to analyze the relationship between cumulative exposure to occupational hazards and ERS and FBG levels. Finally, the association between occupational hazards and FBG trajectories was further analyzed by stratification and subgroups based on factory category, factory entry date, age, education level, shift work, and BMI. The CEM and PDI scores were divided into groups according to quantity and tertile, respectively. Additionally, we conducted a sensitivity

analysis using the same method after excluding patients with diabetes.

Statistical analysis was performed using R 4.4.1 and SPSS 23.0, and a statistical difference of P < 0.05 was considered statistically significant.

RESULTS

A total of 3,728 male workers were included in this study (Figure 1), with an average age of 42 years (range, 23–57 years); 61.5% were aged > 40 years. A total of 90% of the participants were of Han nationality and married. 56.2% had attended high school or secondary specialized school, and 55.9% had worked for \geq 20 years. The proportions of participants exposed to noise, dust, high temperature, and CO were 56.6%, 55.6%, 61.6%, and 51.8%, respectively. The study population was married and educated, had a higher monthly family income per capita, and was younger than the population lost to follow-up (Supplementary Table S1).

The Trajectory of FBG

The average FBG level from 2017 to 2022 was 6.03 \pm 1.1 mmol/L, 6.04 \pm 1.2 mmol/L, 5.77 \pm 1.1 mmol/L, 5.8 6 \pm 1.2 mmol/L, 5.34 \pm 1.4 mmol/L, and 5.36 \pm 1.5 mmol/L respectively. A statistically significant change was noted in FBG levels over time (F = 377.530, *P* < 0.001). Based on the GBTM, three different trajectories were identified according to the BIC, sample size proportion, and AvePP (Table S2



Figure 1. Flow chart of the study population.

and Figure 2). FBG of around 5.8 mmol/L at five-time points was defined as the "low-level group" (trajectory 1: n = 3,312, 88.7%). FBG of around 8 mmol/L at five-time points was defined as the "medium-level group" (trajectory 2: n = 215, 5.9%). FBG of about 12 mmol/L at five-time points was



Figure 2. Trajectories in FBG.

defined as the "high-level group" (trajectory 3: n = 201, 5.4%).

Baseline Characteristics Of Different FBG Trajectory Groups

There were significant differences in the factory category, age, education level, factory entry date, shift work, diabetes, hypertension, medication history, family history of diabetes, and BMI among the three FBG trajectories. The proportion of participants with diabetes, hypertension, obesity, and medication use in Trajectory 3 was higher than those in Trajectories 1 and 2 (Table 1).

Association with Occupational Hazards and FBG Trajectories

Figure 3A and Table S3 show that participants exposed to high temperatures were prone to

Variables	Low-level group (n = 3,312)	Medium-level group (n = 215)	High-level group (n = 201)	χ²	P-value
Product-line factories, <i>n</i> (%)	1,645 (49.7)	107 (49.8)	76 (37.8)	10.709	0.005
Age (years), n (%)				72.399	< 0.001
< 30	263 (7.9)	5 (2.3)	5 (2.5)		
30-	1,073 (32.4)	51 (23.7)	40 (19.9)		
40-	1,464 (44.2)	121 (56.3)	89 (44.3)		
≥ 50	512 (15.5)	38 (17.7)	67 (33.3)		
Han, n (%)	3,222 (97.3)	209 (97.2)	195 (97.0)	0.054	0.974
married, n (%)	2,987 (90.2)	194 (90.2)	186 (92.5)	6.916	0.140
education level, n (%)				35.974	< 0.001
middle school or below	628 (19.0)	72 (33.5)	54 (26.9)		
high school or secondary specialized school	1,891 (57.1)	110 (51.2)	111 (55.2)		
college or above	793 (23.9)	33 (15.3)	36 (17.9)		
monthly family income per capita (¥), n (%)				1.731	0.785
< 2000	1,686 (50.9)	102 (47.4)	105 (52.3)		
2000–2999	1,119 (33.8)	74 (34.4)	66 (32.8)		
≥ 3000	507 (15.3)	39 (18.2)	30 (14.9)		
entry factory date, n (%)				28.111	< 0.001
before 1998	1,988 (60.0)	148 (68.8)	154 (76.6)		
1998–2007	957 (28.9)	52 (24.2)	35 (17.4)		
after 2007	367 (11.1)	15 (7.0)	12 (6.0)		
work categories, n (%)				28.615	0.432
ironworker	126 (3.8)	9 (4.2)	11 (5.5)		
steelworker	241 (7.3)	18 (8.4)	11 (5.5)		
others	2,945 (88.9)	188 (87.4)	179 (89.0)		

Table 1. Baseline characteristics according to FBG trajectory groups

Variables	Low-level group	Medium-level group	High-level group	χ²	P-value
Years of service, n (%)	((1) 220)	(,, 201)	28.519	< 0.001
1-	128 (3.9)	3 (1.4)	4 (2.0)		
5-	263 (7.9)	12 (5.6)	9 (4.5)		
10-	1,118 (33.8)	59 (27.4)	48 (23.9)		
≥ 20	1,803 (54.4)	141 (65.6)	140 (69.6)		
shift work, n (%)				10.038	0.040
never	542 (16.4)	30 (14.0)	32 (15.9)		
ever	483 (14.6)	33 (15.3)	45 (22.4)		
present	2,287 (69.0)	152 (70.7)	124 (61.7)		
diabetes, n (%)	0 (0.0)	13 (6.0)	102 (50.7)	1,639.004	< 0.001
hypertension, n (%)	416 (12.6)	34 (15.8)	57 (28.4)	41.201	< 0.001
medication history, n (%)	0 (0.0)	13 (6.0)	27 (13.4)	375.310	< 0.001
family history of diabetes, n (%)	346 (10.4)	37 (17.2)	50 (24.9)	45.387	< 0.001
smoking, n (%)	1,989 (60.1)	132 (61.4)	128 (63.7)	1.150	0.563
drinking, n (%)	2,506 (75.7)	173 (80.5)	161 (80.1)	4.363	0.113
PDI, <i>n</i> (%)				1.941	0.747
low	1,095 (33.0)	71 (33.0)	76 (37.8)		
medium	1,108 (33.5)	72 (33.5)	63 (31.3)		
high	1,109 (33.5)	72 (33.5)	62 (30.9)		
PA, n (%)				3.755	0.440
low	329(9.9)	28 (13.0)	20 (10.0)		
medium	486(14.7)	30 (14.0)	23 (11.4)		
high	2,497(75.4)	157 (73.0)	158 (78.6)		
BMI (kg/m²), <i>n</i> (%)				37.082	< 0.001
< 24	1,176 (35.5)	48 (22.4)	41 (20.4)		
24-	1,448 (43.7)	102 (47.4)	104 (51.7)		
≥ 28	688 (20.8)	65 (30.2)	56 (27.9)		
usage of dustproof masks, <i>n</i> (%)	2,398 (72.4)	163 (75.8)	142 (70.6)	1.546	0.462
usage of cooling drinks, n (%)	2,722 (82.2)	181 (84.2)	166 (82.6)	0.565	0.754
usage of noise-proof earplugs, n (%)	1,588 (47.9)	111 (51.6)	95 (47.3)	1.159	0.560

Continued

Note. PDI, the plant-based diet index; PA, physical activity; BMI, body mass index.

elevated FBG trajectories; however, no significant relationship was observed between occupational noise, dust, CO exposure, and FBG trajectories. The FBG trajectory level increased with increasing cumulative high-temperature exposure ($P_{trend} < 0.001$) (Supplementary Table S7).

Compared to participants not exposed to noise, the risk of a median FBG trajectory with noise exposure decreased before adjusting for age and other factors. However, no relationship was found between occupational noise and the median FBG trajectory after adjusting for age and other factors. Furthermore, no statistically significant differences were noted between occupational exposure and high FBG trajectories, regardless of whether other factors were unadjusted or adjusted (P > 0.05) (Figure 3B and Supplementary Table S4). The risk of a median or high FBG trajectory increased with cumulative dust or CO exposure (both P = 0.009) (Supplementary Table S8).



Figure 3. Association with occupational hazards and FBG trajectories by Logistic regression analysis. (A) compares the medium- and low-level groups, and (B) compares the high- and low-level groups. Model 1 was unadjusted; Model 2 was adjusted for age, nationality, education level, monthly family income per capita, factory entry date, work categories, shift work, usage of dustproof masks, usage of cooling drinks, usage of noise-proof earplugs, personal and family disease history, medication history, BMI, PA, smoking, drinking, and PDI scores.

Stratified Analysis of Occupational Hazards and FBG Trajectories

The association between occupational hazards and FBG trajectories was inconsistent across factory categories, factory entry date, age, education level, shift work, and BMI. Working in a product-line factory, the entry factory dates after 1998, being aged \geq 40 years, being in the ever-shift group, and high-temperature exposure can increase the risk of median or high FBG trajectories. Working in a product-line factory, factory entry dates after 1998, educated to junior high school and below, and noise exposure decreased the risk of median or high FBG levels (Supplementary Tables S9–S20).

Associations with ERS and FBG Trajectories

The ERS1 and ERS2 were constructed using single occupational hazards and FBG trajectories using logistic regression analysis (Table 2). The median ERS1 and ERS2 scores in the medium- and low-level were 5.3 (range, 0-11.8), and 5.0 (range, 0-14.7), respectively. In the high-versus low-level groups, the median ERS1 and ERS2 were 3.9 (range, 0-10.5), and 4.2 (range, 0-10.3), respectively.

The FBG trajectory levels increased with

Variables	Medium-level group ^a		High-level group ^a	
	ERS1	ERS2	ERS1	ERS2
cumulative noise exposure, [dB(A)·year]				
never exposed	0.0	0.0	0.0	0.0
33.64–	2.3	1.3	0.8	0.8
90.70-	2.2	1.9	0.6	0.3
93.56-	0.3	0.1	2.8	2.0
96.25–111.76	3.2	4.0	0.7	0.6
cumulative dust exposure, [mg/m ³ ·year]				
never exposed	0.0	0.0	0.0	0.0
59.22-	0.8	0.1	2.1	1.4
138.70-	0.9	0.4	1.7	2.6
271.30-	0.3	1.7	2.2	1.0
456.82–1,464.48	0.4	1.1	2.6	1.8
cumulative high-temperature exposure, [°C·year]				
never exposed	0.0	0.0	0.0	0.0
374.44-	2.0	1.7	1.0	1.8
951.20-	5.0	4.8	0.3	0.6
2,316.34–	1.5	2.2	2.3	3.6
3,590.92–7,608.06	6.2	7.0	2.8	3.7
cumulative CO exposure, [mg/m ³ ·year]				
never exposed	0.0	0.0	0.0	0.0
76.76-	1.2	0.2	1.5	0.6
277.72-	0.6	0.8	0.4	1.1
544.88-	1.3	2.0	2.3	2.0
888.50–2,684.10	2.0	1.6	0.1	1.6

Note. ERS1 was constructed using ten times the regression coefficients of occupational hazards in the single occupational hazard model as weights, and ERS2 was constructed using ten times the regression coefficients of occupational hazards in the multiple occupational hazard model as weights. ^a represents a low–level of FBG group as a reference.

increasing ERS1 and ERS2 levels (both P_{trend} < 0.05) (Supplementary Table S7). Figure 4A and Supplementary Table S5 show that the risk of developing medium-level FBG in the highest ERS1 and ERS2 was irrespectively 2.17 times (1.39–3.40) and 2.06 times (1.31–3.24) with comparison to that in the lowest group in Model 1. After adjusting for factors such as age, BMI, drinking, and smoking, the risk of developing medium-level FBG decreased from 2.17 (1.39–3.40) to 1.90 (1.17–3.10) in the highest ERS1 group, and from 2.02 (1.28–3.18) to 1.77 (1.07–2.95) in the ERS2 group with scores of 6.1–8.4.

The risk of elevated FBG levels did not increase with an increase in ERS (Supplementary Table S8). In Model 2, Figure 4B and Supplementary Table S6 demonstrate that the risk of high-level FBG in the highest ERS1 score and ERS2 of 4.7–6.7 score is 2.21 times (1.09–4.45) and 2.59 times (1.25–5.33)

compared to that in the lowest group.

Sensitivity Analysis

After excluding participants with confirmed diabetes, the analysis showed stable results (Supplementary Table S21–S22). However, after adjusting for covariates, the high-level group is more likely to have a high-temperature exposure compared to the low-level group (P = 0.023, OR = 1.85, 95% *CI*: 1.09–3.14), whereas there was no association before the exclusion.

DISCUSSION

Our study analyzed the longitudinal FBG changing trajectories from 2017 to 2022 among male steelworkers in the TGOC region. The FBG trajectories of the steelworkers were categorized



Figure 4. The relationship between ERS and FBG trajectories by Logistics regression analysis. (A) compares the medium-level and low-level groups, and (B) compares the high-level and low-level groups. Model 1 was unadjusted; Model 2 was adjusted for age, nationality, education level, monthly family income per capita, factory entry date, work categories, shift work, usage of dustproof masks, usage of cooling drinks, usage of noise-proof earplugs, personal and family disease history, medication history, BMI, PA, smoking, drinking, and PDI scores.

into three groups. Single and multiple occupational hazards are related to the FBG trajectory, particularly high temperatures and noise, which play important roles in FBG trajectories. In addition, our findings may help identify high-risk individuals with high FBG levels and provide measures to minimize the increase in FBG levels.

FBG levels increased with cumulative hightemperature exposure in our study. These results are consistent with those of previous studies^[7,27,28]. A possible explanation for this observation is that high temperatures are unfavorable for heat dissipation in the body. As the temperature rises, the body's temperature regulation ability weakens, leading to fluid loss and resulting in pathological blood concentration, which, in turn, increases FBG levels. This increase is harmful to glucose metabolism and tends to gradually increase the degree of damage^[7,29,30].

Our study showed that occupational noise exposure can decrease FBG trajectory. Our findings contradict those of previous cohort studies showing a positive correlation between noise exposure and high FBG^[6,31]. A possible explanation for this finding is that there may have been a more pronounced healthy worker effect in the study population. The reference group consisted of workers who were not exposed to hazard factors, whereas the exposed group had stricter pre-employment or on-the-job health examination requirements. Those who do not meet the occupational health examination eligibility criteria are restricted from engaging in work that involves noise exposure. Therefore, individuals who worked with occupational risk factors for a long time are more likely to have better physical health or have already adapted to these factors.

Compared to the low-level group, the other two groups had higher ERS. Essentially, the larger the ERS, the higher the FBG level. Moreover, the combined effect of these occupational hazards was greater than the single-factor effects. Many crosssectional studies have demonstrated that exposure to multiple occupational hazards increases the risk of diseases such as diabetes and hypertension^[7,32,33]. A 2019 review indicated that exposure to multiple occupational hazards can have significant adverse effects on health^[34]. However, the underlying mechanism remains unclear. One possible explanation is that simultaneous exposure to multiple occupational hazards may significantly affect hypoglycemic mechanisms in the body. Exposure to multiple occupational hazards not only affects the concentration and activity of pancreatic beta cells but also leads to abnormal excitation of the sympathetic nervous system, which can negatively affect blood glucose regulation and result in elevated blood glucose levels^[7,33,35].

The association between occupational hazards and FBG trajectories was inconsistent across factory categories, factory entry date, age, education level, shift work, and BMI. Workers in productline factories are more likely to be exposed to occupational hazards than those in auxiliary factories. Shift work and obesity have been confirmed as risk factors for T2DM and can affect FBG levels. Shift work disrupts the body's circadian rhythm or triggers metabolic disorders that increase the risk of T2DM^[36,37] and may also alter blood glucose levels by decreasing insulin sensitivity and reducing glucose tolerance^[38].

This study had several strengths. First, we utilized a prospective cohort with stable populations, detailed information on various occupational exposures, and a comprehensive questionnaire survey for potential confounders such as diet and PA. Second, we used the ERS to evaluate single and combined exposure to multiple occupational hazard factors, as well as the effects of occupational exposure on FBG trajectories. However, this study had some limitations. First, the ERS only considered the additive effect of various occupational hazard factors, neglecting the antagonistic and multiplicative interaction effects between them. These interactions can be further analyzed by applying the BKMR and other methods. Second, the healthy worker effect may have led to underestimation of the relationship between exposure and outcomes. Third, considering the influence of drugs on FBG levels, we only adjusted for medication history. However, after excluding patients with diabetes, the results of the repeated analyses remained stable. Finally, when the blood glucose trajectories were divided into five groups, the upward and downward trajectories were observed. Owing to the population sample size limitations, our study selected only three groups of trajectories. It is recommended that the sample size be expanded in future studies to include more trajectory group analyses.

CONCLUSION

Our study revealed an association between high temperature, noise exposure, and FBG levels. As ERS increases, the risk of elevated FBG levels also increases. Our study provides a basis for blood glucose control and prevention among male steelworkers.

Funding This study was supported by the Key Research and Development Program of the Ministry of Science and Technology of China (grant number: 2016YF0900605); the Key Research and Development Program of Hebei Province (grant number: 192777129D); the Joint Fund for Iron and Steel of the Natural Science Foundation of Hebei Province (grant number: H2016209058); and the National Natural Science Foundation for Regional Joint Fund of China (grant number: U22A20364).

Competing Interests The authors declare no conflict of interest.

Ethics This study was approved by the Medical Ethics Committee of North China University of Science and Technology (approval number: 16040). All participants who completed the baseline and follow-up surveys signed an informed consent form.

Authors' Contributions Design: Hongmin Fan; Methodology: Mingxia Zou; Investigation: Mingxia Zou, Wei Du, Qin Kang, Yuhao Xia, Nuoyun Zhang, Liu Feng, Feiyue Li, Tiancheng Ma, and Yajing Bao; Project administration: Mingxia Zou and Wei Du; Software: Mingxia Zou; Validation: Mingxia Zou and Hongmin Fan; Writing the original draft: Mingxia Zou; Writing the review: Mingxia Zou and Hongmin Fan. All authors have responded to the modifications of the study protocol and approved the final manuscript.

Acknowledgments The authors thank all participants and administrators of this study.

Received: ; Accepted: February 24, 2025

REFERENCES

- Lou YM, Zhang YY, Zhao P, et al. Association of fasting plasma glucose change trajectory and risk of hypertension: a cohort study in China. Endocr Connect, 2022; 11, e210464.
- Li DK, Song LL, Wang LL, et al. Association of fasting plasma glucose trajectory with lifetime risk of cardiovascular disease. Eur J Clin Nutr, 2023; 77, 335–41.
- Jun H, Lee J, Lee HA, et al. Fasting blood glucose variability and unfavorable trajectory patterns are associated with the risk of colorectal cancer. Gut Liver, 2022; 16, 423–32.
- Alkhatib A, Tsang C, Tiss A, et al. Functional foods and lifestyle approaches for diabetes prevention and management. Nutrients, 2017; 9, 1310.
- Papakonstantinou E, Oikonomou C, Nychas G, et al. Effects of diet, lifestyle, chrononutrition and alternative dietary interventions on postprandial glycemia and insulin resistance. Nutrients, 2022; 14, 823.
- Kim S, Yun B, Lee S, et al. Occupational noise exposure and incidence of high fasting blood glucose: A 3-Year, multicenter,

retrospective study. Int J Environ Heal R, 2021; 18, 9388.

- Liao YJ, Xu CX, Ma CQ, et al. Effect of occupational factors on pre-diabetes mellitus among iron and steel workers. Chin J Epidemiol, 2020; 41, 929–33 (In Chinese).
- Yun B, Sim J, Lee S, et al. The relationship between occupational dust exposure and incidence of diabetes in male workers: A retrospective cohort study. Diabet Med, 2022; 39, e14837.
- Huang CC, Chen TH, Ho CH, et al. Risks of Developing diabetes and hyperglycemic crisis following carbon monoxide poisoning: A study incorporating epidemiologic analysis and animal experiment. Clin Epidemiol Glob, 2022; 14, 1265–79.
- Yu LL, Liu W, Wang X, et al. A review of practical statistical methods used in epidemiological studies to estimate the health effects of multi-pollutant mixture. Environ Pollut, 2022; 306, 119356.
- Wang X, Mukherjee B, Park SK. Associations of cumulative exposure to heavy metal mixtures with obesity and its comorbidities among U. S. adults in NHANES 2003-2014. Environ Int, 2018; 121, 683–94.
- Guo XW, Li N, Wang H, et al. Combined exposure to multiple metals on cardiovascular disease in NHANES under five statistical models. Environ Res, 2022; 215, 114435.
- 13. Park SK, Tao YB, Meeker JD, et al. Environmental risk score as a new tool to examine multi-pollutants in epidemiologic research: an example from the NHANES study using serum lipid levels. PLoS One, 2014; 9, e98632.
- Chen YY, Yang YZ, Zheng ZW, et al. Influence of occupational exposure on hyperuricemia in steelworkers: a nested casecontrol study. BMC Public Health, 2022; 22, 1508.
- 15. Ministry of Health of the People's Republic of China. Measurement of physical agents in workplace Part 7: Heat stress: GBZ/T 189.7—2007. Beijing: Standards Press of China, 2007 (In Chinese).
- Ministry of Health of the People's Republic of China. Classification of occupational hazards at workplaces Part 3: Occupational exposure to heat stress: GBZ/T 229.3—2010. Beijing: The People's Health Press Co., Ltd, 2010 (In Chinese).
- 17. Ministry of Health of the People's Republic of Chin. Measurement of physical agents in workplace Part 8: Noise: GBZ 189.8—2007. Beijing: Standards Press of China, 2007 (In Chinese).
- Ministry of Health of the People's Republic of Chin. Classification of occupational hazards at workplaces Part 4: Occupational exposure to noise: GBZ/T 229.4—2010. Beijing: Standards Press of China, 2012 (In Chinese).
- Ministry of Health of the People's Republic of Chin. Determination of dust in the air of workplace-Part 1: Total dust concentration: GBZ-T 192.1—2007. Beijing: The People's Health Press Co., Ltd, 2007 (In Chinese).
- National Health Commission of the People's Republic of China. Occupational exposure limits for hazardous agents in the workplace—Part 1: Chemical hazardous agents: GBZ 2.1—2019. Beijing: Standards Press of China, 2019 (In Chinese).
- National Health and Family Planning Commission of the People's Republic of China. Determination of toxic substances in workplace air—Part 37: Carbon monoxide and carbon dioxide: GBZ/T 300.37—2017. Beijing: Standards Press of China, 2017 (In Chinese).
- 22. Lou XW, He Q. Validity and reliability of the international physical activity questionnaire in chinese hemodialysis patients: A multicenter study in China. Med Sci Monit, 2019; 25, 9402–8.
- 23. Satija A, Bhupathiraju SN, Spiegelman D, et al. Healthful and unhealthful plant-based diets and the risk of coronary heart

disease in U. S. Adults. J Am Coll Cardiol, 2017; 70, 411-22.

- 24. Yuan Y, Chen SM, Lin CJ, et al. Association of triglycerideglucose index trajectory and frailty in urban older residents: evidence from the 10-year follow-up in a cohort study. Cardiovasc Diabetol, 2023; 22, 264.
- Nagin DS, Jones BL, Elmer J. Recent advances in group-based trajectory modeling for clinical research. Annu Rev Clin Psycho, 2024; 20, 285–305.
- 26. Park SK, Zhao ZC, Mukherjee B. Construction of environmental risk score beyond standard linear models using machine learning methods: application to metal mixtures, oxidative stress and cardiovascular disease in NHANES. Environ Health-GLOB, 2017; 16, 102.
- 27. Li S, Zhou Y, Williams G, et al. Seasonality and temperature effects on fasting plasma glucose: A population-based longitudinal study in China. Diabetes Metab, 2016; 42, 267–75.
- 28. Liu Y, Wang DM, Huang XZ, et al. Temporal trend and global burden of type 2 diabetes attributable to non-optimal temperature, 1990-2019: an analysis for the Global Burden of Disease Study 2019. Environ Sci Pollut Res, 2023; 30, 82686–95.
- 29. Gao DH, Friedman S, Hosler AS, et al. Ambient heat and diabetes hospitalizations: Does the timing of heat exposure matter?. Sci Total Environ, 2024; 912, 169011.
- 30. Vetter C, Dashti HS, Lane JM, et al. Night shift work, genetic risk, and type 2 diabetes in the UK biobank. Diabetes Care, 2018; 41, 762–9.

- Chang TY, Yu TY, Liu CS, et al. Occupational noise exposure and its association with incident hyperglycaemia: a retrospective cohort study. Sci Rep, 2020; 10, 8584.
- Guo JQ, Liu YF, Hu CX, et al. Long-term exposure to dust and noise can increase the risk of hypertension-a population based, cross-sectional study. J Occup Environ Med, 2023; 65, e128–33.
- Li X, Wang CY, Fan HM, et al. Association between joint of heat and noise and metabolic syndrome in steel workers. J Hyg Res, 2015; 44, 77–81,90 (In Chinese).
- 34. Golmohammadi R, Darvishi E. The combined effects of occupational exposure to noise and other risk factors a systematic review. Noise Health, 2019; 21, 125–41.
- 35. Münzel T, Hahad O, Sørensen M, et al. Environmental risk factors and cardiovascular diseases: a comprehensive expert review. Cardiovasc Res, 2022; 118, 2880–902.
- 36. Gao YY, Gan T, Jiang LL, et al. Association between shift work and risk of type 2 diabetes mellitus: a systematic review and dose-response meta-analysis of observational studies. Chronobiol Int, 2020; 37, 29–46.
- Leproult R, Holmbäck U, Van Cauter E. Circadian misalignment augments markers of insulin resistance and inflammation, independently of sleep loss. Diabetes, 2014; 63, 1860–9.
- Esquirol Y, Bongard V, Ferrieres J, et al. Shiftwork and higher pancreatic secretion: early detection of an intermediate state of insulin resistance?. Chronobiol Int, 2012; 29, 1258–66.