

Original Article



Long-Term National Trends in the Prevalence of Type 2 Diabetes Mellitus and the Association with Social Determinants of Health in Korean Adults, 2007–2024

Jihu Im^{1,2,&}, Juyeong Kim^{1,3,&}, Seoyoung Park^{1,2}, Yesol Yim^{1,2}, Hyunjee Kim^{1,2},
Jihye Choi^{1,3}, and Dongkeon Yon^{1,2,3,4,#}

1. Center for Digital Health, Medical Science Research Institute, Kyung Hee University Medical Center, Kyung Hee University College of Medicine, Seoul, South Korea; 2. Department of Precision Medicine, Kyung Hee University College of Medicine, Seoul, South Korea; 3. Department of Regulatory Science, Kyung Hee University, Seoul, South Korea; 4. Department of Pediatrics, Kyung Hee University College of Medicine, Seoul, South Korea

Abstract

Background Type 2 diabetes mellitus (T2DM) is linked to social determinants of health (SDOH); however, long-term trends in its prevalence and the association with SDOH remain unclear. Thus, we investigated national trends in T2DM prevalence and the cumulative burden of SDOH.

Methods We analyzed data from the Korea National Health and Nutrition Examination Survey (2007–2024), including adults aged ≥ 19 years. T2DM was defined by physician diagnosis, use of anti-diabetic medication, fasting glucose ≥ 126 mg/dL, or hemoglobin A1c $\geq 6.5\%$. SDOH was assessed across seven key domains—education level, marital status, household income, food security, type of health insurance, employment status, and home ownership— and classified into low- (≤ 2) and high-burden (> 2) groups; the cumulative score was also analyzed as ordinal and continuous variables to evaluate dose-response relationships. Trends were examined using weighted linear regression, and adjusted odds ratios (aORs) were estimated using multivariable logistic regression across pre- (2007–2019), intra- (2020–2022), and preliminary post-pandemic (2023–2024) periods. Age-standardized estimates and sensitivity analyses, including inverse probability weighting and mediation analysis, were performed to ensure robustness.

Results A total of 85,695 adults (males; 35,392, 41.30%) were included. The prevalence of T2DM showed increasing trend (8.55%, 95% CI: 7.96% to 9.13% in 2007–2009; 11.66%, 95% CI: 11.11% to 12.22% in 2016–2019), and a surge was observed at the onset of the pandemic in 2020, peaking at 13.66% (95% CI: 12.24% to 15.07%) in 2021 before rebounding to 14.19% (12.99% - 15.39%) in 2024. High cumulative SDOH was associated with T2DM (aOR=1.33, 95% CI: 1.25 to 1.41), showing a clear dose-response relationship where the risk progressively escalated with higher SDOH scores (score ≥ 6 : 1.79 [1.58 to 2.04]). Among individual components, lack of private health insurance showed the strongest association (aOR, 1.30 [95% CI: 1.22 to 1.39]). Other major risk factors included established demographic and clinical predictors such as older age, male, and obesity.

Conclusions T2DM prevalence has increased over time, and a high cumulative burden of SDOH remains consistently associated with this prevalence, highlighting the need for equity-focused public health strategies.

Key words: COVID-19; Social determinants of health; South Korea; Trend; Type 2 diabetes mellitus

Biomed Environ Sci, 2026; 39(x): 1-19

doi: [10.3967/bes2026.090](https://doi.org/10.3967/bes2026.090)

ISSN: 0895-3988

www.besjournal.com (full text)

CN: 11-2816/Q

Copyright ©2026 by China CDC

[&]These authors contributed equally to this work.

[#]Correspondence should be addressed to Dongkeon Yon, MD, PhD, E-mail: yonkkang@gmail.com

INTRODUCTION

T2DM represents a leading global public health challenge, with its escalating prevalence imposing substantial personal, societal, and economic burdens worldwide^[1]. The burden of T2DM is deeply intertwined with social and economic conditions, serving as a critical indicator of population health and health inequality^[1]. While established risk factors for T2DM such as age, sex, and obesity are well known^[2], a comprehensive understanding of how a cumulative burden of social determinants of health (SDOH) influences long-term trends remains less explored^[3]. A deeper analysis focusing on these synergistic effects of multiple social disadvantages is crucial for developing holistic, equity-focused interventions that move beyond individual-level risk modification^[4,5].

The coronavirus disease 2019 (COVID-19) pandemic has acted as a global stress test, causing widespread disruptions to health management^[6,7], altered dietary patterns^[8], and daily life behaviors^[9,10], which may have influenced the prevalence and risk of T2DM, particularly among vulnerable populations^[11]. Specifically, pandemic-driven restrictions on routine medical services and physical activity have disproportionately burdened vulnerable populations, who face greater barriers to adopting technology-based care and maintaining healthy lifestyles^[12,13]. Consequently, while SDOH are associated with T2DM risk^[4,5], the pandemic may have exacerbated these existing inequalities^[12]. Therefore, a detailed examination is essential to understand the long-term trends in T2DM, the specific impact of cumulative SDOH, and how this relationship may have been intensified by the distinct phases of the pandemic.

Therefore, this study aimed to investigate the long-term trends in T2DM prevalence among Korean adults, using nationally representative data from the Korea National Health and Nutrition Examination Survey (KNHANES) from 2007 to 2024. We examined changes in T2DM prevalence across pre-, intra-, and preliminary post-pandemic periods, and identified key risk factors, with a particular focus on the cumulative impact of SDOH. The findings from this study are expected to provide critical evidence to inform the development of integrated public health strategies and policies aimed at reducing health inequalities in T2DM management.

METHODS

Data Source

This study utilized data from the KNHANES, a nationally representative survey conducted by the Korea Disease Control and Prevention Agency (KDCA) from 2007 to 2024^[14]. The KNHANES employs a multistage, stratified, probability-based cluster sampling design, with survey areas and households as the primary and secondary sampling units, respectively^[14]. Sampling weights were applied to ensure national representativeness and to account for differential probabilities of selection and response. The prevalence of T2DM was estimated from a nationally representative sample of 85,695 adults aged 19 years or older in South Korea. We categorized the survey periods into three pandemic-related phases based on the timing of the first confirmed case in South Korea (January 19, 2020), the World Health Organization (WHO)'s pandemic declaration (March 11, 2020), and the official lifting of the pandemic alert by the Korean government in May 2023: pre- (2007–2019), intra- (2020–2022), and preliminary post-pandemic (2023–2024) periods^[15].

The research protocol was approved by the Institutional Review Boards of the KDCA (2007-02CON-04-P, 2008-04EXP-01-C, 2009-01CON-03-2C, 2010-02CON-21-C, 2011-02CON-06-C, 2012-01EXP-01-2C, 2013-07CON-03-4C, 2013-12EXP-03-5C, 2018-01-03-P-A, 2018-01-03-C-A, 2018-01-03-2C-A, 2018-01-03-5C-A, 2018-01-03-4C-A, 2022-11-16-R-A, and 2022-11-16-R-03). Written informed consent was obtained from all participants prior to their participation. The study was conducted in accordance with the ethical principles of the Declaration of Helsinki.

Ascertainment of T2DM

In this study, T2DM was defined as meeting at least one of the following criteria: a self-reported physician diagnosis, current use of any anti-diabetic medication, a fasting glucose level of 126 mg/dL or higher, or a hemoglobin A1c level of 6.5% or higher. Participants not meeting any of these criteria were classified as the non-diabetic group^[16].

Definition of Social Determinants of Health

SDOH refer to the social, economic, and physical factors that influence the environment in which individuals are born, grow up, and live^[17]. These factors play a substantial role in shaping health outcomes and have a significant impact on health equity^[17]. SDOH in this study included education

level (high school graduate or higher and less than high school), marital status (married and single, separated, widowed, or divorced), household income (high [third or highest quartile] and low [lowest or second quartile]), food security (full security and marginal, low, or very low food security), type of health insurance (classified solely by the presence of private health insurance: private and government or none), employment status (employed, student, or retired and unemployed), and home ownership (own home and rent home or other arrangement).

To assess the cumulative burden of SDOH, each of the seven SDOH variables was first dichotomized into either a favorable or unfavorable condition. The unfavorable conditions were defined as: high school or lower education level; single, separated, widowed, or divorced; low household income; marginal, low, or very low food security; having government or no insurance; unemployment; and rent home or other arrangement. A cumulative score was then calculated based on the number of these unfavorable conditions (range, 0–7) for each participant. Due to the small number of participants with a score of 7, scores of 6 and 7 were combined into a single category, resulting in a final score range of 0 to ≥ 6 ^[18]. Finally, this cumulative score was dichotomized into two groups for analysis: high-burden (more than 2 unfavorable SDOH) vs. low-burden (2 or fewer), using the median value of the score which was 2 as the cut-point^[19]. Since there is no universally established clinical cut-off for SDOH burden, a data-driven threshold, which was 2, based on the median was selected. Additionally, to evaluate the dose-response relationship and the graded impact of social disadvantage on T2DM risk, the raw cumulative SDOH score (range, 0 to ≥ 6) was further analyzed as both an ordinal and a continuous variable in our models.

Covariates

The covariates included in this study were sex (male and female), age group (19–39, 40–59, 60–69, and ≥ 70 years)^[20], region of residence (urban and rural), body mass index (BMI) group, smoking status (current smoker, ex-smoker, and never-smoker), drinking status (current drinker, ex-drinker, and never-drinker), blood pressure (normal, elevated, stage 1 hypertension, and stage 2 hypertension), and physical activity (sufficient and insufficient). BMI was classified according to the Asia-Pacific guidelines as underweight (< 18.5 kg/m²), normal weight (18.5–22.9 kg/m²), overweight (23.0–24.9 kg/m²),

and obese (≥ 25.0 kg/m²)^[21]. Blood pressure was classified according to the 2025 American College of Cardiology and the American Heart Association guidelines. Systolic blood pressure (SBP) and diastolic blood pressure (DBP) were used to define categories as follows: normal (SBP < 120 and DBP < 80 mmHg), elevated (120–129 and < 80 mmHg), stage 1 hypertension (130–139 or 80–89 mmHg), and stage 2 hypertension (≥ 140 or ≥ 90 mmHg)^[22]. Physical activity was evaluated based on the framework of the WHO's Global Physical Activity Questionnaire. Participants were considered to have sufficient physical activity if they engaged in ≥ 150 minutes/week of moderate-intensity activity, ≥ 75 minutes/week of vigorous-intensity activity, or an equivalent combination, where one minute of vigorous activity was counted as two minutes of moderate activity^[23]. BMI and blood pressure were obtained through direct anthropometric and clinical measurements, whereas all other covariate data were obtained through self-reported questionnaires administered by trained interviewers.

Statistical Analysis

The outcomes of this study were presented using qualitative data expressed as proportions or percentages^[21]. To ensure accurate population-level estimates, a weighted complex sampling analysis was applied to account for the stratified sampling design and to minimize the influence of differences in participant numbers across survey years. The association between the COVID-19 pandemic period and the prevalence of T2DM was evaluated using weighted linear regression models. We also applied weighted linear regression to assess how the prevalence of T2DM varied across levels of cumulative unfavorable SDOH within each pandemic period. β coefficients with 95% confidence intervals (CIs) were estimated, and differences in β coefficients (β_{diff}) across pandemic phases were compared to assess temporal changes in both overall prevalence and SDOH-related disparities. To test the statistical significance of these temporal variations, p-values for interaction were calculated. To identify risk factors associated with T2DM, multivariable logistic regression models were employed, and the results were presented as adjusted odds ratios (aORs) along with their corresponding 95% CIs. In addition, age-standardized prevalence and aORs were estimated to account for potential demographic shifts across survey years.

Several sensitivity analyses were performed to ensure robustness. To address potential selection

bias, inverse probability weighting (IPW) was performed, with inclusion probabilities estimated via multivariable logistic regression. To further validate the SDOH score, variance inflation factors (VIFs) were derived from weighted multiple linear regression models to assess multicollinearity among SDOH variables. Furthermore, to account for the differential contributions of individual SDOH components, a weighted SDOH score was constructed, with weights assigned based on the log-odds ratios from multivariable logistic regression models. This weighted score was analyzed as both a continuous variable and categorized into quartiles. Additionally, mediation analysis was conducted to evaluate the potential pathways between lifestyle factors (overweight or obesity, current smoking, current drinking, and insufficient physical activity) and T2DM. All statistical analyses were performed using two-sided tests, and a P -value < 0.05 was considered statistically significant, utilizing SAS software (version 9.4; SAS Institute, Cary, NC, USA), with the exception of the mediation analysis, which was performed using R software (version 4.5.2; R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

Baseline Characteristics

Of the 140,372 participants surveyed during the study period, participants under the age of 19 were excluded (Supplementary Figure S1). Among the remaining 111,381 adults, an additional 25,686 participants were excluded due to missing data on BMI, smoking status, drinking status, blood pressure, physical activity, education level, marital status, household income, food security, type of health insurance, employment status, home ownership, and T2DM status. As a result, the final survey participants consisted of 85,695 individuals (males, 35,392 [41.30%]; 19–39, 23,460 [27.38%]) (Table 1). To minimize selection bias, a comparison of baseline characteristics between included and excluded participants is provided in Supplementary Table S1, and additional IPW analysis did not indicate meaningful selection bias arising from these exclusions (Supplementary Table S2).

Trends in T2DM Prevalence across Pandemic Periods and Cumulative SDOH

Figure 1 and Table 2 present the weighted prevalence of T2DM across the pre-, intra-, and

preliminary post-pandemic periods. The overall prevalence of T2DM showed a consistent increasing trend, rising from 8.55% (95% CI: 7.96 to 9.13) in 2007–2009 to a peak of 13.66% (12.24 to 15.07) in 2021. Notably, the onset of the COVID-19 pandemic marked a significant surge in prevalence, with the odds of having T2DM in 2020 being higher than in the 2016–2019 period ($aOR = 1.18$, 95% CI: 1.04 to 1.35) (Supplementary Table S3); however, the rate of increase was significantly decelerated at the transition from the pre- to intra-pandemic period prevalence (β_{diff} , -0.051 [95% CI: -0.097 to -0.004]; P for interaction < 0.05) (Supplementary Table S4). Furthermore, following the peak of 2021, prevalence declined through 2022–2023 before rebounding to 14.19% (12.99 to 15.39) in 2024. These trends remained consistent generally when age-standardized prevalence was calculated (Supplementary Tables S5).

Consistent with these overall trends, the burden of T2DM was disproportionately higher among socially disadvantaged groups. When stratified by cumulative unfavorable SDOH (Figure 2), both low- and high-burden groups exhibited the same overarching pattern of rising T2DM prevalence across periods. However, individuals with a cumulative SDOH score greater than 2 consistently showed substantially higher prevalence levels than those with lower SDOH burden across all time points. Furthermore, a clear dose-response relationship was observed, with T2DM prevalence increasing progressively with higher SDOH burden scores across all survey periods.

Association Between SDOH and T2DM Prevalence

Table 3 summarizes the aOR for sociodemographic factors associated with T2DM prevalence. Individuals with high-burden group had a higher likelihood of T2DM compared to those with low-burden group (aOR , 1.33 [95% CI: 1.25 to 1.41]), highlighting the cumulative influence of social disadvantage on T2DM risk. This association exhibited a significant graded impact, where the risk of T2DM progressively escalated with increasing raw SDOH scores (continuous value: aOR , 1.11 [95% CI: 1.09 to 1.13]). Specifically, compared to a score of 0, the risk significantly increased starting from a score of 3 (aOR , 1.30 [95% CI: 1.17 to 1.45]) and peaking at a score of ≥ 6 (1.79 [1.58 to 2.04]). Age-standardized aOR s for the cumulative SDOH score and burden group yielded consistent associations (Supplementary Table S6). A similar graded pattern was also observed using the weighted SDOH score,

Table 1. General characteristics of Korean adults based on data obtained from the KNHANES, 2007–2024 (*n*=85,695)

Characteristic	Total	Pre-pandemic				Intra-pandemic			Preliminary post-pandemic	
		2007–2009	2010–2012	2013–2015	2016–2019	2020	2021	2022	2023	2024
Overall, <i>n</i> (%)	85,695 (100.00)	13,841 (16.15)	15,328 (17.89)	13,290 (15.51)	20,108 (23.46)	4,077 (4.76)	4,249 (4.96)	4,426 (5.16)	5,140 (6.00)	5,236 (6.11)
Crude rate, <i>n</i> (%)										
Sex, <i>n</i> (%)										
Male	35,392 (41.30)	5,439 (6.35)	6,140 (7.16)	5,390 (6.29)	8,436 (9.84)	1,769 (2.06)	1,821 (2.13)	1,924 (2.25)	2,224 (2.60)	2,249 (2.62)
Female	50,303 (58.70)	8,402 (9.80)	9,188 (10.72)	7,900 (9.22)	11,672 (13.62)	2,308 (2.69)	2,428 (2.83)	2,502 (2.92)	2,916 (3.40)	2,987 (3.49)
Age group (years), <i>n</i> (%)										
19–	23,460 (27.38)	4,393 (5.13)	4,399 (5.13)	3,713 (4.33)	5,401 (6.30)	1,103 (1.29)	964 (1.12)	1,097 (1.28)	1,168 (1.36)	1,222 (1.43)
40–	31,300 (36.52)	5,039 (5.88)	5,607 (6.54)	4,998 (5.83)	7,462 (8.71)	1,482 (1.73)	1,502 (1.75)	1,518 (1.77)	1,874 (2.19)	1,818 (2.12)
60–	16,076 (18.76)	2,359 (2.75)	2,733 (3.19)	2,375 (2.77)	3,617 (4.22)	808 (0.94)	878 (1.02)	983 (1.15)	1,186 (1.38)	1,137 (1.33)
≥ 70	14,859 (17.34)	2,050 (2.39)	2,589 (3.02)	2,204 (2.57)	3,628 (4.23)	684 (0.80)	905 (1.06)	828 (0.97)	912 (1.06)	1,059 (1.24)
Region of residence, <i>n</i> (%)										
Urban	38,840 (45.32)	5,870 (6.85)	6,991 (8.16)	6,153 (7.18)	9,368 (10.93)	1,805 (2.11)	1,999 (2.33)	1,908 (2.23)	2,291 (2.67)	2,455 (2.86)
Rural	46,855 (54.68)	7,971 (9.30)	8,337 (9.73)	7,137 (8.33)	10,740 (12.53)	2,272 (2.65)	2,250 (2.63)	2,518 (2.94)	2,849 (3.32)	2,781 (3.25)
BMI group, <i>n</i> (%)*										
Underweight	3,673 (4.29)	658 (0.77)	701 (0.82)	558 (0.65)	775 (0.90)	135 (0.16)	183 (0.21)	181 (0.21)	236 (0.28)	246 (0.29)
Normal	33,254 (38.81)	5,525 (6.45)	6,225 (7.26)	5,274 (6.15)	7,789 (9.09)	1,477 (1.72)	1,515 (1.77)	1,648 (1.92)	1,923 (2.24)	1,878 (2.19)
Overweight	19,902 (23.22)	3,324 (3.88)	3,564 (4.16)	3,129 (3.65)	4,624 (5.40)	925 (1.08)	1,001 (1.17)	1,007 (1.18)	1,155 (1.35)	1,173 (1.37)
Obese	28,866 (33.68)	4,334 (5.06)	4,838 (5.65)	4,329 (5.05)	6,920 (8.08)	1,540 (1.80)	1,550 (1.81)	1,590 (1.86)	1,826 (2.13)	1,939 (2.26)
Smoking status, <i>n</i> (%)										
Never smoker	52,753 (61.56)	8,377 (9.78)	9,383 (10.95)	8,325 (9.71)	12,410 (14.48)	2,502 (2.92)	2,632 (3.07)	2,671 (3.12)	3,157 (3.68)	3,296 (3.85)
Ex-smoker	18,259 (21.31)	2,671 (3.12)	3,141 (3.67)	2,648 (3.09)	4,402 (5.14)	915 (1.07)	983 (1.15)	1,085 (1.27)	1,189 (1.39)	1,225 (1.43)
Current smoker	14,683 (17.13)	2,793 (3.26)	2,804 (3.27)	2,317 (2.70)	3,296 (3.85)	660 (0.77)	634 (0.74)	670 (0.78)	794 (0.93)	715 (0.83)
Drinking status, <i>n</i> (%)										
Never drinker	10,895 (12.71)	2,129 (2.48)	2,243 (2.62)	1,740 (2.03)	2,283 (2.66)	455 (0.53)	525 (0.61)	440 (0.51)	571 (0.67)	509 (0.59)
Ex-drinker	14,619 (17.06)	2,121 (2.48)	2,343 (2.73)	2,258 (2.63)	3,418 (3.99)	784 (0.91)	923 (1.08)	854 (1.00)	933 (1.09)	985 (1.15)
Current drinker	60,181 (70.23)	9,591 (11.19)	10,742 (12.54)	9,292 (10.84)	14,407 (16.81)	2,838 (3.31)	2,801 (3.27)	3,132 (3.65)	3,636 (4.24)	3,742 (4.37)
Blood pressure, <i>n</i> (%)										
Normal	37,774 (44.08)	6,365 (7.43)	6,514 (7.60)	6,185 (7.22)	8,636 (10.08)	1,717 (2.00)	1,877 (2.19)	1,940 (2.26)	2,337 (2.73)	2,203 (2.57)
Elevated	5,409 (6.31)	725 (0.85)	839 (0.98)	829 (0.97)	1,239 (1.45)	255 (0.30)	362 (0.42)	350 (0.41)	404 (0.47)	406 (0.47)
Stage 1 hypertension	14,831 (17.31)	2,588 (3.02)	2,831 (3.30)	2,198 (2.56)	3,651 (4.26)	824 (0.96)	561 (0.65)	637 (0.74)	728 (0.85)	813 (0.95)
Stage 2 hypertension	27,681 (32.30)	4,163 (4.86)	5,144 (6.00)	4,078 (4.76)	6,582 (7.68)	1,281 (1.49)	1,449 (1.69)	1,499 (1.75)	1,671 (1.95)	1,814 (2.12)

Continued

Characteristic	Total	Pre-pandemic				Intra-pandemic			Preliminary post-pandemic	
		2007–2009	2010–2012	2013–2015	2016–2019	2020	2021	2022	2023	2024
Physical activity, <i>n</i> (%)										
Insufficient	38,895 (45.39)	3,631 (4.24)	5,026 (5.87)	5,845 (6.82)	11,497 (13.42)	2,336 (2.73)	2,502 (2.92)	2,378 (2.78)	2,766 (3.23)	2,914 (3.40)
Sufficient	46,800 (54.61)	10,210 (11.91)	10,302 (12.02)	7,445 (8.69)	8,611 (10.05)	1,741 (2.03)	1,747 (2.04)	2,048 (2.39)	2,374 (2.77)	2,322 (2.71)
Education level, <i>n</i> (%)										
High	60,323 (70.39)	9,889 (11.54)	9,721 (11.34)	8,956 (10.45)	14,345 (16.74)	3,072 (3.58)	3,075 (3.59)	3,341 (3.90)	3,946 (4.60)	3,978 (4.64)
Low	25,372 (29.61)	3,952 (4.61)	5,607 (6.54)	4,334 (5.06)	5,763 (6.73)	1,005 (1.17)	1,174 (1.37)	1,085 (1.27)	1,194 (1.39)	1,258 (1.47)
Marital status, <i>n</i> (%)										
Married	60,567 (70.68)	10,172 (11.87)	11,485 (13.40)	9,486 (11.07)	14,073 (16.42)	2,703 (3.15)	2,809 (3.28)	2,958 (3.45)	3,471 (4.05)	3,410 (3.98)
Single, separated, widowed, or divorced	25,128 (29.32)	3,669 (4.28)	3,843 (4.48)	3,804 (4.44)	6,035 (7.04)	1,374 (1.60)	1,440 (1.68)	1,468 (1.71)	1,669 (1.95)	1,826 (2.13)
Household income, <i>n</i> (%)										
High	47,576 (55.52)	7,290 (8.51)	8,262 (9.64)	7,359 (8.59)	11,272 (13.15)	2,426 (2.83)	2,406 (2.81)	2,475 (2.89)	2,989 (3.49)	3,097 (3.61)
Low	38,119 (44.48)	6,551 (7.64)	7,066 (8.25)	5,931 (6.92)	8,836 (10.31)	1,651 (1.93)	1,843 (2.15)	1,951 (2.28)	2,151 (2.51)	2,139 (2.50)
Food security, <i>n</i> (%)										
Full security	45,951 (53.62)	5,527 (6.45)	6,688 (7.80)	6,441 (7.52)	11,136 (12.99)	2,568 (3.00)	2,749 (3.21)	3,078 (3.59)	3,646 (4.25)	4,118 (4.81)
Marginal, low, or very low security	39,744 (46.38)	8,314 (9.70)	8,640 (10.08)	6,849 (7.99)	8,972 (10.47)	1,509 (1.76)	1,500 (1.75)	1,348 (1.57)	1,494 (1.74)	1,118 (1.30)
Type of health insurance, <i>n</i> (%)										
Private	64,538 (75.31)	9,053 (10.56)	10,715 (12.50)	9,829 (11.47)	15,690 (18.31)	3,395 (3.96)	3,377 (3.94)	3,647 (4.26)	4,328 (5.05)	4,504 (5.26)
Government or none	21,157 (24.69)	4,788 (5.59)	4,613 (5.38)	3,461 (4.04)	4,418 (5.16)	682 (0.80)	872 (1.02)	779 (0.91)	812 (0.95)	732 (0.85)
Employment status, <i>n</i> (%)										
Employed, student, or retired	50,525 (58.96)	7,837 (9.15)	8,800 (10.27)	7,637 (8.91)	12,024 (14.03)	2,418 (2.82)	2,556 (2.98)	2,724 (3.18)	3,218 (3.76)	3,311 (3.86)
Unemployed	35,170 (41.04)	6,004 (7.01)	6,528 (7.62)	5,653 (6.60)	8,084 (9.43)	1,659 (1.94)	1,693 (1.98)	1,702 (1.99)	1,922 (2.24)	1,925 (2.25)
Home ownership, <i>n</i> (%)										
Own home	60,394 (70.48)	9,703 (11.32)	11,512 (13.43)	9,103 (10.62)	13,945 (16.27)	2,756 (3.22)	2,889 (3.37)	3,067 (3.58)	3,618 (4.22)	3,801 (4.44)
Rent home or other arrangement	25,301 (29.52)	4,138 (4.83)	3,816 (4.45)	4,187 (4.89)	6,163 (7.19)	1,321 (1.54)	1,360 (1.59)	1,359 (1.59)	1,522 (1.78)	1,435 (1.67)
SDOH burden score, <i>n</i> (%)										
0	10,037 (11.71)	1,129 (1.32)	1,356 (1.58)	1,245 (1.45)	2,557 (2.98)	553 (0.65)	628 (0.73)	710 (0.83)	856 (1.00)	1,003 (1.17)
1	18,739 (21.87)	2,690 (3.14)	3,126 (3.65)	2,787 (3.25)	4,591 (5.36)	1,004 (1.17)	956 (1.12)	1,027 (1.20)	1,253 (1.46)	1,305 (1.52)
2	19,216 (22.42)	3,069 (3.58)	3,501 (4.09)	3,075 (3.59)	4,415 (5.15)	908 (1.06)	908 (1.06)	976 (1.14)	1,201 (1.40)	1,163 (1.36)
3	15,138 (17.67)	2,679 (3.13)	2,819 (3.29)	2,420 (2.82)	3,389 (3.95)	694 (0.81)	696 (0.81)	770 (0.90)	847 (0.99)	824 (0.96)
4	11,300 (13.19)	2,030 (2.37)	2,231 (2.60)	1,870 (2.18)	2,534 (2.96)	492 (0.57)	533 (0.62)	514 (0.60)	549 (0.64)	547 (0.64)
5	6,693 (7.81)	1,314 (1.53)	1,428 (1.67)	1,068 (1.25)	1,502 (1.75)	265 (0.31)	324 (0.38)	265 (0.31)	274 (0.32)	253 (0.30)

Continued

Characteristic	Total	Pre-pandemic				Intra-pandemic			Preliminary post-pandemic	
		2007–2009	2010–2012	2013–2015	2016–2019	2020	2021	2022	2023	2024
≥6	4,572 (5.34)	930 (1.09)	867 (1.01)	825 (0.96)	1,120 (1.31)	161 (0.19)	204 (0.24)	164 (0.19)	160 (0.19)	141 (0.16)
Cumulative unfavorable SDOH, n (%)										
Low-burden (≤ 2)	47,992 (56.00)	6,888 (8.04)	7,983 (9.32)	7,107 (8.29)	11,563 (13.49)	2,465 (2.88)	2,492 (2.91)	2,713 (3.17)	3,310 (3.86)	3,471 (4.05)
High-burden (> 2)	37,703 (44.00)	6,953 (8.11)	7,345 (8.57)	6,183 (7.22)	8,545 (9.97)	1,612 (1.88)	1,757 (2.05)	1,713 (2.00)	1,830 (2.14)	1,765 (2.06)
Weighted rate (95% CI)										
Sex, weighted % (95% CI)										
Male	49.56 (49.15 to 49.97)	48.95 (48.11 to 49.79)	49.35 (48.51 to 50.19)	48.92 (48.06 to 49.79)	49.54 (48.86 to 50.22)	49.98 (48.44 to 51.52)	49.70 (48.09 to 51.31)	49.83 (48.29 to 51.37)	49.34 (48.09 to 50.60)	50.25 (49.02 to 51.48)
Female	50.44 (50.03 to 50.85)	51.05 (50.21 to 51.89)	50.65 (49.81 to 51.49)	51.08 (50.21 to 51.94)	50.46 (49.78 to 51.14)	50.02 (48.48 to 51.56)	50.30 (48.69 to 51.91)	50.17 (48.63 to 51.71)	50.66 (49.40 to 51.91)	49.75 (48.52 to 50.98)
Age group (years), weighted % (95% CI)										
19-	35.48 (34.81 to 36.16)	42.30 (40.72 to 43.87)	40.01 (38.61 to 41.40)	38.19 (36.80 to 39.57)	35.47 (34.26 to 36.67)	34.99 (32.33 to 37.66)	33.45 (31.03 to 35.86)	32.75 (30.20 to 35.30)	32.39 (30.01 to 34.76)	31.56 (29.20 to 33.93)
40-	39.30 (38.71 to 39.90)	39.13 (37.81 to 40.44)	40.11 (38.98 to 41.25)	40.77 (39.57 to 41.96)	40.38 (39.33 to 41.42)	40.37 (38.01 to 42.74)	38.99 (36.88 to 41.09)	38.36 (35.97 to 40.75)	38.17 (36.29 to 40.05)	37.67 (35.49 to 39.85)
60-	13.70 (13.31 to 14.09)	10.40 (9.76 to 11.04)	10.67 (10.02 to 11.33)	11.28 (10.61 to 11.94)	12.28 (11.67 to 12.90)	13.69 (12.24 to 15.15)	14.60 (13.29 to 15.91)	16.26 (14.60 to 17.92)	16.84 (15.49 to 18.20)	16.26 (14.74 to 17.78)
≥70	11.51 (11.10 to 11.93)	8.18 (7.57 to 8.79)	9.21 (8.57 to 9.85)	9.77 (9.09 to 10.45)	11.87 (11.17 to 12.58)	10.94 (9.15 to 12.74)	12.97 (11.26 to 14.68)	12.63 (10.97 to 14.29)	12.60 (11.20 to 14.01)	14.51 (12.85 to 16.18)
Region of residence, weighted % (95% CI)										
Urban	47.04 (45.83 to 48.25)	48.03 (45.82 to 50.23)	48.43 (46.15 to 50.70)	48.05 (45.98 to 50.12)	49.36 (47.03 to 51.69)	46.55 (38.96 to 54.15)	48.39 (41.26 to 55.53)	44.01 (37.16 to 50.86)	45.41 (38.74 to 52.07)	45.49 (38.73 to 52.25)
Rural	52.96 (51.75 to 54.17)	51.97 (49.77 to 54.18)	51.57 (49.30 to 53.85)	51.95 (49.88 to 54.02)	50.64 (48.31 to 52.97)	53.45 (45.85 to 61.04)	51.61 (44.47 to 58.74)	55.99 (49.14 to 62.84)	54.59 (47.93 to 61.26)	54.51 (47.75 to 61.27)
BMI group, weighted % (95% CI)*										
Underweight	4.50 (4.30 to 4.70)	4.99 (4.50 to 5.47)	4.84 (4.41 to 5.28)	4.43 (3.98 to 4.87)	4.14 (3.79 to 4.48)	3.63 (2.86 to 4.40)	4.55 (3.81 to 5.29)	4.47 (3.77 to 5.16)	4.83 (4.19 to 5.47)	4.71 (4.05 to 5.38)
Normal	37.52 (37.05 to 37.99)	39.82 (38.81 to 40.82)	40.58 (39.51 to 41.64)	40.10 (39.08 to 41.12)	38.77 (37.92 to 39.63)	35.59 (33.75 to 37.43)	35.80 (34.12 to 37.48)	36.22 (34.44 to 37.99)	36.75 (35.24 to 38.26)	34.95 (33.49 to 36.40)
Overweight	22.65 (22.27 to 23.04)	23.86 (22.99 to 24.72)	22.63 (21.81 to 23.46)	22.89 (22.05 to 23.73)	22.59 (21.91 to 23.28)	23.08 (21.58 to 24.59)	21.88 (20.45 to 23.31)	22.75 (21.36 to 24.15)	22.21 (20.96 to 23.45)	22.21 (21.14 to 23.27)
Obese	35.33 (34.83 to 35.83)	31.34 (30.31 to 32.37)	31.95 (30.91 to 32.98)	32.58 (31.58 to 33.58)	34.49 (33.67 to 35.32)	37.70 (35.92 to 39.49)	37.77 (35.81 to 39.73)	36.57 (34.55 to 38.59)	36.22 (34.59 to 37.84)	38.13 (36.59 to 39.68)
Smoking status, weighted % (95% CI)										
Never smoker	56.96 (56.46 to 57.46)	53.08 (52.09 to 54.06)	54.27 (53.33 to 55.20)	57.55 (56.60 to 58.51)	57.43 (56.61 to 58.24)	57.37 (55.48 to 59.26)	57.99 (56.17 to 59.81)	56.73 (54.92 to 58.53)	58.34 (56.71 to 59.98)	59.10 (57.40 to 60.80)
Ex-smoker	23.12 (22.72 to 23.52)	20.43 (19.65 to 21.21)	20.58 (19.76 to 21.39)	20.28 (19.49 to 21.07)	22.52 (21.89 to 23.15)	24.07 (22.65 to 25.49)	24.05 (22.61 to 25.49)	26.45 (24.89 to 28.00)	23.64 (22.41 to 24.87)	25.14 (23.69 to 26.58)
Current smoker	19.92 (19.47 to 20.37)	26.50 (25.50 to 27.50)	25.16 (24.16 to 26.16)	22.17 (21.18 to 23.16)	20.05 (19.27 to 20.84)	18.56 (16.86 to 20.27)	17.96 (16.44 to 19.48)	16.83 (15.04 to 18.61)	18.02 (16.62 to 19.41)	15.76 (14.35 to 17.18)

Continued

Characteristic	Total	Pre-pandemic				Intra-pandemic			Preliminary post-pandemic	
		2007–2009	2010–2012	2013–2015	2016–2019	2020	2021	2022	2023	2024
Drinking status, weighted % (95% CI)										
Never drinker	8.88 (8.60 to 9.16)	11.28 (10.63 to 11.92)	10.56 (9.88 to 11.23)	9.76 (9.13 to 10.39)	8.42 (7.94 to 8.90)	8.16 (7.04 to 9.27)	9.11 (7.92 to 10.29)	7.43 (6.45 to 8.41)	8.42 (7.46 to 9.39)	7.38 (6.47 to 8.29)
Ex-drinker	15.79 (15.42 to 16.17)	12.70 (12.01 to 13.38)	12.89 (12.17 to 13.60)	14.55 (13.80 to 15.29)	14.76 (14.17 to 15.35)	17.07 (15.51 to 18.62)	18.89 (17.43 to 20.35)	17.52 (16.16 to 18.89)	16.18 (14.95 to 17.41)	16.70 (15.54 to 17.86)
Current drinker	75.33 (74.84 to 75.82)	76.03 (75.10 to 76.96)	76.56 (75.60 to 77.51)	75.69 (74.72 to 76.66)	76.82 (76.04 to 77.59)	74.77 (72.72 to 76.83)	72.00 (70.11 to 73.89)	75.05 (73.28 to 76.81)	75.40 (73.68 to 77.12)	75.92 (74.25 to 77.59)
Blood pressure, weighted % (95% CI)										
Normal	47.98 (47.36 to 48.60)	49.80 (48.38 to 51.23)	47.40 (46.12 to 48.68)	51.37 (50.12 to 52.61)	46.74 (45.67 to 47.82)	45.64 (43.21 to 48.06)	48.55 (46.37 to 50.72)	47.37 (44.78 to 49.97)	49.96 (47.78 to 52.14)	45.58 (43.61 to 47.55)
Elevated	6.74 (6.47 to 7.00)	4.24 (3.83 to 4.65)	5.21 (4.76 to 5.67)	5.75 (5.26 to 6.24)	5.69 (5.31 to 6.07)	6.05 (5.05 to 7.05)	8.88 (7.83 to 9.92)	8.36 (7.36 to 9.36)	8.06 (7.31 to 8.80)	7.75 (6.87 to 8.62)
Stage 1 hypertension	18.00 (17.62 to 18.39)	20.81 (19.78 to 21.84)	20.04 (19.15 to 20.94)	18.26 (17.39 to 19.13)	20.20 (19.46 to 20.94)	21.78 (20.29 to 23.27)	15.06 (13.75 to 16.37)	15.04 (13.68 to 16.40)	14.75 (13.63 to 15.87)	16.76 (15.76 to 17.76)
Stage 2 hypertension	27.28 (26.75 to 27.82)	25.15 (24.07 to 26.24)	27.34 (26.27 to 28.41)	24.62 (23.62 to 25.63)	27.37 (26.49 to 28.24)	26.53 (24.47 to 28.60)	27.52 (25.54 to 29.49)	29.23 (27.00 to 31.45)	27.24 (25.38 to 29.09)	29.91 (28.05 to 31.77)
Physical activity, weighted % (95% CI)										
Insufficient	46.30 (45.71 to 46.88)	24.56 (23.35 to 25.77)	30.79 (29.66 to 31.92)	40.10 (38.81 to 41.40)	53.04 (52.02 to 54.06)	55.25 (53.15 to 57.35)	54.76 (52.80 to 56.71)	49.97 (47.81 to 52.13)	50.57 (48.58 to 52.57)	52.32 (50.16 to 54.48)
Sufficient	53.70 (53.12 to 54.29)	75.44 (74.23 to 76.65)	69.21 (68.08 to 70.34)	59.90 (58.61 to 61.19)	46.96 (45.94 to 47.98)	44.75 (42.65 to 46.85)	45.24 (43.29 to 47.20)	50.03 (47.87 to 52.19)	49.43 (47.43 to 51.42)	47.68 (45.52 to 49.84)
Education level, weighted % (95% CI)										
High	80.82 (80.24 to 81.40)	81.68 (80.65 to 82.71)	73.07 (71.72 to 74.41)	77.04 (75.84 to 78.23)	79.83 (78.77 to 80.89)	83.18 (80.65 to 85.70)	82.33 (79.98 to 84.67)	83.45 (81.04 to 85.86)	83.29 (81.28 to 85.30)	82.62 (80.55 to 84.70)
Low	19.18 (18.60 to 19.76)	18.32 (17.29 to 19.35)	26.93 (25.59 to 28.28)	22.96 (21.77 to 24.16)	20.17 (19.11 to 21.23)	16.82 (14.30 to 19.35)	17.67 (15.33 to 20.02)	16.55 (14.14 to 18.96)	16.71 (14.70 to 18.72)	17.38 (15.30 to 19.45)
Marital status, weighted % (95% CI)										
Married	65.26 (64.57 to 65.95)	69.88 (68.50 to 71.26)	68.46 (67.26 to 69.66)	67.65 (66.30 to 69.01)	66.28 (65.13 to 67.42)	63.80 (60.98 to 66.62)	63.27 (60.64 to 65.90)	63.66 (60.96 to 66.36)	63.02 (60.69 to 65.36)	62.55 (59.81 to 65.29)
Single, separated, widowed, or divorced	34.74 (34.05 to 35.43)	30.12 (28.74 to 31.50)	31.54 (30.34 to 32.74)	32.35 (30.99 to 33.70)	33.72 (32.58 to 34.87)	36.20 (33.38 to 39.02)	36.73 (34.10 to 39.36)	36.34 (33.64 to 39.04)	36.98 (34.64 to 39.31)	37.45 (34.71 to 40.19)
Household income, weighted % (95% CI)										
High	62.11 (61.25 to 62.97)	58.84 (57.02 to 60.67)	57.00 (55.41 to 58.59)	61.00 (59.24 to 62.76)	61.67 (60.18 to 63.17)	65.21 (61.59 to 68.82)	64.42 (61.02 to 67.82)	62.68 (59.41 to 65.95)	63.67 (60.73 to 66.60)	63.48 (60.32 to 66.64)
Low	37.89 (37.03 to 38.75)	41.16 (39.33 to 42.98)	43.00 (41.41 to 44.59)	39.00 (37.24 to 40.76)	38.33 (36.83 to 39.82)	34.80 (31.18 to 38.41)	35.58 (32.18 to 38.98)	37.32 (34.05 to 40.59)	36.33 (33.40 to 39.27)	36.52 (33.36 to 39.68)
Food security, weighted % (95% CI)										
Full security	61.24 (60.41 to 62.06)	41.67 (39.99 to 43.34)	43.16 (41.66 to 44.67)	49.19 (47.56 to 50.81)	56.90 (55.45 to 58.36)	64.94 (61.74 to 68.13)	66.99 (63.67 to 70.31)	70.76 (68.28 to 73.25)	71.59 (69.02 to 74.17)	80.02 (77.68 to 82.36)
Marginal, low, or very low security	38.76 (37.94 to 39.59)	58.33 (56.66 to 60.01)	56.84 (55.33 to 58.34)	50.81 (49.19 to 52.44)	43.10 (41.64 to 44.55)	35.06 (31.87 to 38.26)	33.01 (29.69 to 36.33)	29.24 (26.75 to 31.72)	28.41 (25.83 to 30.98)	19.98 (17.64 to 22.32)

Continued

Characteristic	Total	Pre-pandemic				Intra-pandemic			Preliminary post-pandemic	
		2007–2009	2010–2012	2013–2015	2016–2019	2020	2021	2022	2023	2024
Type of health insurance, weighted % (95% CI)										
Private	82.63 (82.16 to 83.09)	72.23 (71.01 to 73.45)	75.09 (73.97 to 76.21)	79.65 (78.59 to 80.71)	82.57 (81.68 to 83.47)	86.11 (84.33 to 87.89)	83.62 (81.81 to 85.43)	86.34 (84.84 to 87.84)	86.98 (85.40 to 88.55)	88.52 (87.25 to 89.79)
Government or none	17.37 (16.91 to 17.84)	27.77 (26.55 to 28.99)	24.91 (23.79 to 26.03)	20.35 (19.29 to 21.41)	17.43 (16.53 to 18.32)	13.89 (12.11 to 15.67)	16.38 (14.57 to 18.19)	13.66 (12.16 to 15.16)	13.02 (11.45 to 14.60)	11.48 (10.21 to 12.75)
Employment status, weighted % (95% CI)										
Employed, student, or retired	64.16 (63.61 to 64.71)	60.40 (59.18 to 61.61)	64.19 (63.06 to 65.31)	62.63 (61.54 to 63.72)	63.82 (62.85 to 64.79)	62.98 (60.78 to 65.17)	63.65 (61.72 to 65.57)	64.89 (62.80 to 66.98)	66.68 (64.95 to 68.41)	67.45 (65.64 to 69.26)
Unemployed	35.84 (35.29 to 36.39)	39.60 (38.39 to 40.82)	35.81 (34.69 to 36.94)	37.37 (36.28 to 38.46)	36.18 (35.21 to 37.15)	37.02 (34.83 to 39.22)	36.35 (34.43 to 38.28)	35.11 (33.02 to 37.20)	33.32 (31.59 to 35.05)	32.55 (30.74 to 34.36)
Home ownership, weighted % (95% CI)										
Own home	67.76 (66.71 to 68.81)	66.17 (64.08 to 68.26)	68.66 (66.96 to 70.36)	65.73 (63.69 to 67.77)	67.34 (65.67 to 69.01)	67.48 (63.39 to 71.56)	66.56 (61.79 to 71.32)	68.65 (65.04 to 72.27)	68.35 (64.87 to 71.83)	70.50 (66.97 to 74.02)
Rent home or other arrangement	32.24 (31.19 to 33.29)	33.83 (31.74 to 35.92)	31.34 (29.64 to 33.04)	34.27 (32.23 to 36.31)	32.66 (30.99 to 34.33)	32.52 (28.44 to 36.61)	33.44 (28.68 to 38.21)	31.35 (27.73 to 34.96)	31.65 (28.17 to 35.13)	29.50 (25.98 to 33.03)
SDOH burden score, weighted % (95% CI)										
0	14.87 (14.28 to 15.46)	9.43 (8.59 to 10.27)	9.06 (8.35 to 9.76)	10.57 (9.71 to 11.42)	13.72 (12.82 to 14.62)	15.53 (13.23 to 17.83)	17.12 (14.79 to 19.46)	17.96 (15.63 to 20.29)	17.51 (15.60 to 19.41)	21.09 (18.79 to 23.40)
1	24.42 (23.86 to 24.99)	21.25 (20.21 to 22.29)	20.96 (20.02 to 21.91)	23.17 (22.09 to 24.26)	24.74 (23.82 to 25.67)	26.90 (24.53 to 29.28)	24.48 (22.13 to 26.83)	25.15 (23.05 to 27.24)	26.02 (24.17 to 27.88)	26.24 (24.41 to 28.07)
2	23.67 (23.21 to 24.13)	24.10 (23.19 to 25.02)	24.70 (23.69 to 25.70)	24.47 (23.46 to 25.47)	23.28 (22.54 to 24.02)	22.92 (21.09 to 24.75)	23.24 (21.63 to 24.85)	23.33 (21.60 to 25.06)	24.36 (22.95 to 25.76)	22.88 (21.40 to 24.37)
3	16.90 (16.45 to 17.34)	19.44 (18.57 to 20.32)	19.15 (18.30 to 19.99)	18.32 (17.41 to 19.22)	17.05 (16.34 to 17.76)	16.18 (14.54 to 17.83)	15.46 (13.66 to 17.25)	16.37 (14.61 to 18.14)	16.12 (14.61 to 17.62)	14.74 (13.23 to 16.25)
4	11.14 (10.77 to 11.51)	13.32 (12.47 to 14.16)	13.72 (12.91 to 14.54)	12.61 (11.82 to 13.40)	11.55 (10.87 to 12.24)	10.44 (8.95 to 11.93)	10.71 (9.28 to 12.14)	10.15 (8.79 to 11.52)	9.48 (8.33 to 10.63)	8.98 (7.75 to 10.20)
5	5.76 (5.49 to 6.03)	7.89 (7.22 to 8.55)	8.01 (7.44 to 8.59)	6.72 (6.05 to 7.38)	5.91 (5.45 to 6.37)	5.40 (4.18 to 6.62)	5.52 (4.53 to 6.50)	4.68 (3.76 to 5.60)	4.24 (3.45 to 5.03)	4.12 (3.29 to 4.95)
≥ 6	3.24 (3.05 to 3.44)	4.57 (4.09 to 5.05)	4.41 (3.98 to 4.83)	4.15 (3.58 to 4.72)	3.74 (3.35 to 4.13)	2.62 (1.94 to 3.31)	3.47 (2.60 to 4.35)	2.36 (1.70 to 3.02)	2.28 (1.72 to 2.85)	1.95 (1.49 to 2.41)
Cumulative unfavorable SDOH, weighted % (95% CI)										
Low-burden (≤ 2)	62.96 (62.10 to 63.82)	54.78 (53.13 to 56.43)	54.72 (53.33 to 56.11)	58.21 (56.46 to 59.95)	61.75 (60.31 to 63.19)	65.35 (61.79 to 68.92)	64.84 (61.15 to 68.53)	66.43 (63.17 to 69.69)	67.88 (65.12 to 70.65)	70.21 (67.11 to 73.32)
High-burden (> 2)	37.04 (36.18 to 37.90)	45.22 (43.57 to 46.87)	45.28 (43.89 to 46.67)	41.79 (40.05 to 43.54)	38.25 (36.81 to 39.69)	34.65 (31.08 to 38.21)	35.16 (31.47 to 38.85)	33.57 (30.31 to 36.83)	32.12 (29.35 to 34.88)	29.79 (26.68 to 32.89)

Note. *According to Asian-Pacific guidelines, BMI is divided into four groups: underweight (< 18.5 kg/m²), normal weight (18.5–22.9 kg/m²), overweight (23.0–24.9 kg/m²), and obese (≥ 25.0 kg/m²); BMI, body mass index; CI: confidence interval; KNHANES, Korea National Health and Nutrition Examination Survey; SDOH, social determinants of health.

both as a continuous variable and across quartiles (Supplementary Table S7). Further mediation analysis suggested that lifestyle factors may not fully account for the observed association between SDOH burden and T2DM risk. Overweight or obesity and insufficient physical activity showed no significant mediating effects, while smoking and drinking status showed only minimal indirect effects (Supplementary Table S8).

Examining the contribution of each component underlying this cumulative burden, unfavorable conditions across all individual SDOH indicators were consistently associated with higher odds of T2DM compared to their more advantaged counterparts. Among the individual SDOH indicators, government or no health insurance coverage showed the strongest association with T2DM (*aOR*, 1.30 [95% *CI*: 1.22 to 1.39]), followed by unemployment (1.25 [1.18 to 1.33]), lower household income (1.25 [1.18 to 1.32]), and lower education level (1.25 [1.17 to 1.33]). In addition, all VIFs were below 2.0, confirming that the estimates were not influenced by multicollinearity among SDOH variables (Supplementary Table S9). These findings emphasize the important role of socioeconomic and structural disadvantages in shaping vulnerability to T2DM.

DISCUSSION

Key Findings

This study examined the long-term trends in T2DM prevalence according to SDOH among Korean




Figure 1. Trends in T2DM prevalence during the pre-, intra-, and preliminary post-pandemic periods from 2007 to 2024 ($n=85,745$), and shaded regions indicate 95% *CI*s. Abbreviations: *CI*: confidence intervals; T2DM, type 2 diabetes mellitus.

adults from 2007 to 2024, utilizing nationally representative data. The prevalence of T2DM continued to rise through the early intra-pandemic period, reaching its peak in 2021, followed by a modest decline during 2022–2023 before rebounding in 2024. T2DM prevalence was higher among individuals with a high-burden of SDOH group, with government or no health insurance coverage emerging as the most prominent individual risk factor. Similar patterns were observed in groups defined as having unfavorable levels of employment status, household income, education level, home ownership, and marital status. These associations remained consistent across multiple sensitivity analyses, including age-standardized estimates and weighted SDOH scores. Other major risk factors included older age, male, obesity, and smoking status. These findings highlight the necessity of an integrated public health strategy that simultaneously addresses both structural inequalities and individual-level risk factors to effectively manage the growing burden of T2DM in South Korea.

Plausible Underlying Mechanisms

Our findings indicate that a higher burden of unfavorable SDOH is associated with an increased prevalence of T2DM^[24]. This association was evident both in the cumulative SDOH burden and in several individual sociodemographic indicators, suggesting that social disadvantage plays a central role in shaping T2DM risk. The cumulative effects of socioeconomic adversity can manifest through chronic psychosocial stress. This stress stimulates the release of glucocorticoids and catecholamines, activates pro-inflammatory pathways (i.e., interleukin-6, tumor necrosis factor- α), and disrupts glucose homeostasis^[25]. Together, these biological responses might contribute to insulin resistance and increase the likelihood of developing T2DM^[25,26].

Among all SDOH domains examined, government or no health insurance emerged as the greatest predictor of T2DM prevalence. Individuals without adequate insurance coverage may face barriers to timely diagnosis, routine screening, and ongoing disease management, potentially contributing to undetected or poorly controlled T2DM^[27]. Lower education level also showed a substantial association^[28], as it influences a wide range of downstream factors—including health literacy, self-management behaviors, preventive care utilization, and the ability to interpret health information—which collectively affect an individual's capacity to maintain metabolic health^[4]. A lower

Table 2. Weighted trends in the prevalence of T2DM pre-, intra-, preliminary post-pandemic (weighted % [95% CI]) based on data obtained from the KNHANES

Group	Pre-pandemic				Intra-pandemic			Preliminary post-pandemic	
	2007–2009	2010–2012	2013–2015	2016–2019	2020	2021	2022	2023	2024
Overall	8.55 (7.96 to 9.13)	9.69 (9.08 to 10.29)	10.97 (10.31 to 11.63)	11.66 (11.11 to 12.22)	13.51 (12.23 to 14.80)	13.66 (12.24 to 15.07)	12.69 (11.64 to 13.74)	12.55 (11.44 to 13.66)	14.19 (12.99 to 15.39)
Sex									
Male	9.49 (8.59 to 10.39)	10.73 (9.82 to 11.64)	12.33 (11.37 to 13.29)	13.17 (12.35 to 13.98)	15.79 (13.86 to 17.72)	15.85 (13.77 to 17.93)	15.23 (13.56 to 16.90)	15.12 (13.50 to 16.73)	16.55 (14.90 to 18.20)
Female	7.64 (6.97 to 8.32)	8.67 (7.93 to 9.41)	9.67 (8.91 to 10.43)	10.19 (9.52 to 10.85)	11.24 (9.82 to 12.67)	11.49 (9.91 to 13.07)	10.16 (8.85 to 11.46)	10.05 (8.83 to 11.26)	11.81 (10.35 to 13.27)
Age group (years)									
19–	1.87 (1.39 to 2.35)	2.02 (1.51 to 2.53)	1.71 (1.26 to 2.16)	2.09 (1.68 to 2.51)	2.19 (1.27 to 3.10)	1.87 (0.82 to 2.91)	2.23 (1.27 to 3.20)	2.18 (1.24 to 3.11)	2.70 (1.51 to 3.90)
40–	9.52 (8.56 to 10.49)	10.60 (9.58 to 11.61)	11.14 (10.13 to 12.16)	11.35 (10.51 to 12.19)	14.07 (11.92 to 16.23)	12.72 (10.72 to 14.72)	13.12 (11.13 to 15.11)	11.67 (10.08 to 13.26)	12.23 (10.84 to 13.62)
60–	22.84 (20.78 to 24.91)	22.09 (20.28 to 23.91)	25.89 (23.74 to 28.04)	22.55 (20.98 to 24.13)	26.94 (23.02 to 30.85)	25.68 (22.48 to 28.89)	21.84 (18.94 to 24.74)	23.56 (20.92 to 26.20)	26.21 (23.32 to 29.11)
≥ 70	20.23 (17.85 to 22.61)	24.65 (22.52 to 26.77)	29.23 (26.90 to 31.56)	30.05 (28.24 to 31.85)	30.87 (26.85 to 34.90)	33.34 (29.45 to 37.23)	26.71 (23.68 to 29.74)	27.13 (24.04 to 30.23)	30.78 (27.58 to 33.98)
Region of residence									
Urban	8.67 (7.79 to 9.55)	8.12 (7.32 to 8.92)	10.99 (9.99 to 11.98)	11.20 (10.39 to 12.00)	12.06 (10.02 to 14.11)	12.39 (10.27 to 14.51)	11.81 (10.28 to 13.33)	11.81 (10.30 to 13.33)	13.76 (11.83 to 15.68)
Rural	8.44 (7.66 to 9.21)	11.15 (10.26 to 12.05)	10.96 (10.07 to 11.84)	12.12 (11.33 to 12.90)	14.78 (13.17 to 16.39)	14.84 (12.93 to 16.75)	13.38 (11.93 to 14.83)	13.16 (11.56 to 14.77)	14.55 (13.04 to 16.06)
BMI group*									
Underweight	2.66 (1.42 to 3.91)	3.51 (1.96 to 5.06)	2.68 (1.34 to 4.02)	3.10 (1.72 to 4.49)	2.92 (0.00 to 6.73)	2.19 (0.70 to 3.68)	4.61 (1.38 to 7.84)	7.05 (3.66 to 10.44)	3.29 (0.67 to 5.90)
Normal	4.90 (4.22 to 5.58)	6.39 (5.66 to 7.12)	7.25 (6.46 to 8.04)	6.98 (6.37 to 7.59)	8.07 (6.48 to 9.66)	9.39 (7.60 to 11.18)	8.49 (7.16 to 9.83)	7.71 (6.45 to 8.97)	8.56 (7.19 to 9.94)
Overweight	8.26 (7.21 to 9.30)	9.55 (8.41 to 10.68)	11.62 (10.40 to 12.85)	11.70 (10.70 to 12.70)	14.71 (12.06 to 17.36)	12.20 (9.93 to 14.46)	11.37 (9.39 to 13.35)	13.20 (11.22 to 15.17)	14.93 (12.82 to 17.04)
Obese	14.34 (13.08 to 15.59)	14.91 (13.63 to 16.18)	16.22 (14.96 to 17.48)	17.93 (16.90 to 18.96)	18.94 (16.79 to 21.09)	19.92 (17.43 to 22.41)	18.65 (16.50 to 20.80)	17.79 (15.99 to 19.59)	20.26 (18.23 to 22.29)
Smoking status									
Never smoker	7.80 (7.12 to 8.48)	8.35 (7.63 to 9.07)	9.59 (8.82 to 10.35)	10.06 (9.42 to 10.70)	11.51 (10.09 to 12.92)	11.33 (9.75 to 12.91)	10.24 (9.01 to 11.46)	10.55 (9.29 to 11.81)	11.75 (10.29 to 13.22)
Ex-smoker	11.53 (10.08 to 12.97)	12.97 (11.52 to 14.42)	13.74 (12.25 to 15.23)	15.47 (14.30 to 16.64)	16.67 (14.11 to 19.22)	18.22 (15.59 to 20.85)	15.29 (13.05 to 17.52)	15.64 (13.25 to 18.02)	19.03 (16.69 to 21.38)
Current smoker	7.75 (6.65 to 8.85)	9.88 (8.68 to 11.08)	12.04 (10.63 to 13.44)	11.98 (10.78 to 13.18)	15.63 (12.61 to 18.66)	15.05 (11.44 to 18.66)	16.87 (14.05 to 19.68)	14.97 (12.47 to 17.47)	15.60 (12.84 to 18.36)
Drinking status									
Never drinker	13.44 (11.65 to 15.22)	16.59 (14.70 to 18.48)	19.68 (17.46 to 21.90)	21.52 (19.63 to 23.42)	22.66 (17.30 to 28.02)	21.91 (17.75 to 26.07)	21.38 (16.92 to 25.84)	18.78 (15.34 to 22.23)	23.77 (19.44 to 28.09)
Ex-drinker	14.36 (12.68 to 16.05)	13.34 (11.56 to 15.12)	14.12 (12.53 to 15.71)	15.72 (14.28 to 17.15)	17.32 (14.44 to 20.19)	19.23 (16.08 to 22.38)	17.83 (14.91 to 20.75)	18.59 (15.97 to 21.21)	18.51 (15.85 to 21.16)
Current drinker	6.85 (6.23 to 7.47)	8.12 (7.45 to 8.79)	9.24 (8.57 to 9.92)	9.80 (9.22 to 10.39)	11.65 (10.28 to 13.02)	11.15 (9.74 to 12.56)	10.63 (9.42 to 11.83)	10.55 (9.36 to 11.75)	12.31 (11.05 to 13.56)
Blood pressure									
Normal	3.39 (2.89 to 3.89)	4.05 (3.51 to 4.60)	4.59 (4.01 to 5.18)	4.86 (4.35 to 5.37)	5.86 (4.53 to 7.19)	5.72 (4.49 to 6.95)	5.30 (4.33 to 6.28)	5.05 (4.15 to 5.95)	5.75 (4.68 to 6.82)
Elevated	10.01 (7.48 to 12.55)	10.04 (7.70 to 12.38)	13.37 (10.88 to 15.86)	14.60 (12.34 to 16.85)	14.35 (10.28 to 18.42)	11.69 (8.06 to 15.32)	9.25 (6.37 to 12.13)	11.71 (8.51 to 14.90)	11.31 (8.20 to 14.42)
Stage 1 hypertension	7.54 (6.22 to 8.86)	8.38 (7.11 to 9.65)	10.12 (8.70 to 11.54)	8.13 (7.17 to 9.10)	10.94 (8.70 to 13.19)	11.82 (9.03 to 14.62)	12.42 (9.24 to 15.61)	10.50 (8.32 to 12.68)	10.31 (8.11 to 12.51)
Stage 2 hypertension	19.34 (17.87 to 20.80)	20.34 (18.99 to 21.70)	24.35 (22.70 to 25.99)	25.28 (24.02 to 26.54)	28.59 (25.48 to 31.70)	29.30 (26.57 to 32.03)	25.77 (23.17 to 28.38)	27.66 (25.37 to 29.95)	29.96 (27.45 to 32.48)

Continued

Group	Pre-pandemic				Intra-pandemic			Preliminary post-pandemic	
	2007–2009	2010–2012	2013–2015	2016–2019	2020	2021	2022	2023	2024
Physical activity									
Insufficient	9.00 (7.95 to 10.05)	11.32 (10.29 to 12.34)	13.28 (12.28 to 14.29)	13.72 (12.95 to 14.48)	14.77 (12.99 to 16.55)	16.79 (14.91 to 18.67)	15.43 (13.79 to 17.07)	15.29 (13.89 to 16.69)	17.51 (15.90 to 19.13)
Sufficient	8.40 (7.75 to 9.05)	8.96 (8.24 to 9.69)	9.42 (8.64 to 10.21)	9.34 (8.66 to 10.03)	11.96 (10.27 to 13.65)	9.86 (8.24 to 11.48)	9.95 (8.61 to 11.28)	9.75 (8.34 to 11.15)	10.54 (8.95 to 12.13)
Education level									
High	6.20 (5.65 to 6.75)	6.02 (5.43 to 6.60)	7.44 (6.82 to 8.06)	8.21 (7.70 to 8.73)	10.47 (9.29 to 11.64)	10.34 (9.04 to 11.65)	10.13 (9.05 to 11.21)	9.85 (8.73 to 10.96)	11.13 (10.07 to 12.19)
Low	19.02 (17.41 to 20.64)	19.64 (18.32 to 20.96)	22.80 (21.23 to 24.38)	25.31 (23.97 to 26.65)	28.58 (25.25 to 31.90)	29.08 (26.03 to 32.13)	25.57 (22.61 to 28.54)	26.02 (23.38 to 28.66)	28.73 (25.65 to 31.81)
Marital status									
Married	9.60 (8.85 to 10.35)	10.65 (9.91 to 11.39)	11.94 (11.15 to 12.74)	12.85 (12.17 to 13.53)	15.50 (13.76 to 17.25)	15.36 (13.46 to 17.26)	14.03 (12.62 to 15.45)	14.39 (12.88 to 15.89)	16.57 (15.07 to 18.07)
Single, separated, widowed, or divorced	6.11 (5.28 to 6.94)	7.59 (6.68 to 8.50)	8.94 (7.94 to 9.93)	9.34 (8.51 to 10.17)	10.00 (8.40 to 11.61)	10.72 (8.94 to 12.49)	10.33 (8.63 to 12.03)	9.41 (7.89 to 10.94)	10.21 (8.50 to 11.93)
Household income									
High	6.22 (5.57 to 6.87)	7.49 (6.75 to 8.23)	7.89 (7.21 to 8.57)	8.47 (7.89 to 9.05)	10.95 (9.55 to 12.35)	9.28 (7.91 to 10.65)	9.70 (8.52 to 10.89)	10.10 (8.86 to 11.34)	11.99 (10.73 to 13.26)
Low	11.88 (10.90 to 12.86)	12.60 (11.72 to 13.48)	15.79 (14.67 to 16.90)	16.80 (15.87 to 17.74)	18.33 (16.10 to 20.55)	21.58 (19.33 to 23.83)	17.70 (15.84 to 19.56)	16.84 (15.05 to 18.63)	18.00 (15.61 to 20.39)
Food security									
Full security	7.45 (6.68 to 8.22)	8.85 (7.96 to 9.75)	10.14 (9.30 to 10.97)	10.57 (9.88 to 11.25)	12.85 (11.29 to 14.41)	12.43 (10.74 to 14.11)	11.46 (10.34 to 12.58)	11.57 (10.34 to 12.81)	13.43 (12.25 to 14.62)
Marginal, low, or very low security	9.33 (8.55 to 10.12)	10.32 (9.52 to 11.12)	11.78 (10.85 to 12.71)	13.11 (12.29 to 13.93)	14.74 (12.81 to 16.67)	16.15 (14.19 to 18.11)	15.66 (13.25 to 18.07)	15.01 (13.05 to 16.97)	17.21 (14.56 to 19.86)
Type of health insurance									
Private	6.29 (5.73 to 6.86)	7.01 (6.39 to 7.63)	8.34 (7.69 to 8.99)	9.41 (8.87 to 9.95)	11.91 (10.73 to 13.08)	11.36 (10.02 to 12.69)	11.37 (10.26 to 12.49)	11.27 (10.13 to 12.42)	12.93 (11.72 to 14.14)
Government or none	14.41 (13.09 to 15.74)	17.75 (16.34 to 19.17)	21.29 (19.59 to 22.98)	22.34 (20.88 to 23.80)	23.48 (20.01 to 26.95)	25.39 (21.97 to 28.82)	21.00 (18.12 to 23.87)	21.06 (18.26 to 23.85)	23.89 (20.40 to 27.38)
Employment status									
Employed, student, or retired	7.15 (6.42 to 7.88)	8.34 (7.59 to 9.09)	9.14 (8.40 to 9.89)	10.01 (9.39 to 10.63)	11.82 (10.35 to 13.28)	11.57 (10.10 to 13.04)	10.67 (9.28 to 12.06)	11.47 (10.29 to 12.65)	12.52 (11.28 to 13.76)
Unemployed	10.68 (9.71 to 11.65)	12.10 (11.14 to 13.05)	14.04 (12.92 to 15.15)	14.58 (13.63 to 15.53)	16.40 (14.43 to 18.37)	17.31 (14.96 to 19.65)	16.42 (14.59 to 18.25)	14.71 (12.87 to 16.55)	17.64 (15.39 to 19.90)
Home ownership									
Own home	9.22 (8.50 to 9.93)	10.23 (9.52 to 10.95)	11.63 (10.86 to 12.40)	11.99 (11.35 to 12.63)	14.31 (12.84 to 15.78)	13.07 (11.34 to 14.80)	12.72 (11.40 to 14.04)	13.50 (12.16 to 14.84)	14.80 (13.51 to 16.10)
Rent home or other arrangement	7.24 (6.30 to 8.18)	8.49 (7.46 to 9.51)	9.71 (8.66 to 10.75)	10.99 (10.02 to 11.96)	11.86 (9.68 to 14.04)	14.82 (12.74 to 16.90)	12.62 (10.82 to 14.41)	10.48 (8.78 to 12.19)	12.72 (10.50 to 14.95)
SDOH burden score									
0	7.02 (5.34 to 8.70)	7.83 (5.89 to 9.78)	7.74 (6.02 to 9.46)	9.04 (7.77 to 10.30)	12.98 (9.81 to 16.15)	9.28 (6.49 to 12.07)	9.49 (6.85 to 12.13)	11.22 (9.10 to 13.33)	13.25 (11.03 to 15.48)
1	6.64 (5.58 to 7.71)	6.55 (5.53 to 7.58)	7.88 (6.72 to 9.04)	8.78 (7.84 to 9.73)	11.36 (9.27 to 13.45)	9.68 (7.22 to 12.15)	9.49 (7.62 to 11.35)	9.67 (7.67 to 11.66)	11.99 (10.09 to 13.89)
2	5.70 (4.75 to 6.65)	6.98 (5.96 to 8.01)	8.17 (7.13 to 9.20)	8.86 (7.96 to 9.75)	9.89 (7.51 to 12.28)	10.71 (8.46 to 12.95)	12.31 (10.03 to 14.59)	11.91 (9.90 to 13.91)	10.94 (8.92 to 12.97)
3	8.12 (6.86 to 9.38)	8.91 (7.65 to 10.17)	10.33 (9.08 to 11.59)	10.93 (9.77 to 12.09)	13.85 (10.74 to 16.96)	16.21 (12.96 to 19.46)	14.19 (11.43 to 16.94)	12.72 (10.50 to 14.94)	15.49 (12.34 to 18.65)

Continued

Group	Pre-pandemic				Intra-pandemic			Preliminary post-pandemic	
	2007–2009	2010–2012	2013–2015	2016–2019	2020	2021	2022	2023	2024
4	10.42 (8.81 to 12.04)	13.27 (11.59 to 14.94)	15.24 (13.27 to 17.22)	16.67 (15.01 to 18.33)	18.24 (14.19 to 22.28)	19.25 (15.44 to 23.06)	14.04 (10.42 to 17.65)	16.44 (13.26 to 19.62)	21.37 (16.81 to 25.93)
5	14.72 (12.34 to 17.10)	16.01 (13.59 to 18.43)	21.30 (18.45 to 24.15)	21.10 (18.89 to 23.32)	20.81 (15.60 to 26.01)	28.97 (22.79 to 35.14)	22.82 (17.39 to 28.25)	23.45 (18.17 to 28.73)	25.17 (18.47 to 31.88)
≥ 6	21.27 (17.97 to 24.58)	24.29 (20.67 to 27.90)	26.12 (22.61 to 29.62)	30.81 (27.64 to 33.98)	34.48 (26.29 to 42.66)	30.01 (23.53 to 36.49)	38.56 (30.75 to 46.37)	24.82 (18.12 to 31.52)	25.81 (17.83 to 33.79)
Cumulative unfavorable SDOH									
Low-burden (≤ 2)	6.29 (5.62 to 6.97)	6.96 (6.23 to 7.69)	7.98 (7.29 to 8.67)	8.87 (8.28 to 9.46)	11.23 (9.78 to 12.69)	9.94 (8.48 to 11.41)	10.48 (9.26 to 11.70)	10.87 (9.61 to 12.14)	12.03 (10.87 to 13.19)
High-burden (> 2)	11.28 (10.31 to 12.25)	12.98 (12.07 to 13.90)	15.14 (14.07 to 16.22)	16.18 (15.24 to 17.12)	17.82 (15.70 to 19.94)	20.50 (18.28 to 22.72)	17.06 (15.11 to 19.01)	16.09 (14.27 to 17.92)	19.28 (16.56 to 21.99)

Note. *According to Asian-Pacific guidelines, BMI is divided into four groups: underweight (< 18.5 kg/m²), normal weight (18.5–22.9 kg/m²), overweight (23.0–24.9 kg/m²), and obese (≥ 25.0 kg/m²). BMI, body mass index; *CI*: confidence interval; KNHANES, Korea National Health and Nutrition Examination Survey; SDOH, social determinants of health; T2DM, type 2 diabetes mellitus.

education level is also closely tied to broader structural disadvantages—such as unstable employment, food insecurity, lack of health insurance coverage, and limited access to environments that support healthy dietary patterns or physical activity—which may help explain the higher T2DM prevalence observed in the high-burden group^[29].

The pandemic may have intensified these disparities through a diagnostic delay followed by an increase during the preliminary post-pandemic period. In our data, T2DM prevalence increased through 2021, showed a modest decline in 2022–2023, and rose again in 2024, a pattern that may reflect delayed detection followed by accumulated metabolic burden. This trend may reflect not only accumulated metabolic debt but also the virus’s potential effect, where direct pancreatic injury and persistent insulin resistance can facilitate new-onset T2DM^[12,30]. These biological risks, combined with limited access to technology-based care, potentially imposed a disproportionate burden populations with unfavorable SDOH, leading to the long-term metabolic consequences observed in our study^[12,13].

Comparison to Previous Studies

Contrary to previous small-cohort studies, our study analyzed a large, nationally representative sample of 81,065 adults, ensuring sufficient statistical power and generalizability (*n*=60,098, 2011–2020^[31]; *n*=79,753, 1998–2018^[32] vs. *n*=85,695, 2007–2024). A key strength is our extended timeframe, which captures the entire trajectory of

the COVID-19 pandemic up to 2024, including the preliminary post-pandemic period. This allowed for a more nuanced interpretation of the pandemic’s long-term consequences on T2DM management, distinguishing immediate shocks from persistent effects, in contrast to prior studies limited to the pandemic’s initial phase (2019–2022^[6], 2011–2022^[33] vs. 2007–2024). Furthermore, this is one of the first long-term studies in South Korea to investigate the




Figure 2. Cumulative SDOH-specific trends in T2DM prevalence during pre-, intra-, preliminary post-pandemic periods from 2007 to 2024, and shaded regions indicate 95% *CIs*. Groups were categorized based on the cumulative count of adverse SDOH factors: low burden (≤ 2) and high burden (> 2). Abbreviations: *CIs*, confidence intervals; SDOH, social determinants of health; T2DM, type 2 diabetes mellitus.

Continued

Variables	Overall (2007–2024)		Pre-pandemic (2007–2019)		Intra-pandemic (2020–2022)		Preliminary post-pandemic (2023–2024)	
	aOR (95% CI) ^a	P-value	aOR (95% CI)	P-value	aOR (95% CI)	P-value	aOR (95% CI)	P-value
High	1.00 (ref)		1.00 (ref)		1.00 (ref)		1.00 (ref)	
Low	1.25 (1.17 to 1.33)	< 0.001	1.33 (1.23 to 1.43)	< 0.001	1.28 (1.16 to 1.41)	< 0.001	1.17 (1.02 to 1.34)	0.027
Marital status								
Married	1.00 (ref)		1.00 (ref)		1.00 (ref)		1.00 (ref)	
Single, separated, widowed, or divorced	1.09 (1.03 to 1.16)	0.005	1.04 (0.97 to 1.13)	0.293	1.17 (1.06 to 1.28)	0.001	1.14 (1.01 to 1.30)	0.042
Household income								
High	1.00 (ref)		1.00 (ref)		1.00 (ref)		1.00 (ref)	
Low	1.25 (1.18 to 1.32)	< 0.001	1.30 (1.21 to 1.39)	< 0.001	1.34 (1.23 to 1.47)	< 0.001	1.12 (0.98 to 1.27)	0.090
Food security								
Full security	1.00 (ref)		1.00 (ref)		1.00 (ref)		1.00 (ref)	
Marginal, low, or very low security	1.05 (0.99 to 1.10)	0.083	1.07 (1.00 to 1.14)	0.045	1.06 (0.98 to 1.15)	0.122	1.15 (1.02 to 1.30)	0.028
Type of health insurance								
Private	1.00 (ref)		1.00 (ref)		1.00 (ref)		1.00 (ref)	
Government or none	1.30 (1.22 to 1.39)	< 0.001	1.40 (1.29 to 1.51)	< 0.001	1.31 (1.19 to 1.44)	< 0.001	1.10 (0.96 to 1.27)	0.173
Employment status								
Employed, student, or retired	1.00 (ref)		1.00 (ref)		1.00 (ref)		1.00 (ref)	
Unemployed	1.25 (1.18 to 1.33)	< 0.001	1.27 (1.18 to 1.36)	< 0.001	1.26 (1.15 to 1.38)	< 0.001	1.17 (1.04 to 1.33)	0.012
Home ownership								
Own home	1.00 (ref)		1.00 (ref)		1.00 (ref)		1.00 (ref)	
Rent home or other arrangement	1.14 (1.08 to 1.21)	< 0.001	1.11 (1.03 to 1.19)	0.005	1.25 (1.14 to 1.36)	< 0.001	1.16 (1.02 to 1.31)	0.022
SDOH burden score								
0	1.00 (ref)		1.00 (ref)		1.00 (ref)		1.00 (ref)	
1	1.08 (0.98 to 1.19)	0.141	1.08 (0.95 to 1.24)	0.244	1.15 (0.99 to 1.34)	0.062	1.11 (0.92 to 1.34)	0.272
2	1.13 (1.02 to 1.26)	0.019	1.13 (0.99 to 1.30)	0.074	1.26 (1.08 to 1.47)	0.004	1.27 (1.04 to 1.55)	0.021
3	1.30 (1.17 to 1.45)	< 0.001	1.34 (1.17 to 1.54)	< 0.001	1.43 (1.22 to 1.68)	< 0.001	1.27 (1.03 to 1.56)	0.027
4	1.48 (1.32 to 1.65)	< 0.001	1.59 (1.38 to 1.85)	< 0.001	1.62 (1.37 to 1.93)	< 0.001	1.32 (1.04 to 1.67)	0.022
5	1.62 (1.43 to 1.83)	< 0.001	1.68 (1.43 to 1.96)	< 0.001	1.77 (1.47 to 2.13)	< 0.001	1.68 (1.30 to 2.17)	< 0.001
≥ 6	1.79 (1.58 to 2.04)	< 0.001	1.92 (1.63 to 2.26)	< 0.001	2.26 (1.87 to 2.72)	< 0.001	1.66 (1.24 to 2.23)	< 0.001
Continuous value	1.11 (1.09 to 1.13)	< 0.001	1.12 (1.10 to 1.15)	< 0.001	1.13 (1.10 to 1.16)	< 0.001	1.09 (1.05 to 1.13)	< 0.001
Cumulative unfavorable SDOH								
Low-burden (≤ 2)	1.00 (ref)		1.00 (ref)		1.00 (ref)		1.00 (ref)	
High-burden (> 2)	1.33 (1.25 to 1.41)	< 0.001	1.39 (1.29 to 1.50)	< 0.001	1.38 (1.26 to 1.51)	< 0.001	1.19 (1.05 to 1.35)	0.007

Note. *According to Asian-Pacific guidelines, BMI is divided into four groups: underweight (< 18.5 kg/m²), normal weight (18.5–22.9 kg/m²), overweight (23.0–24.9 kg/m²), and obese (≥ 25.0 kg/m²). The figures in bold indicate statistical significance (*P*-value < 0.05). ^aThis model was adjusted for all sociodemographic variables. aOR, adjusted odds ratio; BMI, body mass index; CI: confidence interval; KNHANES, Korea National Health and Nutrition Examination Survey; SDOH, social determinants of health; T2DM, type 2 diabetes mellitus.

association between a cumulative count of unfavorable SDOH and T2DM prevalence. While previous studies focused on individual socioeconomic factors^[3], our approach using a cumulative SDOH score reveals the synergistic and graded impact of social disadvantages on T2DM risk.

Clinical and Policy Implications

T2DM results from a complex interplay between individual lifestyle factors and powerful structural determinants^[5,10]. The profound impact of cumulative burden of social disadvantage on T2DM prevalence requires a paradigm shift from individual behavioral modifications to structural transformations^[34]. From a clinical perspective, lack of health insurance coverage represents a critical access barrier that may be associated with delayed diagnosis and reduced continuity of care. Expanding coverage and reducing out-of-pocket costs for high-risk populations should therefore be prioritized as a frontline clinical strategy^[35]. Lower educational attainment may further be associated with lower health literacy, potentially limiting health literacy and limiting effective engagement with both conventional and digital health interventions^[36,37]. Digital health interventions must move beyond merely expanding technological access and instead embed equity through digital literacy support and user-centered design that accommodates diverse levels of health literacy^[38,39]. At the policy level, addressing upstream drivers like food insecurity and unemployment demands a "Health in All Policies" approach. This includes incorporating health equity impact assessments across non-health sectors such as urban planning and labor market policies^[40,41]. Despite South Korea's near-universal health insurance system, individuals who remain underserved by this infrastructure may face disproportionate barriers to timely diabetes screening and management^[42]. With respect to insurance coverage, South Korea's National Health Insurance Service could consider expanding its targeted outreach programs, such as the Community Health Care Center initiative, to actively identify and enroll uninsured or underinsured individuals in high-risk SDOH burden groups, coupled with waived or reduced copayments for diabetes screening and management^[35]. Furthermore, systemic solutions to obesogenic environments should include fiscal policies, specifically taxation on sugar-sweetened beverages combined with subsidies for fresh produce. These integrated strategies are critical for dismantling the structural roots of T2DM inequalities

in the post-pandemic era.

Strengths and Limitations

This study has several limitations. First, given the multifactorial nature of T2DM and the interrelated structure of the covariates, interpreting the effect of any single variable in isolation requires caution. To mitigate this limitation, we adopted a cumulative risk approach based on constructing an index of unfavorable SDOH. This allowed us to capture the synergistic burden of multiple interconnected social disadvantages, rather than relying solely on the isolated associations of individual variable. Second, because KNHANES data are derived from self-reported questionnaires, they are inherently subject to response bias, and the results may vary depending on the participant's recall. To mitigate these limitations, we attempted to include variables defined by objective measures, such as experience with a diagnosis from a physician, a fasting glucose level, and a hemoglobin A1c level, whenever possible. Third, because this study used only Korean data, the applicability of the results to other ethnic groups or countries may be limited. Therefore, similar follow-up studies using data from other countries are needed. Fourth, the cross-sectional design of our study precludes the establishment of causal relationships between the cumulative burden of unfavorable SDOH and T2DM^[43]. Nevertheless, this design is effective for identifying significant associations and generating hypotheses for future longitudinal research. Fifth, the cross-sectional nature of the data prevented longitudinal follow-up of the same individuals. As a result, direct within-person changes associated with the evolving pandemic context could not be assessed. Nevertheless, this limitation was partially mitigated by the use of nationally representative samples collected with consistent survey methodology across survey years and by adjustment for a comprehensive set of sociodemographic and health-related covariates to enhance comparability between survey cycles. Finally, many of the SDOH variables were measured using categorized or simplified indicators; household income was grouped into quartiles, and food security was a binary measure. Such categorization, while necessary for large-scale survey analysis, may not fully capture the complex and continuous nature of these social constructs, potentially leading to some loss of information.

Despite these limitations, to the best of our knowledge, this study is among the first in South Korea to investigate the association between the

cumulative number of unfavorable SDOH and the prevalence of T2DM using a large, nationally representative sample over an extended period. This approach moves beyond analyzing individual socioeconomic factors in isolation to reveal the synergistic and graded impact of social disadvantages on T2DM risk, providing a more holistic view of health inequality.

CONCLUSION

This study analyzed long-term trends in T2DM prevalence and its association with SDOH in South Korea using nationally representative data from 2007 to 2024. The prevalence of T2DM continued to rise through the intra-pandemic period and was closely associated with a greater cumulative burden of unfavorable SDOH. Among the individual SDOH indicators, government or no health insurance showed the strongest association. These findings highlight the need for proactive, integrated public health strategies that target both individual risk factors and social inequalities to manage the growing T2DM burden.

Funding This research was supported by the Ministry of Science and ICT (RS-2024-00509257 and IITP-2026-RS-2024-00438239) and the Ministry of Health & Welfare (RS-2025-02220492), Republic of Korea. The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Conflict of Interest Statement The authors have no conflicts of interest to declare.

Ethics The research protocol was approved by the Institutional Review Boards of the KDCA (2007-02CON-04-P, 2008-04EXP-01-C, 2009-01CON-03-2C, 2010-02CON-21-C, 2011-02CON-06-C, 2012-01EXP-01-2C, 2013-07CON-03-4C, 2013-12EXP-03-5C, 2018-01-03-P-A, 2018-01-03-C-A, 2018-01-03-2C-A, 2018-01-03-5C-A, 2018-01-03-4C-A, 2022-11-16-R-A, and 2022-11-16-R-03). Written informed consent was obtained from all participants prior to their participation. The study was conducted in accordance with the ethical principles of the Declaration of Helsinki.

Authors' Contributions Dr. Dongkeon Yon had full access to all of the data in the study and took responsibility for the integrity of the data and the accuracy of the data analysis. All authors approved the final version of the manuscript before submission. Study concept and design: Jihu Im, Juyeong Kim, and Dongkeon Yon; acquisition,

analysis, or interpretation of data: Jihu Im, Juyeong Kim, and Dongkeon Yon; drafting of the manuscript: Jihu Im, Juyeong Kim, and Dongkeon Yon; critical revision of the manuscript for important intellectual content: all authors; statistical analysis: Jihu Im, Juyeong Kim, and Dongkeon Yon; study supervision: Dongkeon Yon. Dongkeon Yon supervised the study and served as a guarantor. Jihu Im and Juyeong Kim contributed equally as first authors. Dongkeon Yon contributed as corresponding author. The corresponding author attests that all listed authors meet the authorship criteria and that no one meeting the criteria has been omitted. All authors had full access to the study data and had final responsibility for the decision to submit for publication.

Data Availability Statement Data are publicly available on legal and ethical grounds from the KDCA as open data (<https://www.kdca.go.kr/yhs/home.jsp>).

Received:

Accepted:

REFERENCES

1. Zhong JL, Zhang YB, Zhu K, et al. Associations of social determinants of health with life expectancy and future health risks among individuals with type 2 diabetes: two nationwide cohort studies in the UK and USA. *Lancet Healthy Longev*, 2024; 5, e542–51.
2. Lu X, Xie QX, Pan XH, et al. Type 2 diabetes mellitus in adults: pathogenesis, prevention and therapy. *Signal Transduct Target Ther*, 2024; 9, 262.
3. Park EJ, Ji NJ, You CH, et al. Healthcare utilization and discrepancies by income level among patients with newly diagnosed type 2 diabetes in Korea: an analysis of national health insurance sample cohort data. *J Prev Med Public Health*, 2024; 57, 471–9.
4. Coughlin SS, Vernon M, Hatzigeorgiou C, et al. Health literacy, social determinants of health, and disease prevention and control. *J Environ Health Sci*, 2020; 6, 3061.
5. Hill-Briggs F, Adler NE, Berkowitz SA, et al. Social determinants of health and diabetes: a scientific review. *Diabetes Care*, 2020; 44, 258–79.
6. Kim Y, Kim B, Rajaguru V, et al. Association between diabetes mellitus education and self-management pre- and post-COVID-19. *Am J Health Promot*, 2025; 39, 777–85.
7. Green H, Fernandez R, MacPhail C. The social determinants of health and health outcomes among adults during the COVID-19 pandemic: a systematic review. *Public Health Nurs*, 2021; 38, 942–52.
8. Odoms-Young A, Brown AGM, Agurs-Collins T, et al. Food insecurity, neighborhood food environment, and health disparities: state of the science, research gaps and opportunities. *Am J Clin Nutr*, 2024; 119, 850–61.
9. Zhao Z, Li L, Sang Y. The COVID-19 pandemic increased poor lifestyles and worsen mental health: a systematic review. *Am J Transl Res*, 2023; 15, 3060–6.

10. Park YS, Kang SH, Jang SI, et al. Association between lifestyle factors and the risk of metabolic syndrome in the South Korea. *Sci Rep*, 2022; 12, 13356.
11. Pantea Stoian A, Bica IC, Salmen T, et al. New-onset diabetes mellitus in COVID-19: a scoping review. *Diabetes Ther*, 2024; 15, 33–60.
12. Pranata R, Henrina J, Raffaello WM, et al. Diabetes and COVID-19: the past, the present, and the future. *Metabolism*, 2021; 121, 154814.
13. Kang E, Yun J, Hwang SH, et al. The impact of the COVID-19 pandemic in the healthcare utilization in Korea: analysis of a nationwide survey. *J Infect Public Health*, 2022; 15, 915–21.
14. Kim J, Yim Y, Park S, et al. National trends in asthma prevalence among adolescents in South Korea, 2007-2023: a national representative serial study. *Int Arch Allergy Immunol*, 2025; 186, 967–83.
15. Park S, Kim K, Kim M, et al. Trends in adolescent violence victimization pre-, intra-, and post-COVID-19 pandemic in South Korea, 2012-2023: a nationwide cross-sectional study. *Psychiatry Res*, 2025; 348, 116429.
16. Baek JH, Park YM, Han KD, et al. Comparison of operational definition of type 2 diabetes mellitus based on data from Korean national health insurance service and Korea national health and nutrition examination survey. *Diabetes Metab J*, 2023; 47, 201–10.
17. Frank J, Abel T, Campostrini S, et al. The social determinants of health: time to re-think? *Int J Environ Res Public Health*, 2020; 17, 5856.
18. Bundy JD, Mills KT, He H, et al. Social determinants of health and premature death among adults in the USA from 1999 to 2018: a national cohort study. *Lancet Public Health*, 2023; 8, e422–31.
19. Zhu RX, Wang R, He JL, et al. Prevalence of cardiovascular-kidney-metabolic syndrome stages by social determinants of health. *JAMA Netw Open*, 2024; 7, e2445309.
20. Lee H, Kim M, Woo S, et al. National and regional trends in the prevalence of hypertension in South Korea amid the pandemic, 2009-2022: nationwide study of over 3 million individuals. *JMIR Public Health Surveill*, 2024; 10, e51891.
21. Kim H, Park J, Kang J. National trends in osteoporosis in the general older population and postmenopausal women, to the COVID-19 pandemic by related factors, 2001-2021: a nationwide study in South Korea. *Life Cycle*, 2025; 5, e1.
22. Writing Committee Members, Jones DW, Ferdinand KC, et al. 2020 ACC/AHA/AAP/AABC/ACCP/ACPM/AGS/AMA/ASPC/NM A/PCNA/SGIM guideline for the prevention, detection, evaluation and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association joint committee on clinical practice guidelines. *Circulation*, 2025; 152, e114–218.
23. Strain T, Flaxman S, Guthold R, et al. National, regional, and global trends in insufficient physical activity among adults from 2000 to 2022: a pooled analysis of 507 population-based surveys with 5·7 million participants. *Lancet Global Health*, 2024; 12, e1232–43.
24. Zhong JL, Zhang YB, Zhu K, et al. Associations of social determinants of health with life expectancy and future health risks among individuals with type 2 diabetes: two nationwide cohort studies in the UK and USA. *Lancet Healthy Longev*, 2024; 5, e542–51.
25. Lisco G, Giagulli VA, de Pergola G, et al. Chronic stress as a risk factor for type 2 diabetes: endocrine, metabolic, and immune implications. *Endocr Metab Immune Disord Drug Targets*, 2024; 24, 321–2.
26. Chung J, Kim JH, Lee JY, et al. The association between occupational stress level and health-related productivity loss among Korean employees. *Epidemiol Health*, 2023; 45, e2023009.
27. Shin JI, Wang D, Fernandes G, et al. Trends in receipt of American Diabetes Association guideline-recommended care among U. S. adults with diabetes: NHANES 2005-2018. *Diabetes Care*, 2021; 44, 1300–8.
28. Hill-Briggs F, Adler NE, Berkowitz SA, et al. Social determinants of health and diabetes: a scientific review. *Diabetes Care*, 2020; 44, 258–79.
29. Baek SU, Kim MS, Lim MH, et al. Multidimensional employment precariousness mediates the association between low educational attainment and poor subjective well-being: results from a nationwide cross-sectional study in South Korea. *Scand J Work Environ Health*, 2023; 49, 506–17.
30. Al-Aly Z, Cao B. The enduring effect of the COVID-19 pandemic on diabetes. *Lancet Diabetes Endocrinol*, 2024; 12, 508–10.
31. Lee GB, Kim Y, Park S, et al. Obesity, hypertension, diabetes mellitus, and hypercholesterolemia in Korean adults before and during the COVID-19 pandemic: a special report of the 2020 Korea National Health and Nutrition Examination Survey. *Epidemiol Health*, 2022; 44, e2022041.
32. Kim Y, Nho SJ, Woo G, et al. Trends in the prevalence and management of major metabolic risk factors for chronic disease over 20 years: findings from the 1998-2018 Korea National Health and Nutrition Examination Survey. *Epidemiol Health*, 2021; 43, e2021028.
33. Ha KH, Kim DJ. Changes in the epidemiological landscape of diabetes in South Korea: trends in prevalence, incidence, and healthcare expenditures. *Endocrinol Metab*, 2024; 39, 669–77.
34. Konnyu KJ, Yogasingam S, Lépine J, et al. Quality improvement strategies for diabetes care: effects on outcomes for adults living with diabetes. *Cochrane Database Syst Rev*, 2023; 5, Cd014513.
35. American Diabetes Association Professional Practice Committee for Diabetes. 1. Improving care and promoting health in populations: standards of care in diabetes—2026. *Diabetes Care*, 2026; 49, S13–26.
36. Sacerdote C, Ricceri F, Rolandsson O, et al. Lower educational level is a predictor of incident type 2 diabetes in European countries: the EPIC-InterAct study. *Int J Epidemiol*, 2012; 41, 1162–73.
37. Rahim NE, Flood D, Marcus ME, et al. Diabetes risk and provision of diabetes prevention activities in 44 low-income and middle-income countries: a cross-sectional analysis of nationally representative, individual-level survey data. *Lancet Global Health*, 2023; 11, e1576–86.
38. Shirvani T, Javadi Z, Azimi S, et al. Community-based educational interventions for prevention of type II diabetes: a global systematic review and meta-analysis. *Syst Rev*, 2021; 10, 81.
39. Moschonis G, Siopis G, Jung J, et al. Effectiveness, reach, uptake, and feasibility of digital health interventions for adults with type 2 diabetes: a systematic review and meta-analysis of randomised controlled trials. *Lancet Digit Health*, 2023; 5, e125–43.
40. Tonelli M, Tang KC, Forest PG. Canada needs a “Health in All Policies” action plan now. *CMAJ*, 2020; 192, E61–7.
41. Pineda E, Poelman MP, Aaspöllu A, et al. Policy implementation and priorities to create healthy food environments using the Healthy Food Environment Policy Index (Food-EPI): a pooled level analysis across eleven European countries. *Lancet Reg Health Eur*, 2022; 23, 100522.
42. Jung B, Ha IH. Determining the reasons for unmet healthcare needs in South Korea: a secondary data analysis. *Health Qual Life Outcomes*, 2021; 19, 99.

-
43. Shah S, Cho U, Lee W, et al. Coffee intake and hypertension in Korean adults: results from KNHANES 2012-2016. [Clin Hypertens](#), 2023; 29, 20.