

Original Article



Association between Dietary Live Microbe Intake and Sarcopenia: A Cross-sectional Study in Shanghai

Xiaoyu Sun^{1,2,&}, Zhengyuan Wang^{3,&}, Jiayi Wu^{1,4}, Wei Lu³, Zehuan Shi³, Liping Shen³, Chenyu Shen¹,
Chenchen Hu¹, Xuesong Xiang^{1,#}, Jiajie Zang^{1,3,#}, and Guiju Sun^{2,#}

1. Key Laboratory of Public Nutrition and Health, National Health Commission of the People's Republic of China, National Institute for Nutrition and Health, Chinese Center for Disease Control and Prevention, Beijing 100050, China; 2. Department of Nutrition and Food Hygiene, Key Laboratory of Environmental Medicine Engineering of Ministry of Education, School of Public Health, Southeast University, Nanjing 210009, Jiangsu, China; 3. Shanghai Municipal Center for Disease Control and Prevention, Shanghai 201107, China; 4. Key Laboratory of Precision Nutrition and Food Quality, Key Laboratory of Functional Dairy, Ministry of Education, College of Food Science and Nutritional Engineering, China Agricultural University, Beijing 100083, China

Abstract

Objective This study aimed to investigate the relationship between the dietary intake of live microbes and sarcopenia in adults living in Shanghai.

Methods Data were collected from 1,236 participants aged 50 years or more from the Shanghai Diet and Health Surveillance (SDHS). Dietary data were assessed using a standardized questionnaire. The participants were categorized into the low-, medium-, and high-dietary live microbial groups. Sarcopenia was defined according to the European Working Group on Sarcopenia in Older People (EWGSOP2019).

Results Yogurt and traditional Chinese fermented foods (pickled mustard and fermented tofu) were the primary dietary sources of live microbes in this cohort. Overall, participants without sarcopenia consumed foods with higher levels of live microbes than people with sarcopenia ($P < 0.001$). Following stepwise adjustment for confounders, we found that individuals consuming a higher abundance of live dietary microbes were less likely to develop sarcopenia than individuals consuming a lower abundance of live dietary microbes. In the crude model, the odds ratio and 95% confidence interval for high live microbe intake were 0.46 (0.27, 0.76) ($P < 0.001$) and the adjusted odds ratio (95% confidence interval) was 0.41 (0.22, 0.77) ($P = 0.007$). Subgroup analyses identified no significant interaction effects for sex, age, diabetes, fatty liver disease, or overweight/obesity ($P > 0.05$).

Conclusion A high dietary intake of live microbes may be associated with a lower risk of sarcopenia in the Chinese population.

Key words: Sarcopenia; Live microbe; Cross-sectional study; Shanghai

Biomed Environ Sci, 2026; 39(9): 1119-1129

doi: [10.3967/bes2026.095](https://doi.org/10.3967/bes2026.095)

ISSN: 0895-3988

www.besjournal.com (full text)

CN: 11-2816/Q

Copyright ©2026 by China CDC

[&]These authors contributed equally to this work.

[#]Correspondence should be addressed to Xuesong Xiang, PhD, Tel: 13601219682, E-mail: xiangxs@nih.chinacdc.cn; Jiajie Zang, PhD, Tel: 15821992450, E-mail: zangjiajie@scdc.sh.cn; Guiju Sun, PhD, Tel: 13951928860, E-mail: gjsun@seu.edu.cn

Biographical notes of the first authors: Xiaoyu Sun, Master's degree, majoring in probiotics, and food-medicine homology, E-mail: xysun2000@163.com; Zhengyuan Wang, PhD, majoring in nutrition epidemiology and disease prevention, E-mail: wangzhengyuan@scdc.sh.cn

INTRODUCTION

As global aging accelerates, the proportion of individuals aged 65 years and older is steadily increasing. According to the China Statistical Yearbook 2024^[1], 216.76 million individuals were aged 65 years or older by the end of 2023, representing 15.4% of the population. Shanghai, a city with a high life expectancy, is struggling with population aging. By the end of 2023, it had a population of 4.38 million people aged 65 or over, accounting for 28.9% of the total registered permanent residents of the city^[2], which is much higher than the standard of China's aging cities. A gradual decline in muscle mass and quantity occurs with increasing age. Sarcopenia, a progressive skeletal muscle disorder closely associated with aging, has accordingly become increasingly prevalent^[3]. This deterioration increases the risk of adverse events such as falls, frailty, functional limitations, and even mortality. Notably, a recent prospective cohort study across different cultural contexts found that the combination of sarcopenia and social isolation synergistically elevated the all-cause mortality risk among older Chinese adults^[4]. This rising incidence, combined with population aging, not only increases health risks for older adults, but also imposes a substantial socioeconomic burden on healthcare systems and societies worldwide. A recent meta-analysis of nine studies encompassing 7,656 participants aged 50 years and older from community settings identified an overall sarcopenia prevalence of 14%^[5]. Furthermore, data from the China Health and Retirement Longitudinal Study (CHARLS) revealed a substantial increase in the prevalence of sarcopenia among the Chinese population, from 8.5% in 2015 to 29.6% in 2018^[6]. This marked increase highlights the urgent need for intervention strategies targeting sarcopenia.

Multiple factors may affect sarcopenia progression, among which the gut microbiota plays a particularly significant role^[7]. The human gut microbiota plays a critical role in overall health by modulating numerous physiological systems, including muscular functions. By regulating energy balance and nutrient metabolism, the gut microbiota supports the production of short-chain fatty acids (SCFAs), which may promote muscle repair and sustain muscle function^[8]. Gut dysbiosis may impair these processes, consequently contributing to the development and progression of sarcopenia^[9]. The ingestion of probiotic-rich foods has further been found to affect the composition of the gut

microbiota, and existing studies have linked this association to a lower risk of various diseases^[10]. Additionally, numerous studies have shown that probiotic supplementation regulates the composition and function of the gut microbiota, thereby influencing the gut-muscle axis and contributing to the prevention or treatment of sarcopenia^[11,12]. However, only a limited number of individuals regularly consume probiotics. Most live microbes ingested by individuals over time are derived from their daily food.

In addition to probiotics, live dietary microbes directly affect the gut microbiota^[13]. Sanders et al^[14] first used cross-sectional data from the NHANES and Nutrition Examination Survey to categorize foods into three groups. Several studies^[15,16] have since utilized Sanders' classification system and data from the NHANES database to investigate the relationship between the consumption of live microbes and various health outcomes, including sarcopenia, offering invaluable insights into the potential health effects of live microbes.

Although one study used data from the NHANES database to investigate the relationship between live microbial intake and sarcopenia, significant research gaps remain in this field, particularly when considering the substantial variations in dietary patterns, lifestyle habits, and population characteristics across different countries and regions. No studies have yet been conducted on the relationship between live dietary microbes and sarcopenia in the Chinese population. Therefore, in the present study, we conducted surveys on the dietary intake and incidence of sarcopenia to investigate the association between the consumption of live microbes and sarcopenia using a classification system for live microbes in Chinese foods (CLMCS).

MATERIALS AND METHODS

Study Design and Methodology

Data were obtained from the Shanghai Diet and Health Surveillance (SDHS), a representative cross-sectional study conducted by the Shanghai Municipal Centre for Disease Control and Prevention. The study population comprised adults aged 18 years and older who had resided in the city for at least six months. A multistage stratified random sampling method was used. First, 48 surveillance points across the city were selected using a population proportional probability sampling method. Second, one to three

neighborhood/administrative villages were selected from the towns/streets using a simple random sampling method according to the proportion of the population. A total of 71 neighbourhood/administrative villages were selected. A simple random sampling method was applied to select 40 residents from each community and administrative village. Participants were selected from the following age groups, with each group comprising 5 males and 5 females: 18 to 44 years old, 45 to 59 years old, 60 to 74 years old, and 75 years old and above. As sarcopenia is now diagnosed at a younger age, many studies have explored the risk factors for sarcopenia in people aged 50 years and older^[17]. Thus, the present study included all participants aged 50 and above from the SDHS.

Ethics Approval and Consent to Participate

This study was approved by the Ethics Committee of the Shanghai Center for Disease Control and Prevention (KY-2024-16) and was conducted in accordance with the Declaration of Helsinki. Written informed consent was obtained from all the participants before their inclusion.

Data Collection

A standardized questionnaire was used to collect information on basic characteristics, personal health, behavioral habits, and diet. Trained investigators collected the data in a one-to-one manner. A 24-hour dietary recall was conducted over three consecutive days, with each participant reporting all foods consumed over the last 24 h, including two weekdays and one weekend. Information collected using the questionnaire was as follows: whether all consumed foods were pre-packaged foods, names of the raw materials, quality of consumption, and primary cooking methods.

Uniformly trained investigators measured each participant's physical variables, including height, weight, and blood pressure, using standard methods. All tools used for the physical examination were tested and qualified for use in the program. Blood pressure was measured using an electronic sphygmomanometer, with a precision of 1 mmHg. Measurements were conducted three times, at one minute apart.

Collection of Blood Samples and Detection of Indicators

In total, 6 mL of venous blood was collected from each participant after 10–14 h of fasting. The samples were stored in a refrigerator immediately

after collection and the test was completed within 8 h. Hematological indicators included fasting blood glucose (FBG) (hexokinase method), serum triglycerides (TG) (enzymatic method), high-density lipoprotein cholesterol (HDL-C) (direct method-catalase clearance method), and glycosylated hemoglobin (HbA1c) (high-performance liquid chromatography method). FBG, TG, and HDL-C were all measured using a Hitachi 008α instrument, while HbA1c was measured using a Tosoh G11 instrument.

Measurement of Dietary Live Microbe Intake

Dietary intake data were derived from face-to-face 24-hour dietary recall interviews. Participants provided comprehensive details regarding all of their consumed foods and beverages. Foods containing live microbes were classified using the CLMCS, which is based on the Chinese Food Composition Table. Food samples were categorized into low (< 10 CFU/g), medium (10–10 CFU/g), and high (> 10 CFU/g) microbial groups. This method is consistent with the classification criteria proposed by Professor Sanders^[3]. Most foods ($n = 2,303$) were classified as low-level foods, including commonly consumed items such as pork and noodles. Medium-level foods ($n = 173$) predominantly included fresh fruits consumed with the skin and fermented condiments, such as apples and fermented bean paste. High-level foods ($n = 47$) consisted primarily of fermented products, including yogurt and milk skin (nai pi zi). The participants were subsequently categorized into three dietary groups: low dietary intake of live microbes (exclusive consumption of low-level foods), medium intake of dietary live microbes (consumption of medium-level foods without high-level foods), and high intake of dietary live microbes (consumption of any high-level foods).

Diagnosis of Sarcopenia

Sarcopenia was diagnosed based on the 2019 criteria established by the European Working Group on Sarcopenia in Older People (EWGSOP2019) using a three-step algorithm^[18]. First, participants were screened using the SARC-F questionnaire, with a score of ≥ 4 indicating potential sarcopenia. Second, muscle strength and physical performance were evaluated. Low muscle strength was defined as a grip strength < 27 kg for males and < 16 kg for females. Impaired physical performance was defined as a gait speed of ≤ 0.8 m/s. Finally, the presence of low muscle mass was confirmed by bioelectrical impedance analysis (BIA) using a body composition analyzer (Inbody270).

Covariates

A questionnaire survey was conducted to collect demographic data, including age and sex. Educational level was classified into three categories: below high school, high school or equivalent, and above high school. Marital status was categorized into three groups: married, unmarried, and other (widowed or divorced). Annual income was categorized as $\leq 49,000$ CNY, 50,000–199,900 CNY, and $\geq 200,000$ CNY. Body mass index (BMI) was calculated as weight (kg) divided by height squared (m^2). BMI values of less than 18.5 kg/m^2 , between 18.5 kg/m^2 and 24 kg/m^2 , from 24 kg/m^2 to less than 28 kg/m^2 , and 28 kg/m^2 or higher were defined as underweight, normal, overweight, and obese, respectively. Physical activity levels were categorized as insufficient (< 500 metabolic equivalent task (MET) minutes/week) or moderate to high (> 500 MET-minutes per week)^[19]. Hypertension was defined as blood pressure $\geq 140/90$ mmHg, measured more than three times on different days. Subjects with fasting plasma glucose (FPG) ≥ 7.0 mmol/L were defined as having diabetes. The prevalence of fatty liver disease was also self-reported.

Statistical Analysis

All analyses were conducted using R (version 4.4.2), with p -values less than 0.05 considered statistically significant. Categorical variables are summarized as the frequencies and percentages (%), while continuous variables are reported as means and standard errors (SE). The chi-square test was applied to compare categorical variables, and the t -test was used to compare continuous variables. Univariate analyses were conducted to identify the variables with significant differences between the sarcopenic and non-sarcopenic groups.

The relationship between live dietary microbial intake and sarcopenia was investigated using binary logistic regression analysis. To investigate whether this association was modified by baseline characteristics, the interaction terms between the dietary live microbe group and each stratifying variable (sex, age, diabetes, fatty liver disease, and overweight/obesity) were introduced separately into the fully adjusted logistic regression model. The statistical significance of these interactions was determined by evaluating the p -value of the corresponding interaction terms. Stratified analyses were subsequently conducted to determine specific effect sizes within each subgroup.

RESULTS

Clinical Characteristics of the Participants

Among the 1,293 initially enrolled participants, 57 were excluded due to incomplete or invalid nutritional data (Figure 1). In total, 1,236 participants were enrolled in this study. The prevalence of sarcopenia in this population was 18.28%. The baseline characteristics of the participants are summarized in Table 1. Significant differences in terms of age, sex, educational level, TC level, diabetes, fatty liver, and BMI were observed between the sarcopenic and non-sarcopenic individuals. Advanced age ($P < 0.001$) and diabetes mellitus ($P = 0.002$) were identified as the risk factors for sarcopenia. Females ($P = 0.020$) and individuals with high educational levels ($P = 0.006$) exhibited a lower risk of sarcopenia. Importantly, a high dietary intake of live microbes was found to be associated with a lower risk of sarcopenia ($P < 0.001$). Additionally, individuals with sarcopenia exhibited lower BMI ($P < 0.001$), TC levels ($P = 0.006$), and a reduced risk of fatty liver disease ($P = 0.005$). The relationship between sarcopenia and nutrient intake is shown in Table 2. People without sarcopenia consumed more zinc than those with sarcopenia ($P = 0.041$).

Dietary Sources of Live Microbes

To further elucidate the specific dietary patterns contributing to live microbe intake in this cohort, we analyzed both the consumption prevalence (Figure 2A) and average daily intake among consumers (Figure 2B) for the medium and high live microbial food categories. Yogurt was the most widely consumed live microbial source, with a prevalence of 12.70% and an average intake of 81.73 g/day among consumers. Notably, traditional Chinese fermented foods demonstrated substantial penetration in the older population, with pickled mustard (11.33%) and




Figure 1. Flowchart illustrating the sample selection from SDHS.

Table 1. Baseline characteristics of participants, stratified by sarcopenia status

Characteristic	Total	Sarcopenia		P value
		No (1,010)	Yes (226)	
Age (years)	67.18 ± 9.93	66.15 ± 9.65	71.77 ± 9.91	< 0.001
Gender (%)				0.020
Male	592 (47.90)	468 (46.34)	124 (54.87)	
Female	644 (52.10)	542 (53.66)	102 (45.13)	
Education level (%)				0.006
Low	284 (22.98)	214 (21.19)	70 (30.97)	
Medium	803 (64.97)	670 (66.34)	133 (58.85)	
High	149 (12.05)	126 (12.48)	23 (10.18)	
Marital status (%)				0.099
Unmarried	26 (2.10)	19 (1.88)	7 (3.10)	
Married	981 (79.37)	813 (80.50)	168 (74.34)	
Other	229 (18.53)	178 (17.62)	51 (22.56)	
Annual income (%)				0.288
≤ 49,000 CNY	200 (16.18)	156 (15.45)	44 (19.47)	
50,000 – 199,900 CNY	839 (67.88)	689 (68.22)	150 (66.37)	
≥ 200,000 CNY	197 (15.94)	165 (16.34)	32 (14.16)	
BMI(%)				< 0.001
< 18.5	31 (2.51)	6 (0.59)	25 (11.06)	
18.5 ≤ BMI < 24.0	527 (42.64)	369 (36.53)	158 (69.91)	
24.0 ≤ BMI < 28.0	506 (40.94)	463 (45.84)	43 (19.03)	
BMI ≥ 28.0	172 (13.91)	172 (17.03)	0 (0.00)	
TG (mmol/L)	1.67 ± 1.13	1.71 ± 1.16	1.48 ± 0.95	0.986
TC (mmol/L)	5.14 ± 1.03	5.14 ± 1.04	5.13 ± 0.99	0.006
Smoking status (%)				0.506
No	881 (71.28)	724 (71.68)	157 (69.47)	
Yes	355 (28.32)	286 (28.32)	69 (30.57)	
Alcohol consumption (%)				0.954
No	806 (65.21)	659 (65.25)	147 (65.04)	
Yes	430 (34.79)	351 (34.75)	79 (34.96)	
PA (%)				0.500
< 500 MET-min/week	576 (46.60)	463 (45.84)	113 (50.00)	
≥ 500 MET-min/week	660 (53.40)	547 (54.16)	113 (50.00)	
Dietary live microbe group (%)				< 0.001
Low	442 (35.76)	337(33.37)	105 (46.46)	
Medium	633(51.21)	532(52.67)	101 (44.69)	
High	161 (13.03)	141(13.96)	20 (8.85)	
Hypertension (%)				0.183
No	689 (55.74)	572(56.63)	117 (51.77)	
Yes	547 (44.26)	438(43.37)	109 (48.23)	
Diabetes (%)				0.002
No	1,086 (87.86)	901 (89.21)	185(81.86)	
Yes	150 (12.14)	109 (10.79)	41 (18.14)	
Fatty liver (%)				0.005
No	989 (80.02)	793 (78.51)	196 (86.73)	
Yes	247 (19.98)	217 (21.49)	30 (13.27)	

fermented tofu (8.82%) classified as the second- and third-most prevalent sources, respectively. When examining the intake volume among actual consumers, items typically consumed as side dishes or condiments showed realistic portion sizes, such as pickled cabbage (45.00 g/day), fermented rice (28.72 g/day), and fermented tofu (8.53 g/day, equivalent to approximately one standard cube). Consistent with traditional dietary habits, the consumption of typical Western fermented foods, such as cheese, was exceptionally rare (0.57%).

Association between Dietary Live Microbes and Sarcopenia

The association between live microbe intake and

the risk of sarcopenia was investigated using both unadjusted and adjusted logistic regression analyses (Table 3). In the crude model, both medium dietary live microbe intake ($OR = 0.61$, 95% CI : 0.45-0.83) and high dietary live microbe intake ($OR = 0.46$, 95% CI : 0.27-0.76) groups showed significantly lower risk of sarcopenia compared to the low intake group. The protective effects observed for both intake levels remained significant in Model 1, which was adjusted for age and sex ($P < 0.001$). Model 2 was further adjusted for educational level, marital status, and BMI ($P = 0.007$). Model 3 was further adjusted for chronic diseases (diabetes and fatty liver), and showed no substantial changes in the results ($P = 0.008$). In the fully adjusted Model 4, which was

Table 2. Intake of dietary nutrients of participants classified by sarcopenia status

Nutrient	Total	Sarcopenia		P value
		No	Yes	
Energy (kcal)	1705.76 ± 647.77	1718.89 ± 662.05	1647.08 ± 577.37	0.132
Protein (g)	74.77 ± 36.75	75.59 ± 37.34	71.08 ± 33.79	0.095
Fat (g)	68.97 ± 37.31	69.11 ± 36.68	68.31 ± 40.1	0.769
Cholesterol (mg)	498.35 ± 442.28	501.06 ± 431.8	486.22 ± 487.22	0.649
Carbohydrate (g)	197.99 ± 88.31	200.01 ± 91.25	188.94 ± 73.31	0.088
Dietary Fiber (g)	8.14 ± 8.40	8.32 ± 8.88	7.35 ± 5.74	0.115
Calcium (mg)	531.47 ± 408.22	538.2 ± 415.41	501.39 ± 373.8	0.221
Phosphorus (mg)	942.03 ± 431.69	953.36 ± 435.35	891.41 ± 412.06	0.051
Potassium (mg)	1809.28 ± 1092.33	1832.24 ± 1111.25	1706.63 ± 999.25	0.118
Sodium (mg)	4504.28 ± 960.69	4480.03 ± 3611.3	4612.64 ± 5254.15	0.649
Magnesium (mg)	267.15 ± 147.96	270.85 ± 151.78	250.62 ± 128.51	0.063
Iron (mg)	20.67 ± 14.71	21.01 ± 15.57	19.17 ± 9.88	0.090
Zinc (mg)	9.79 ± 4.96	9.93 ± 5.09	9.18 ± 4.32	0.041
Selenium (µg)	50.21 ± 32.78	51.02 ± 33.68	46.6 ± 28.22	0.067
Copper (mg)	1.71 ± 1.17	1.74 ± 1.23	1.58 ± 0.85	0.068
Manganese (mg)	5.18 ± 12.46	5.29 ± 13.32	4.67 ± 7.51	0.499
Iodine (µg)	190.67 ± 739.12	165.74 ± 430.11	302.09 ± 1467.54	0.168
Vitamin A (µgRAE)	545.38 ± 602.61	557.82 ± 636.05	489.76 ± 418.59	0.125
Carotene (µg)	2095.42 ± 2740.11	2133.7 ± 2890.07	1924.36 ± 1928.59	0.299
Thiamine (mg)	0.79 ± 0.41	0.8 ± 0.42	0.74 ± 0.38	0.088
Riboflavin (mg)	0.92 ± 0.57	0.93 ± 0.57	0.86 ± 0.55	0.069
Niacin (mg)	14.8 ± 8.15	14.94 ± 8.11	14.14 ± 8.32	0.180
Vitamin C (mg)	80.66 ± 127.85	82.58 ± 139.54	72.09 ± 48.09	0.265
Vitamin E (mg)	23.06 ± 16.59	23.15 ± 15.34	22.68 ± 21.34	0.701
Folate (µg)	84.69 ± 113.54	85.25 ± 116.35	82.19 ± 100.2	0.714
Total 25(OH)D	22.29 ± 7.85	22.31 ± 7.77	22.18 ± 8.23	0.825

adjusted for all covariates, the medium intake of dietary live microbes ($OR = 0.64$, 95% CI : 0.44–0.93) and high intake of dietary live microbes ($OR = 0.41$, 95% CI : 0.22–0.77) groups were still correlated with a reduced risk of sarcopenia ($P = 0.007$).

Subgroup Analysis and Interaction Analysis

To evaluate whether the association between dietary live microbe intake and sarcopenia was modified by potential confounding factors, subgroup analyses were conducted for sex, age, diabetes, fatty liver disease, overweight, and obesity (Figure 3). The results revealed no statistically significant interactions between live dietary microbial groups and these factors ($P > 0.05$). However, a higher dietary intake of live microbes was found to be significantly associated with a reduced risk of sarcopenia in participants aged ≥ 65 years ($OR = 0.497$, 95% CI : 0.243–0.989); however, this association was not statistically significant in those aged < 65 years ($OR = 0.459$, 95% CI : 0.131–1.600).

Interestingly, a significant inverse association was found between a high dietary intake of live microbes and sarcopenia in the non-overweight subgroup ($OR = 0.382$, 95% CI : 0.195–0.751), while this association was not statistically significant in the overweight and obese subgroups ($OR = 0.524$, 95% CI : 0.164–1.677). Among participants without diabetes, compared to the group with a low intake of live dietary microbes, the OR for sarcopenia was 0.601 (95% CI : 0.398–0.909) for the medium intake group and 0.362 (95% CI : 0.184–0.714) for the high intake group. Among participants with diabetes, compared with the low-intake group, the OR was 0.783 (95% CI : 0.282–2.174) for the medium-intake group and 0.812 (95% CI : 0.126–5.216) for the high-intake group. Among individuals without fatty liver, high dietary microbe intake was significantly associated with a lower risk of sarcopenia ($OR = 0.429$, 95% CI : 0.221–0.836); however, this association was not statistically significant in those with fatty liver ($OR = 0.402$, 95% CI : 0.061–2.627).




Figure 2. Dietary sources of live microbes among the study participants. (A) Consumption prevalence of various fermented food categories. (B) Average daily intake among consumers.

Table 3. Logistic regression analysis of the association between sarcopenia and dietary live microbe group

Model	Low dietary live microbe group	Medium dietary live microbe group	High dietary live microbe group	P trend
Crude model	1 [Reference]	0.61 (0.45, 0.83)	0.46 (0.27, 0.76)	< 0.001
Model 1	1 [Reference]	0.58 (0.42, 0.79)	0.43 (0.25, 0.73)	< 0.001
Model 2	1 [Reference]	0.66 (0.46, 0.96)	0.41 (0.22, 0.75)	0.007
Model 3	1 [Reference]	0.66 (0.45, 0.95)	0.42 (0.23, 0.78)	0.008
Model 4	1 [Reference]	0.64 (0.44, 0.93)	0.41 (0.22, 0.77)	0.007

Note. Crude model: unadjusted model. Model 1: adjusted for gender, age. Model 2: adjusted for gender, age, education level and BMI. Model 3: adjusted for gender, age, education level, BMI, diabetes and fatty liver. Model 4: adjusted for gender, age, education level, BMI, diabetes, fatty liver, TC, protein, phosphorus, calcium and zinc intake.

DISCUSSION

Sarcopenia has a significant effect on the quality of life of older adults; however, few studies have investigated the relationship between sarcopenia and the intake of live microbes. In our study, the prevalence of sarcopenia was found to be 18.28%, which is higher than 16.70% reported in a previous study based on the West China Health and Aging Trend (WCHAT)^[20]. This also suggests the need for further research into sarcopenia in Shanghai, a representative of the eastern coastal cities of China.

Recent studies based on the NHANES database have focused on the role of the total intake of live dietary microbes in sarcopenia^[21-23]. The classification systems used in these studies were largely based on the consumption patterns of typical American diets and fermented foods such as yogurt,

cheese, and sauerkraut. In contrast, the Chinese diet features a broad range of fermented foods, including fermented tofu, pickled vegetables, and other region-specific products that differ in terms of live microbial composition and processing methods. Compared with cheese, which predominantly contains *Lactobacillus* and *Streptococcus* species^[24], fermented black soybeans (Dou chi) harbor a more diverse microbial community, including *Bacillus*, *Aspergillus*, and *Mucor*^[9]. These dietary differences may lead to distinct compositions of the gut microbiota and subsequent differences in health effects. Our findings confirm the association between a higher intake of live microbes and a reduced risk of sarcopenia in the Chinese population. Nevertheless, despite these differences in food sources, our study similarly found that a high dietary intake of live microbes was independently associated




Figure 3. Subgroup analyses for sarcopenia and live microbe intake in participants stratified by gender, age, diabetes, hypertension, fatty liver, and overweight/obesity.

with lower odds of sarcopenia ($OR = 0.41$, $P^J = 0.007$), even after controlling for a comprehensive range of demographic, nutritional, and health-related covariates. Prior studies based on the NHANES data have also reported a significant inverse association between live microbe intake and sarcopenia, with an OR of approximately 0.63 in the highest vs. lowest intake group^[25]. Dietary intake of live microbes has additionally been suggested to affect muscle health via the gut-muscle axis, based on existing research^[26]. Live microbes may improve muscle health largely by modulating gut function, as supported by relevant evidence^[27]. Specifically, live microbes can reshape the gut microbiota composition and enhance microbial diversity, primarily by increasing the abundance of beneficial commensal microbes, such as *Lactobacillus* and *Bifidobacterium* species^[28]. These microbes can outcompete pathogenic or pro-inflammatory bacteria, thereby restoring the microbial balance and promoting intestinal homeostasis. Healthy and diverse microbiota can promote the production of SCFAs, such as butyrate, acetate, and propionate, which can improve muscle function. SCFAs serve as energy sources for colonocytes and as signaling molecules that affect systemic metabolic and immune responses. Prior studies have shown that live microbe intake upregulates the anti-inflammatory cytokine IL-10 and downregulates the levels of ROS and the pro-inflammatory cytokine TNF- α ^[11]. These alterations minimize chronic inflammation-mediated muscle damage and decelerate muscle protein breakdown. Moreover, SCFAs may directly affect the skeletal muscle. In particular, butyric acid has been reported to activate several signaling pathways, such as AMP-activated protein kinase (AMPK) and PGC-1 α , both of which play critical roles in mitochondrial function, oxidative metabolism, and resistance to muscle atrophy^[29]. SCFAs may also affect the systemic availability of energy substrates and muscle-related anabolic signals, thereby contributing to muscle maintenance. In brief, ingestion of live microbes can affect muscle function through microbial colonization, thereby affecting the gut microbiota. Although a substantial body of research has focused on the association between probiotics and gut microbiota, the long-term use of probiotic supplements remains limited in the general population. Consuming foods that are naturally rich in live microbes not only enhances the diversity of the gut microbiota, but also provides substrates to support their growth. In fact, compared to probiotic supplements, the dietary

intake of live microbes through habitual foods may offer a more feasible approach for achieving sustained or even permanent colonization in the gut. As such, incorporating live microbe-rich foods into the diet may be a promising strategy for managing sarcopenia. However, the present study did not assess these metabolites and inflammatory markers, which precludes confirmation of the existence of these mechanisms.

Sarcopenia is controlled by many factors, including dietary intake of live microbes^[30]. In addition to exploring the relationship between sarcopenia and live microbes, the present study assessed the correlation between sarcopenia and other influencing factors, thereby strengthening our results. High consumption of live microbes was found to be significantly associated with a lower risk of sarcopenia in individuals aged > 65 years. Although this association was not statistically significant in those under 65 years of age, the lack of any significant overall interaction indicates that the protective effect may still be broadly relevant but more easily detectable in older populations^[31]. Furthermore, a significant decrease in the risk of sarcopenia was observed among non-overweight individuals, while this association was not statistically significant in the overweight and obese subgroups. This suggests that the metabolic burden associated with obesity may attenuate the beneficial effects of live microbes^[32]. Interestingly, a higher prevalence of fatty liver was observed in individuals without sarcopenia, indicating the presence of a complex interplay in which fatty liver may diminish the benefits of live microbes *via* metabolic dysregulation, inflammation, or alterations in the gut-liver axis^[33]. A stronger inverse association between live microbe intake and sarcopenia was found in individuals without diabetes than in those with diabetes, indicating that an individual's metabolic health status affects the efficacy of interventions based on live microbes^[34]. These findings highlight the potential interaction between metabolic health status and the efficacy of live microbe-based interventions in mitigating muscle loss. In individuals without diabetes, the intake of live microbes may have beneficial effects by reshaping the composition of the gut microbiota^[35]. However, these pathways may be compromised in patients with diabetes. These findings suggest that the protective effects of live microbes against sarcopenia vary depending on an individual's metabolic condition, highlighting the need for personalized strategies based on each individual's

metabolic profiles.

Our study has several noteworthy features. Firstly, we used a scientific sampling method and a validated assessment tool to improve the representativeness of the samples and the accuracy of the results. Further, this is the first study to investigate the association between the intake of live dietary microbes and sarcopenia in a Chinese population, and the second study to do so on a global scale. Therefore, it is important to study the relationship between live microbes, dietary habits, and sarcopenia development. However, this study has some limitations. First, due to the cross-sectional nature of this study, we could not establish a causal relationship between live microbial intake and the development of sarcopenia. Second, dietary data were collected using a 24-hour dietary recall method, which relies on memory and may therefore lead to recall bias or omissions, particularly among elderly participants. Third, our assessment of physical activity was categorized into only two levels, and did not distinguish between exercise types (e.g., resistance vs. aerobic exercise) or intensity, even though resistance exercise has been well-established as a core factor directly related to muscle mass and sarcopenia prevention. The diabetic subgroup in the present study had a relatively small sample size, which may have limited the statistical power. Currently, the classification of live dietary microbes is still in the estimation stage, and precise quantitative analyses of foods containing live microbes are lacking. In addition, the survey questionnaire did not collect information on the use of anti-inflammatory drugs, which may be a potential confounding factor. Therefore, future studies should strengthen the accuracy of the assessment of live microbes and their intake to improve the scientific validity and reproducibility of our results.

CONCLUSIONS

In conclusion, based on data from a Chinese population and a classification system for live microbial content in Chinese foods, the present study indicated that individuals with a higher consumption of live microbes exhibit a lower risk of developing sarcopenia, suggesting a potential protective role of a high intake of dietary microbes in maintaining skeletal muscle health. These findings emphasize the importance of live microbe-rich foods in health education and highlight the need for further studies on their underlying mechanisms.

Funding This research was supported by grants from the National Key R&D Program of China (Grant number: 2023YFF1104402) and the Chinese Nutrition Society (NO. CNS-BMSG2021-78), key projects in the three-year plan of the Shanghai Municipal Public Health system (2023–2025) (NO. GWVI-4), and key disciplines in the three-year Plan of Shanghai Municipal Public Health System (2023–2025) (NO. GWVI-11.1-42).

Competing Interests The authors declare that they have no conflicts of interest.

Ethics This study was approved by the Ethics Committee of the Shanghai Center for Disease Control and Prevention (KY-2024-16). Informed consent was obtained from all participants before enrollment, and all procedures were conducted in accordance with the Declaration of Helsinki.

Authors' Contributions X.S. contributed to the conceptualization, data curation, statistical analysis, original draft writing, and visualization. Z.W. was responsible for resources, writing, reviewing, and editing. J.W. participated in the conceptualization and methodology. W.L., Z.S., and L.S. were involved in data collection. C.S. and C.Y. conducted literature searches. G.S. supervised the study and contributed to the writing, review, and editing of the manuscript. J.Z. contributed to writing, reviewing, and editing the manuscript. X.X. was responsible for conceptualization, supervision, writing, review, and editing.

Acknowledgements The authors thank all participants for their contributions to this study.

Data Sharing Owing to participant privacy regulations, we were unable to provide public access to individual-level data in accordance with the ethical guidelines. Additionally, the written informed consent obtained from the study participants did not include provisions for public data sharing. Qualified researchers can request access to the aggregated dataset upon reasonable request by contacting the corresponding author. The supplementary materials will be available in www.besjournal.com.

Consent for publication Not applicable.

Clinical Trial Number Not applicable.

Received: February 13, 2026;

Accepted: August 11, 2026

REFERENCES

1. National Bureau of Statistics of China. China statistical yearbook 2024. China Statistics Press. 2024. (In Chinese)
2. Shanghai Municipal Bureau of Statistics. Shanghai statistical

- yearbook 2024. Shanghai Municipal Bureau of Statistics. 2024. (In Chinese)
3. de Sire A, Ferrillo M, Lippi L, et al. Sarcopenic dysphagia, malnutrition, and oral frailty in elderly: a comprehensive review. *Nutrients*, 2022; 14, 982.
 4. Li JJ, Zhang Z, Zhang JJ, et al. Joint associations of sarcopenia and social isolation with mortality: two prospective cohort studies across different cultural contexts. *Biomed Environ Sci*, 2026; 39, 3–14.
 5. Almohaisen N, Gittins M, Todd C, et al. Prevalence of undernutrition, frailty and sarcopenia in community-dwelling people aged 50 years and above: systematic review and meta-analysis. *Nutrients*, 2022; 14, 1537.
 6. Hu YS, Peng WJ, Ren RJ, et al. Sarcopenia and mild cognitive impairment among elderly adults: the first longitudinal evidence from CHARLS. *J Cachexia Sarcopenia Muscle*, 2022; 13, 2944–52.
 7. Papadopoulou SK, Papadimitriou K, Voulgaridou G, et al. Exercise and nutrition impact on osteoporosis and sarcopenia: the incidence of osteosarcopenia: a narrative review. *Nutrients*, 2021; 13, 4499.
 8. Locantore P, Del Gatto V, Gelli S, et al. The interplay between immune system and microbiota in osteoporosis. *Mediators Inflamm*, 2020; 2020, 3686749.
 9. Zhang T, Cheng JK, Hu YM. Gut microbiota as a promising therapeutic target for age-related sarcopenia. *Ageing Res Rev*, 2022; 81, 101739.
 10. Zmora N, Suez J, Elinav E. You are what you eat: diet, health and the gut microbiota. *Nat Rev Gastroenterol Hepatol*, 2019; 16, 35–56.
 11. Chen LH, Chang SS, Chang HY, et al. Probiotic supplementation attenuates age-related sarcopenia via the gut-muscle axis in SAMP8 mice. *J Cachexia Sarcopenia Muscle*, 2022; 13, 515–31.
 12. Ticinesi A, Lauretani F, Milani C, et al. Aging gut microbiota at the cross-road between nutrition, physical frailty, and sarcopenia: is there a gut-muscle axis?. *Nutrients*, 2017; 9, 1303.
 13. Leeuwendaal NK, Stanton C, O'Toole PW, et al. Fermented foods, health and the gut microbiome. *Nutrients*, 2022; 14, 1527.
 14. Marco ML, Hutkins R, Hill C, et al. A classification system for defining and estimating dietary intake of live microbes in US adults and children. *J Nutr*, 2022; 152, 1729–36.
 15. Ding L, Duan JN, Hou JJ, et al. Association between dietary live microbe intake and chronic diarrhea and fecal incontinence: a cross-sectional NHANES 2005–2010 study. *J Am Nutr Assoc*, 2025; 44, 342–52.
 16. Yuan F. Association of dietary live microbe intake with prevalence of osteoporosis in US postmenopausal women: a cross-sectional study. *Arch Osteoporos*, 2024; 19, 69.
 17. Rosas-Carrasco O, Ruiz-Valenzuela RE, López-Teros MT. Phase angle cut-off points and their association with sarcopenia and frailty in adults of 50–64 years old and older adults in Mexico City. *Front Med*, 2021; 8, 617126.
 18. Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*, 2018; 48, 16–31.
 19. Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*, 2020; 54, 1451–62.
 20. Liu X, Hao Q, Yue J, et al. Sarcopenia, obesity and sarcopenia obesity in comparison: prevalence, metabolic profile, and key differences: results from WCHAT study. *J Nutr Health Aging*, 2020; 24, 429–37.
 21. Fielding RA, Reeves AR, Jasuja R, et al. Muscle strength is increased in mice that are colonized with microbiota from high-functioning older adults. *Exp Gerontol*, 2019; 127, 110722.
 22. Lee K, Kim J, Park SD, et al. *Lactobacillus plantarum* HY7715 Ameliorates sarcopenia by improving skeletal muscle mass and function in aged Balb/c mice. *Int J Mol Sci*, 2021; 22, 10023.
 23. Lee MC, Tu YT, Lee CC, et al. *Lactobacillus plantarum* TWK10 improves muscle mass and functional performance in frail older adults: a randomized, double-blind clinical trial. *Microorganisms*, 2021; 9, 1466.
 24. Burns P, Patrignani F, Serrazanetti D, et al. Probiotic Crescenza cheese containing *Lactobacillus casei* and *Lactobacillus acidophilus* manufactured with high-pressure homogenized milk. *J Dairy Sci*, 2008; 91, 500–12.
 25. Yan KM, Ma XY, Li C, et al. Higher dietary live microbe intake is associated with a Lower risk of sarcopenia. *Clin Nutr*, 2024; 43, 1675–82.
 26. Milani C, Duranti S, Napoli S, et al. Colonization of the human gut by bovine bacteria present in Parmesan cheese. *Nat Commun*, 2019; 10, 1286.
 27. Liu CR, Wong PY, Barua N, et al. From clinical to benchside: *Lactocaseibacillus* and *Faecalibacterium* are positively associated with muscle health and alleviate age-related muscle disorder. *Ageing Cell*, 2025; 24, e14485.
 28. Rashidah NH, Lim SM, Neoh CF, et al. Differential gut microbiota and intestinal permeability between frail and healthy older adults: a systematic review. *Ageing Res Rev*, 2022; 82, 101744.
 29. Jung TW, Park HS, Choi GH, et al. β -aminoisobutyric acid attenuates LPS-induced inflammation and insulin resistance in adipocytes through AMPK-mediated pathway. *J Biomed Sci*, 2018; 25, 27.
 30. Chen LK, Woo J, Assantachai P, et al. Asian working group for sarcopenia: 2019 consensus update on sarcopenia diagnosis and treatment. *J Am Med Dir Assoc*, 2020; 21, 300–7. e2.
 31. Papadopoulou SK. Sarcopenia: a contemporary health problem among older adult populations. *Nutrients*, 2020; 12, 1293.
 32. Merchant RA, Seetharaman S, Au L, et al. Relationship of fat mass index and fat free mass index with body mass index and association with function, cognition and sarcopenia in pre-frail older adults. *Front Endocrinol*, 2021; 12, 765415.
 33. Polyzos SA, Vachliotis ID, Mantzoros CS. Sarcopenia, sarcopenic obesity and nonalcoholic fatty liver disease. *Metabolism*, 2023; 147, 155676.
 34. Sambashivaiah S, Harridge SDR, Sharma N, et al. Asian Indians with prediabetes have similar skeletal muscle mass and function to those with type 2 diabetes. *Front Nutr*, 2019; 6, 179.
 35. Zoncu R, Efeyan A, Sabatini DM. mTOR: from growth signal integration to cancer, diabetes and ageing. *Nat Rev Mol Cell Biol*, 2011; 12, 21–35.