

Review



Emerging Technologies for Viability Assessment of Foodborne Pathogens: From Molecular Dyes to Artificial Intelligence

Zeyu Wang^{1,3,&}, Anum Farid^{2,3,&}, Zhiwei Chen^{2,3}, Jiayi Qi^{1,3}, Xinxin Shen^{4,#}, and Hong Liu^{1,3,#}

Abstract: To critically examine emerging technologies for assessing foodborne-pathogen viability and to evaluate their relevance for improving food safety monitoring and public-health risk assessment. This review examines the literature on viability-detection using biological indicators such as membrane integrity, infectivity, and virulence. It addresses technologies such as viability dye-based PCR, RNA and CRISPR detection methods, biosensors, bacteriophage assays, Raman spectroscopy, and AI-supported analysis. Compared to conventional culture-based and DNA-only molecular methods, viability-focused techniques support more biologically meaningful detection, particularly for viable but non-culturable (VBNC) cells that may retain pathogenic potential. However, assay performance varies depending on food-matrix complexity, the presence of inhibitors, sensitivity limitations, and the degree of validation. Integrated systems combining multi-modal sensing, microfluidics, and machine learning have shown increasing promise for rapid on-site application, although standardization and regulatory validation remain limited. The emerging viability-focused technologies substantially refine microbial-risk assessment by accurately capturing the infectious potential of foodborne pathogens, including traditionally undetected VBNC populations. Nevertheless, integrating these technologies into routine surveillance remains constrained by matrix interference and the absence of standardized protocols. Future efforts must prioritize developing matrix-resilient methodologies and establishing regulatory validation frameworks to facilitate their practical implementation in food safety

management.

Key words: Foodborne microbiology; Viability; VBNC; Food safety; Emerging technologies; Artificial intelligence

INTRODUCTION

Foodborne pathogens continue to pose a significant threat to global food safety and public health, as the consumption of contaminated food can lead to a diverse array of diseases and impose considerable health and socioeconomic burdens^[1]. This ongoing threat necessitates robust, continuous monitoring to remove contaminated products from the supply chain. However, conventional food safety surveillance faces two fundamental and interconnected challenges: the lack of discrimination between viable pathogens and non-viable cells^[2], and the paucity of practical, multiplexed systems capable of detecting the diverse range of pathogens that threaten the food supply without prior knowledge of the contaminant^[3,4]. Culture-based techniques, the historical gold standard, are notoriously slow, requiring lengthy enrichment periods that are incompatible with the rapid distribution of perishable goods like milk^[5]. More critically, they fail to detect bacteria in the viable but non-culturable (VBNC) state, a dormant condition induced by food processing stresses, in which the cells remain metabolically active and potentially pathogenic but resist cultivation^[6].

While molecular techniques like PCR have revolutionized detection speed, they have introduced a different problem: the inability to differentiate between DNA from live cells and

doi: [10.3967/bes2026.109](https://doi.org/10.3967/bes2026.109)

1. School of Life Sciences and Medicine, Shandong University of Technology, Zibo 255049, Shandong, China; 2. School of Agriculture Engineering and Food Science, Shandong University of Technology, Zibo 255049, Shandong, China; 3. Shandong Provincial Innovation Centre for Dairy Technology, Zibo 255049, Shandong, China; 4. National Key Laboratory of Intelligent Tracking and Forecasting for Infectious Diseases, NHC Key Laboratory of Medical Virology and Viral Diseases, National Institute for Viral Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing, 102206, China

residual genetic material from dead cells, leading to false-positive signals that can trigger economically devastating recalls in the absence of a genuine public health risk. The distinction between these food-born pathogen states—culturable, VBNC, and dead—is therefore paramount for accurate risk assessment^[7]. This technological gap has promoted the development of a new generation of detection platforms specifically designed for viability assessment, which can be broadly classified by their underlying mechanistic principles^[8]. One prominent approach focuses on nucleic-acid differentiation, employing agents like propidium monoazide (PMA) in conjunction with quantitative PCR (qPCR) or digital droplet PCR (ddPCR) to selectively intercalate with DNA from membrane-compromised dead cells, thereby inhibiting its amplification and enabling the specific quantification of viable pathogens.

Infectivity-based detection provides a fundamentally different strategy; this approach leverages the natural specificity of bacteriophages—viruses that infect and lyse only metabolically active bacterial cells—to provide a direct functional readout of viability^[9]. Beyond approaches based on nucleic acids and infectivity, another innovative pathway involves metabolic and volatile profiling. Platforms such as machine learning (ML)-enabled paper chromogenic arrays (PCAs) detect unique volatile organic compounds emitted specifically by viable pathogens, offering a non-destructive, recognition-element-free route to multiplex detection that bypasses the need for predefined molecular targets^[10].

The convergence of these biological assays with enabling technologies is defining the future of food-safety diagnostics. Microfluidics facilitates the miniaturization and automation of complex assays, integrating processes like PCR, ELISA, and isothermal amplification into portable "lab-on-a-chip" devices for rapid, on-site analysis^[11]. Simultaneously, artificial intelligence (AI) and deep learning are revolutionizing data interpretation, from analyzing colorimetric patterns to automating the microscopic identification of pathogens, thereby enhancing speed, objectivity, and accuracy^[12].

This review presents a comprehensive, critical analysis of the emerging technologies for assessing foodborne pathogen viability. We examine their underlying principles, systematically categorize them based on methodology, and assess their respective advantages and limitations within complex food matrices. The transformative potential of integrating biosensors with microfluidics and AI is investigated.

Future directions, focusing on the development of intelligent, multiplexed systems designed to provide rapid, precise, and actionable data essential for the proactive protection of the global food supply, are discussed^[13].

LITERATURE SEARCH STRATEGY AND SELECTION CRITERIA

This review employed a structured literature search method, focusing on recent advances in foodborne-pathogen viability-assessment technologies. Relevant articles were retrieved from Scopus, Web of Science, PubMed, ScienceDirect, Wanfang Data, and the China National Knowledge Infrastructure (CNKI). The literature search covered studies published from 2010 to 2026, with particular emphasis on publications from 2020 onward. Search terms included foodborne microbiology, bacterial viability, VBNC, live/dead discrimination, PMA/EMA (referring propidium monoazide and ethidium monoazide), viability PCR, rapid microbial detection, biosensors, CRISPR-Cas, microfluidics, emerging technologies, and AI. Additional studies were identified by manually screening the references of the included articles. Studies were included if they focused on viability assessment, live/dead discrimination, VBNC-state detection, or the rapid monitoring of foodborne pathogens, and reported methods relevant to food safety or food microbiology. Studies were excluded if they were unrelated to foodborne pathogens, lacked sufficient methodological details, were conference abstracts, editorials, or non-peer-reviewed documents, or provided duplicated information with limited relevance to recent technological advances. A critical analysis of selected studies was conducted to evaluate recent advances in emerging detection technologies and AI-assisted methodologies for assessing foodborne microbe viability.

KEY BACTERIAL VIABILITY STATES: A CRITICAL DISTINCTION FOR DETECTION

The accurate detection of pathogens in samples of foods such as milk, dairy products, raw meat, poultry, seafood, fresh produce, and ready-to-eat foods is complicated by the fact that not all viable cells are capable of proliferating on standard culture media. This arises from the fact that the distinct types of physiological viability—reproductive viability, the ability to form a colony, and metabolic viability, the presence of metabolic activity—do not

always coincide^[14]. This distinction is critical, as conventional culture methods exclusively detect reproductively viable cells, thereby missing sublethally injured and VBNC cells that remain metabolically active and pose a potential health risk^[15]. The primary challenge, therefore, lies in differentiating these states. The VBNC state, a survival strategy induced by stressors such as refrigeration or processing, is exhibited by cells that are metabolically active but non-culturable^[16]. Dead cells, in contrast, have lost both culturability and metabolic activity, and cell death is typically indicated by a compromised membrane. This fundamental difference necessitates the use of molecular viability-testing methods that can discriminate between living cells (culturable or not) and dead cells; such methods, including PMA-qPCR, facilitate more accurate safety assessment than culture-based counts^[17]. A side-by-side comparison of these key viability states, including their defining characteristics and biomarker profiles, is presented in Table 1. The relationships among these viability states are visually summarized in Figure 1.

THE VBNC STATE: A CRITICAL JUSTIFICATION FOR ADVANCED VIABILITY DETECTION

The limitations inherent in traditional culture-based methodologies are most prominently

highlighted by the VBNC state. This dormant survival strategy, triggered by food-related stresses such as refrigeration, disinfectants, or nutrient deprivation, causes bacterial pathogens like *Vibrio vulnificus*, *Escherichia coli* O157: H7, and *Listeria monocytogenes* ceasing to proliferate on culture media^[18]; however, they remain metabolically active, with the potential to resuscitate into culturable, infectious cells^[19]. This potential for resuscitation poses a significant challenge to food safety, as it creates a reservoir of "hidden" pathogens that can elude detection during routine monitoring yet cause illness upon consumption. The mechanisms underlying resuscitation are varied and involve environmental cues such as increases in temperature and nutrient availability, the action of chemical agents like surfactants, and notably, passage through a host's gastrointestinal tract, which can trigger the revival of pathogens such as *Campylobacter jejuni* and *Vibrio cholerae*^[20]. This capacity to regain cultural viability and virulence upon entering a favorable niche underscores the fact that non-culturability is not synonymous with non-viability.

The physiological characteristics of the VBNC state have important implications for the design of advanced detection technologies. Given that these cells cannot be cultured, it is imperative that methods do not rely on growth as an indicator of

Table 1. Key bacterial viability states relevant to foodborne pathogen detection

Viability state	Key features	Inducing factors	Detection challenges	Representative detection methods
Reproductively viable cells	Actively proliferating; colony-forming	- Optimal nutrients - Favourable growth conditions	Easily detected, but overestimates dominance when mixed with other states	- Culture-based methods - qPCR (DNA-based)
Intact cells	Preserved membrane integrity but little/no metabolism	- Dormancy - Reversible non-viability	- Not culturable - May persist in biofilms or milk without growth	- Microscopy, viability - Stains (SYTO 9/PI) - FTIR/Raman
Vital cells	Metabolically active but sometimes slow-growing; extended lag phase	Mild stress (nutrient shift, osmotic stress, sublethal treatment)	- Missed by culture - May contribute to spoilage despite low culturability	- Flow cytometry - Metabolic stains - Molecular assays
Dead cells	Loss of membrane integrity and metabolism	- Heat - Disinfectants - Strong chemical stress	DNA from dead cells may give false positives in PCR, complicating viability assessments	- Propidium iodide staining - PMA/EMA-qPCR
VBNC cells	Metabolically active but non-dividing; possible resuscitation	- Pasteurisation - Nutrient limitation - Oxidative/osmotic stress	Undetected by culture; may resuscitate underestimation of risk	- PMA-qPCR, RT-qPCR - Flow cytometry - Predictive AI models
Dormant cells	Metabolism shutdown; long-term persistence	- Starvation - Osmotic stress - Nutrient deprivation	Invisible to culture and many assays, long-term survival in milk/biofilms	- Advanced microscopy - Resuscitation assays - Omics-based tools

viability. Instead, owing to

The metabolic activity of VBNC cells, assays should focus on detecting fundamental indicators of life, such as membrane integrity, the presence of ATP, RNA transcription, and the activity of key enzymes^[21]. The identification of conserved biological mechanisms, such as resuscitation-promoting factor (Rpf) proteins in species like *Micrococcus luteus* and *Mycobacterium* spp., highlights the fact that resuscitation is an active,

regulated process^[22]. Consequently, a comprehensive viability assessment must distinguish between dead cells, dormant-but-resuscitable cells, and actively metabolizing cells. This complex diagnostic requirement is the primary impetus for the development of the sophisticated platforms discussed here; such platforms have been developed to address the critical detection gap left by traditional methods and to mitigate the public health risks posed by the presence of resuscitating VBNC




Figure 1. Key bacterial viability states relevant to foodborne microbiology. Illustration of major bacterial viability states, ranging from reproductively viable, intact, and vital cells to dormant, VBNC, and dead cells. The diagram provides a framework for understanding detection targets in food safety applications. Abbreviation: VBNC, viable but non-culturable.

pathogens in food^[23].

ADVANCED TECHNOLOGICAL PLATFORMS FOR FOODBORNE PATHOGEN VIABILITY DETECTION

The limitations of traditional culture methods have catalyzed the development of sophisticated technological platforms for viability assessment. These approaches move beyond mere detection to provide functional insights into pathogen viability through various mechanistic pathways. This section critically examines these advanced platforms, categorized by their fundamental operating principles and technological maturity. Their key limitations and advantages are summarized in Table 2.

Nucleic Acid-Based Molecular Platforms

Viability Dye-Coupled PCR (vPCR) Viability dye-

coupled PCR overcomes a key limitation of conventional PCR by reducing signals from non-viable cells. In this approach, amplification from dead cells is suppressed by using membrane-impermeant dyes such as PMA and PMAxx to covalently bind DNA in membrane-compromised cells after photoactivation^[24]. PMA/PMAxx-based vPCR has been combined with isothermal amplification methods, including cross-primer amplification (CPA) and real-time fluorescent loop-mediated isothermal amplification (LAMP), for selective viable-pathogen detection by suppressing amplification from dead-cell DNA. PMA-CPA has enabled the rapid detection of *Salmonella enterica*, including VBNC cells, in complex food systems, with detection limits reaching 101 CFU/mL within 2 h. However, vPCR remains limited by its reliance on membrane integrity as a surrogate marker of viability. Dead or irreversibly inactivated cells with

Table 2. Established technological platforms for viable foodborne pathogen detection

Platform category	Specific technology	Target viability marker	Key advantages	Current limitations & challenges
Nucleic acid-based platforms	Viability Dye-PCR (vPCR; e.g., PMA-qPCR)	Membrane integrity	- Quantitative - High throughput - Compatible with standard lab workflows	- Cannot detect VBNC cells with intact membranes - Matrix interference affects dye penetration - Requires extensive optimization
	CRISPR-Cas detection (e.g., Cas13a)	Metabolic activity (via labile mRNA)	- High specificity & programmability - Excellent sensitivity with signal amplification - Potential for portable, rapid formats	- Primarily proof-of-concept for complex foods - Risk of false positives from transient RNA - Requires optimization for food matrices
Biosensor platforms	Electrochemical biosensors	Metabolic activity (e.g., respiration, redox species)	- Real-time, label-free monitoring - High portability for on-site use - Potential for continuous monitoring	- Can lack pathogen specificity without recognition elements - Sensor fouling in complex matrices - Signal stability over time
	Aptamer-based biosensors (Aptasensors)	Specific molecular targets (e.g., surface proteins, enzymes)	- High affinity and specificity - Superior thermal stability vs. antibodies - Tunable and reproducible production	- Susceptible to nuclease degradation - Limited target range for some pathogens - Complex, costly development process
	ATP bioluminescence Biosensors	Metabolic activity (via intracellular ATP)	- Extreme rapidity (results in minutes) - Simple operation and highly portable - Excellent for hygiene monitoring	- Not pathogen-specific (detects all ATP) - Poor sensitivity for VBNC cells with low metabolic activity - High background in some food types
Biological recognition platforms	Bacteriophage-based assays	Infectious Potential / Metabolic Activity	- High biological relevance for viability - Can detect resuscitating VBNC cells - Exceptional specificity for target bacteria	- Narrow host range of phages requires extensive libraries - Can be multi-step and time-consuming - Lower throughput than molecular methods
Label-free analytical platforms	Raman spectroscopy	Biochemical Composition (Molecular Fingerprint)	- Label-free, non-destructive analysis - Single-cell resolution - Can differentiate live, dead, and VBNC states	- Costly instrumentation - Inherently weak signal requires enhancement - Requires sophisticated data analysis (e.g., AI)
	Advanced microscopy with AI (e.g., AFM)	Cell Morphology & Membrane Integrity	- Nanoscale resolution of structural changes - Automated, quantitative classification - Provides definitive morphological data	- Very low throughput - Expensive instrumentation and expertise required - Complex sample preparation

intact membranes may be misclassified as viable, resulting in overestimation of viable cell counts^[25,26]. In complex food matrices, incomplete dye penetration and matrix interference can further increase false-positive risks^[27]. Assay performance depends strongly on optimized dye concentration, incubation, and photoactivation conditions, which may limit routine application.

CRISPR-Cas Enhanced Viability Detection The CRISPR-Cas system has become an important nucleic acid-based platform for foodborne pathogen detection owing to its high sequence specificity, programmability, and compatibility with isothermal amplification^[28,29]. In Cas12- and Cas13-based detection systems, crRNA-guided Cas effectors recognize target nucleic acids and activate collateral cleavage of ssDNA or ssRNA reporter molecules, thereby generating amplified fluorescence signals^[30]. For viability-associated detection, CRISPR-Cas assays often use mRNA as the target because bacterial mRNA generally has a shorter half-life than DNA and can provide a signal related to recent transcriptional activity^[31]. Although this strategy can reduce interference from residual DNA in dead cells, mRNA detection does not directly confirm true cellular viability or infectious potential, as residual RNA may persist transiently after cell death and cause false-positive results^[32,33]. RNA stability, reverse-transcription efficiency, amplification bias, and inhibitors in food matrices may also affect assay accuracy^[34]. Therefore, mRNA-targeting CRISPR assays should be regarded as indicators of recent biological activity rather than as definitive measures of active viability, and further validation with other physiological indicators is required for practical applications^[35]. Current challenges include optimizing the system to prevent false-positive signals from transient RNA molecules and ensuring robust performance in complex food matrices that may contain amplification inhibitors in complex food matrices. Many food-related CRISPR assays remain at the early validation stage, although some platforms have already shown promising performance in real-sample testing.

Biosensor Platforms Electrochemical Biosensors

Electrochemical biosensing enables viability-associated pathogen detection by monitoring metabolism-related electrochemical signals, such as respiration activity and redox cycling^[36,37]. Modern configurations achieve specificity by employing biological recognition interfaces (antibodies, aptamers, or phage receptors) that isolate target organisms, thereby allowing metabolic signals to be

correlated with specific pathogen identities^[38,39]. The sensitivity of these platforms is often boosted by nanomaterials like gold nanoparticles and carbon nanotubes, which improve conductivity and increase bioreceptor-loading capacity^[40]. Recent innovations have focused on multiplexed arrays for simultaneous detection and microfluidic integration to automate sample processing. Despite these technical strides, challenges persist regarding matrix interference, electrode biofouling, and the need for robust calibration across different food products^[41,42]. Ultimately, the capacity for continuous data collection positions electrochemical sensors as a critical tool for real-time safety management in food processing.

Aptamer-Based Biosensors

Aptamer-based biosensors, or aptasensors, represent a notable biological recognition platform that employs synthetic single-stranded DNA or RNA ligands selected via Systematic Evolution of Ligands by EXponential enrichment (SELEX)^[43,44]. These recognition elements offer distinct advantages over traditional antibodies, including enhanced thermal stability, ease of synthesis and modification, and reproducible production. In contemporary aptasensor configurations, sophisticated designs are utilized to augment their detection capability. Sandwich-type aptasensors coordinate with metal ions to improve sensitivity toward metabolic indicators such as cyclic AMP (cAMP), facilitating direct assessment of physiological activity^[45]. Structure-switching aptamers can convert target-binding events into measurable electrochemical or optical signals, enabling the rapid and label-free detection of foodborne pathogens^[46]. However, the practical application of aptasensors remains challenged by aptamer degradation by nucleases, nonspecific adsorption, complex food-matrix interference, and the limited availability of high-affinity aptamers for some pathogens. Ongoing advances in aptamer engineering, modified nucleotides, nanomaterial-assisted signal amplification, and stabilization strategies are expected to improve their robustness for future food safety monitoring^[47,48].

ATP-Based Biosensors

ATP-based biosensors offer a rapid means of assessing microbial viability by quantifying intracellular adenosine triphosphate (ATP), the universal energy currency of living cells^[49]. These platforms typically utilize luciferase-luciferin reactions that produce bioluminescence proportional to the ATP concentration, facilitating detection within minutes rather than hours or days.

Recent technological advancements have improved detection sensitivity to ca. 10^3 – 10^4 CFU/mL, rendering these biosensors valuable for hygiene monitoring and rapid screening applications^[50,51]. The incorporation of enhanced sample concentration methods and more stable enzyme formulations has broadened their applicability to various food matrices^[52]. The primary limitation of ATP-based detection is its lack of pathogen specificity, as it detects ATP from all biological sources, including food cells and background microflora. Although this constrains its utility for detecting specific pathogens in complex food samples, it remains invaluable for general hygiene assessment and process control^[53].

Biological Recognition Platforms

Bacteriophage-Based Assays Bacteriophage-based assays provide a biologically relevant strategy for bacterial viability assessment by exploiting the host specificity of phages and their requirement for physiologically active bacterial hosts^[54,55]. Unlike conventional plaque assays that rely on visible plaque formation, recent phage-based platforms often integrate phage amplification or phage-mediated bacterial lysis with qPCR, enzyme-linked immunosorbent assay (ELISA), luminescence, or fluorescence readouts, thereby reducing detection time while maintaining high specificity^[55–57]. Because productive phage infection generally requires intact host structures and sufficient metabolic activity, these methods preferentially report viable or infection-competent bacterial cells rather than dead cells. This functional recognition is particularly valuable for detecting metabolically active pathogens and may also support the identification of resuscitating VBNC cells under appropriate conditions^[58]. Nevertheless, the practical application of phage-based assays remains limited by the narrow host range of individual phages, strain-level variability in phage susceptibility, and the need to establish well-characterized phage libraries^[59].

Despite their high specificity for viable bacterial hosts, phage-based detection systems remain strongly dependent on host physiological activity and environmental conditions, which may significantly affect their analytical performance in complex food systems. Successful phage infection and subsequent signal amplification require metabolically active host cells capable of supporting phage adsorption, replication, and propagation^[60]. Consequently, environmental stresses commonly encountered during food processing and storage, including low

temperature, high salinity, acidic conditions, oxidative stress, and preservative exposure, may suppress bacterial metabolic activity and reduce phage-amplification efficiency even when viable cells remain present. These limitations may be particularly relevant in refrigerated and cold-chain food environments, where bacterial cells frequently exhibit reduced metabolic rates or transition into stressed physiological states, including sublethal injury and VBNC conditions. Complex food matrices may impair phage-based detection by limiting phage diffusion, promoting surface adsorption, or introducing inhibitory components that reduce phage access to target bacteria. Strain-level variability and phage-host specificity may further affect assay sensitivity and reproducibility. Therefore, although phage-based platforms show strong potential for selective viable pathogen detection, their robustness requires validation under realistic food processing and storage conditions.

Label-Free Analytical Platforms

Raman Spectroscopy-Based Platforms Raman spectroscopy provides a label-free approach for viability assessment by capturing the molecular fingerprints of individual bacterial cells^[61]. Changes in nucleic acids, proteins, lipids, and other biochemical components can be used to distinguish between live, stressed, and dead cells, especially when combined with microfluidic platforms and machine learning-based spectral analysis^[62,63]. Unlike conventional viability assays, Raman spectroscopy offers single-cell resolution, and it provides richer physiological information. However, its practical application is still limited by weak Raman signals, relatively complex instrumentation, and high equipment costs^[64].

Advanced Microscopy with AI Integration The integration of AI with microscopy and spectroscopic platforms has improved the quantitative interpretation of bacterial viability-related images and signals^[65]. High-resolution analytical modalities, including Atomic Force Microscopy (AFM), Cryo-Electron Microscopy (Cryo-EM), fluorescence imaging, and Raman spectroscopy, provide detailed structural and biochemical information associated with cellular viability states. These technologies enable the evaluation of viability-associated characteristics such as membrane integrity, intracellular ultrastructural organization, biochemical composition, and metabolic activity that are not readily distinguishable using conventional optical microscopy alone^[66,67]. AI-based image and spectral

analysis has increasingly been used to classify viable and non-viable microbes, with early applications mainly based on CNN-derived microscopy features^[68,69]. More recent studies have additionally employed computational approaches such as support vector machines (SVMs), random forest classifiers, partial least squares discriminant analysis (PLS-DA), linear discriminant analysis (LDA), and hybrid spectral-feature analysis methods to improve the discrimination of subtle phenotypic and biochemical variations observed in Raman spectroscopy and fluorescence-based datasets^[64,70,71]. These approaches improve the recognition of viability-associated phenotypes, particularly when microbial signals are masked by complex food matrix backgrounds. Despite these advances, AI-assisted viability detection remains limited by dataset size, model generalizability, and external validation. Many models are trained on small and highly controlled datasets, increasing the risk of overfitting and reducing robustness across different food matrices, processing conditions, imaging systems, and pathogen strains. Variations in sample preparation, preprocessing procedures, signal quality, and matrix interference may further affect reproducibility. Future studies should prioritize diverse strain- and matrix-level datasets, standardized validation workflows, transparent model reporting, and robustness testing under realistic food processing and storage conditions. Although still constrained by instrumentation cost, computational demand, and limited field validation, these approaches offer considerable potential for high-resolution viability assessment and automated microbial phenotype interpretation^[72]. **Figures 2 and 3** compare the established and emerging viability detection technologies.

EMERGING TECHNOLOGIES FOR VIABLE-PATHOGEN DETECTION

Although the established detection platforms are reliable for assessing microbial viability, emerging technologies offer new strategies to overcome the key limitations of the conventional methods, including limited physiological resolution, prolonged analysis time, and poor adaptability to complex food matrices^[73]. These platforms integrate advanced-sensing modalities, engineered biomaterials, microfluidic systems, artificial intelligence-assisted analytics, and next-generation nucleic acid technologies to improve the sensitivity, specificity, scalability, and practical applicability of food-safety

monitoring. To facilitate comparison among these emerging approaches, the technological-maturity classifications summarized in **Table 3** are organized according to adapted Technology Readiness Level (TRL)^[74,75] principles and their corresponding validation status. In parallel, **Table 4** compares the representative emerging platforms in terms of detection limit, assay time, multiplexing capability, and estimated cost per test, providing a practical framework for evaluating their applicability in foodborne pathogen viability assessment.

Machine Learning-Enabled Metabolomic Sensing

Paper-based Chromogenic Array with Neural Networks (PCA-NN) The integration of paper-based chromogenic arrays with ML signifies a transformative advancement in non-destructive viability-associated monitoring. In this technology, an array of 23 chromogenic dyes, which selectively react with volatile organic compounds (VOCs) emitted by metabolically active pathogens, is applied to paper substrates. The primary innovation does not reside in the individual dye-pathogen associations, but rather in the utilization of a multi-layer neural network to interpret the comprehensive colorimetric fingerprint produced by the array^[75]. The PCA-NN system has exhibited exceptional capabilities in practical applications, achieving 91%–95% accuracy in the strain-specific identification of pathogens such as *E. coli* O157: H7 and *L. monocytogenes* on fresh-cut romaine lettuce^[76]. The technology functions without the need for enrichment, culturing, or sample preparation, detecting populations as low as 10³ CFU/mL within 2 h of exposure^[76,77]. The neural network architecture, typically comprising five hidden layers with 256 neurons each, processes the digitized RGB values from the array patterns and correlates them with specific pathogens, population levels, and even growth stages. Current research endeavors aim to broaden this technology's applicability across various food commodities and pathogen combinations. With an estimated production cost below US\$0.10 per unit^[77], PCA-NN systems exhibit significant potential for smart packaging applications. Nonetheless, challenges persist in managing background VOC interference from food matrices and establishing comprehensive training datasets for diverse food-pathogen combinations^[77].

Advanced Functional Biosensing Platforms

Nanozyme-Enhanced Microfluidic Biosensors The

integration of nanotechnology with microfluidics has enabled miniaturized and portable biosensing platforms with improved sensitivity, reduced sample consumption, and the potential for on-site food-safety monitoring. Nanozymes—nanomaterials with enzyme-like catalytic properties—provide robust signal amplification and achieve greater operational stability than natural enzymes. As these platforms are engineered to detect the metabolic products and redox-active compounds secreted exclusively by viable cells, they extend beyond simple pathogen detection, directly linking the cell's catalytic signals

to their microbial viability^[78]. In these systems, nanozymes are commonly coupled with immunorecognition, aptamer recognition, or magnetic enrichment to capture target bacteria, while their peroxidase-like or oxidase-like catalytic activity generates colorimetric, electrochemical, or chemiluminescent signals^[79]. Recent studies using *Salmonella* have demonstrated the analytical potential of this strategy. For example, a finger-actuated microfluidic colorimetric biosensor using Au@PtPd nanocatalysts detected *Salmonella* at concentrations as low as 45 CFU/mL within 25




Figure 2. Established and nucleic acid-based methods for viable pathogen detection. Overview of current and advanced methodologies for viability assessment of foodborne pathogens, including Viability Dye-PCR, CRISPR-Cas platform, biosensors, immunological assays, phage-based assays, and Raman spectroscopy integrated with microfluidics.




Figure 3. Emerging and advanced methods for viable pathogen detection. Conceptual framework illustrating how novel detection tools can be embedded into food safety management systems to enable proactive, multiplexed, and real-time monitoring of microbial risks across the dairy and food supply chain.

Table 3. Emerging and advanced methods for viable pathogen detection

Technology category	Specific platform	Key innovation	Technology readiness classification and validation Status	Major challenges
Metabolomic sensing	Paper chromogenic array (PCA-NN)	Non-destructive VOC profiling with neural network decoding	Laboratory prototype validation (TRL 4–5); limited real food matrix validation	• Background VOC interference • Ttraining data requirements
	Nanzyme-enhanced microfluidics	Nanomaterial catalysis with automated fluid handling	Laboratory prototype validation (TRL 4–5); laboratory-scale validation	• Channel fouling • Nanzyme stability
Functional biosensing	Split-protein systems	Direct virulence factor detection	Early laboratory research (TRL 1–3); controlled laboratory feasibility	• Limited pathogen range • Matrix effects
	Phage-argonaute systems	Biological and enzymatic cascade amplification	Early laboratory research (TRL 1–3); laboratory proof-of-feasibility	• Phage availability • Reaction optimization
Nucleic acid platforms	Digital CRISPR-Cas	Absolute RNA quantification via microfluidic partitioning	Laboratory prototype validation (TRL 4–5); artificial matrix and laboratory validation	• Cost • Sample preparation complexity
	IoT-enabled networks	Real-time continuous monitoring with cloud analytics	Early laboratory research (TRL 1–3); pilot-scale feasibility studies	• Sensor drift • Data security • And power management
Integration systems	Multi-modal sensor fusion	Complementary detection principles with AI analysis	Early laboratory research (TRL 1–3); conceptual and laboratory feasibility	• System complexity • Data integration challenges

min^[80], while a nanozyme-catalyzed colorimetric microfluidic immunosensor achieved a limit of detection (LOD) of 9 CFU/mL within 20 min^[80,81]. Similarly, a sliding microfluidic chip integrated with MnO₂ nanoflowers achieved an LOD of 10 CFU/mL within 20 min^[82]. This technology exhibits clear advantages in detection sensitivity, portability, and operational integration. Nevertheless, its practical application in complex food matrices requires further validation, particularly regarding channel fouling, nonspecific adsorption, matrix interference, and the long-term stability of nanozyme activity.

Split-Protein Systems for Virulence Detection Split-protein systems represent an innovative approach for detecting active virulence in pathogenic bacteria, moving beyond general viability assessment. The PoreGlow system exemplifies this advancement by employing liposomes that encapsulate split green fluorescent protein (GFP) fragments. These fragments are released and complemented upon interaction with functional toxins such as listeriolysin O (LLO), which is secreted by viable *L. monocytogenes* cells^[83]. This method enables direct

assessment of pathogenic capability rather than mere presence, offering unprecedented specificity for virulent pathogens^[84,85]. The PoreGlow assay enables pre-enrichment-free detection within ca. 30 min, with detection limits of 0.17 µg/mL for LLO and 10 CFU/mL for *L. monocytogenes*^[83,86]. Compared with conventional methods requiring bacterial proliferation, this system is faster and more viability-relevant, as only actively virulent strains generate positive signals, thereby minimizing false positives from non-pathogenic relatives or dead cells^[87,88]. Although its current limitations include specificity to certain toxin-producing pathogens and matrix-dependent performance, the modular design allows adaptation to other pathogen-toxin systems through appropriate bioreceptor engineering. This innovation paves the way for viability assays that focus on pathogenic potential rather than cell presence alone.

Next-Generation Nucleic Acid Platforms

Phage-Argonaute Fluorescence Systems By integrating bacteriophage-mediated viable-cell recognition utilizing the argonaute protein from

Table 4. Comparison of key analytical parameters of emerging technological platforms for foodborne pathogen detection

Emerging technology	Limit of detection	Assay time	Multiplexing capability	Cost per test	Key advantages	Current limitations
Paper chromogenic array with neural networks (PCA-NN)	10-10 ³ CFU/mL ^[76,77]	1-2 h ^[76,77]	Multi-target recognition based on colorimetric fingerprints and neural-network classification	US\$0.1 per unit ^[77]	• Non-destructive detection • No complex sample pretreatment • Multiplexing capability	• Susceptible to interference from food matrices and background microflora
Nanozyme-enhanced microfluidic biosensors	10-10 ² CFU/mL ^[80,81]	20-25 min ^[81,82]	Multiplexing enabled by chip compartmentalization and nanozyme amplification	NA	• High integration • Fast detection speed^[82]	• Signal cross-talk in multiplex assays
Split-protein systems for virulence detection	10 CFU/mL ^[83,84]	30 min ^[83,84]	Multi-target detection not demonstrated in the current format	NA	• Virulence activity-based detection • Higher detection specificity	• Multiplexing not yet demonstrated • Risk of false negatives/positives
Phage-argonaute fluorescence systems	20-50 CFU/mL ^[89,90]	2.5-3 h ^[89,90]	Promising for multiplex detection of foodborne pathogens	NA	• Extraction-free and amplification-free	• Not suitable for high-throughput detection^[2]
Digital CRISPR-Cas platforms	1.0 aM-10 fM ^[96-98]	5-80 min	Multiplex detection in a single sample using different CRISPR enzymes	NA	• Absolute quantification • Single-molecule sensitivity • High specificity	• Droplet stability issues • High equipment/chip costs

Note. NA, Not Available ; Internet of Things (IoT)-Enabled Sensor Networks and Multi-Modal Sensor Fusion Platforms are not included in this table because they represent system-level integration strategies rather than standalone detection assays. Their detection limits, assay times, and multiplexing capabilities depend on the specific front-end sensors integrated into each system.

Clostridium butyricum (CbAgo), a phage-assisted nucleic-acid detection platform for viable pathogens has been established^[89]. This system combines phage-based enrichment of viable host cells with guide DNA-directed CbAgo cleavage of target bacterial DNA, thereby converting nucleic-acid recognition into an amplified fluorescence signal. The system employs sequential biological and enzymatic amplification: Phage infection ensures specificity for viable cells, while the CbAgo-mediated cleavage of the target DNA amplifies fluorescence. This dual-specificity approach achieves sensitive viable-pathogen detection, with an LOD of ca. 20–50 CFU/mL and a detection time of ca. 2.5–3 h, while preserving the biological relevance of phage-based assays^[90]. The incorporation of machine vision for signal detection and ML for pattern recognition further enhances its accuracy and throughput^[91]. Current research endeavors are directed towards expanding the phage library to achieve broader pathogen coverage and optimizing the CbAgo reaction conditions to ensure consistent performance in food samples. This technology holds significant promise for confirming infectious potential in conjunction with pathogen identification^[92].

Digital CRISPR-Cas Platforms The advancement of CRISPR-Cas technology into digital detection formats has improved nucleic acid-based pathogen quantification. Digital CRISPR platforms partition samples into numerous femtoliter-to-picoliter microreactors and generate endpoint binary readouts, enabling absolute quantification of target nucleic acids without the need for standard curves^[93]. This strategy provides a useful platform for viable-pathogen detection when combined with viability-discriminating approaches such as PMA treatment or viability-associated RNA targets^[94,95]. Amplification-free Cas13a-based digital RNA detection systems such as SATORI have achieved RNA detection at ca. 5 fM, while droplet/digital CRISPR formats further support absolute nucleic-acid quantification through reaction partitioning and endpoint readouts^[96-98]. When combined with viability-associated RNA markers or PMA-based live-cell discrimination, this digital format provides a promising route for accurate viable-pathogen detection. Portable imaging strategies, including smartphone-based readers, further support field deployment and point-of-care testing. However, the practical application of this approach will depend on simplified sample preparation, lower-cost microfluidic chip fabrication, and robust

performance in complex food matrices^[97,98].

Intelligent Integration Systems

Internet of Things (IoT)-Enabled Sensor Networks

The integration of viability sensors with Internet of Things (IoT) technology facilitates real-time, continuous monitoring throughout the food supply chain^[99]. These systems synergistically integrate nanoscale biosensors with wireless communication modules, cloud analytics, and predictive algorithms to establish comprehensive surveillance networks. Sensors are used to perform continuous spatiotemporal surveillance of environmental conditions and pathogen presence, transmitting data to centralized platforms for immediate analysis and action^[100]. Recent implementations of this approach have employed low-power wide-area network (LPWAN) and energy-harvesting technologies to enable long-term operation without the need for battery replacement^[101]. The machine-learning algorithms embedded within these networks both detect contamination and utilize predictive microbiology models to forecast pathogen proliferation based on environmental fluctuations^[102]. Although they are promising, the scalability of these ecosystems faces challenges regarding sensor bio-stability (anti-fouling), data sovereignty, and the standardization of interoperability protocols across heterogeneous platforms^[103].

Multi-Modal Sensor Fusion Platforms The most advanced emerging systems integrate multiple detection principles through sensor fusion approaches, capitalizing on the complementary strengths of nucleic-acid, metabolic, and immunological sensing^[104]. By incorporating modalities that specifically respond to metabolic activity, RNA transcripts, or toxin secretion, these platforms facilitate direct and confirmatory assessments of microbial viability rather than merely detecting presence. Typically, these systems consolidate all sensing layers within a single automated platform, enabling orthogonal verification of viability status. Advanced data-fusion algorithms, including deep-learning networks and Bayesian inference models, process the multi-modal data to generate comprehensive viability assessments with confidence scoring. By correlating DNA-, RNA-, and metabolite-based signals, these algorithms can differentiate between living and dead cells and adapt detection strategies based on the viability probability inferred from the preliminary results^[102]. This adaptive capability significantly

reduces false positives and enhances reliability. Although their complexity and cost currently limit widespread adoption, such platforms provide a roadmap for the next generation of integrated food safety systems—those capable of autonomously detecting, confirming, and quantifying viable pathogens with minimal human intervention. Collectively, these intelligent systems signify the transition from passive detection to active, adaptive, and predictive food-safety monitoring, representing the frontier of viability assessment technology.

AI Algorithms for Microbial Viability Decoding The primary AI algorithms for distinguishing microbial viability can be categorized into two principal branches: conventional ML and deep learning. Ultimately, the optimal architectural selection is determined by the specific data modality.

Machine Learning for Structured Data Analysis ML algorithms, such as SVMs and random forests, are primarily utilized for analyzing structured datasets, including spectral signatures and metabolic indices. For instance, by evaluating feature importance, random forest models can pinpoint critical spectral peaks associated with microbial viability, thereby enabling the precise differentiation of physiological states^[105].

Deep Learning for Spatiotemporal and Multi-Modal Data Analysis Deep-learning algorithms, particularly convolutional neural networks (CNNs) and recurrent neural networks (RNNs), exhibit exceptional proficiency in interpreting unstructured data modalities, including images and timeseries data. In morphological profiling, CNNs convert complex spatial features like membrane integrity and fluorescence into viability probabilities (live, dead, or VBNC). Enhancing U-Net with a position attention module (PAM) improves the Dice coefficient for segmentation^[106], allowing precise cell-membrane delineation and providing a robust computational basis for capturing the cryptic morphological phenotypes of VBNC cells. The Fusion-Net III model, a dual-stream hybrid neural network, exhibited a 100% accuracy rate in determining the viability states of six prevalent foodborne pathogens by concurrently analyzing spectral fingerprints and spatial image features within intricate multi-modal datasets^[107]. RNNs and Long Short-Term Memory networks (LSTMs) can be used to effectively interpret changes in microbial growth over time, facilitating the detection of viable cells from time-lapse microscopy images up to 12 h faster than conventional culture techniques^[108].

Practical Challenges and Next-Generation AI

Solutions The primary challenges in implementing AI models include their inherent "black-box" nature^[109] and issues related to data skewness^[110]. Tools such as SHapley Additive exPlanations (SHAP) address regulatory challenges by clarifying feature contributions, thereby enhancing transparency^[111]. In terms of addressing data skewness, the Synthetic Minority Oversampling Technique (SMOTE) effectively improves the F1 scores of pathogen-detection models from 0.7 to 0.9 by mitigating the majority-class bias caused by sample imbalance, thus improving the detection of foodborne pathogens^[112]. Next-generation AI algorithms, exemplified by self-supervised learning (SSL)^[113] and edge computing^[114], have exhibited significant potential for application. For instance, biological foundation models utilize SSL to mitigate reliance on large labeled datasets, thus enabling autonomous learning from the unannotated raw data. In terms of application, the integration of lightweight AI with edge devices such as smartphone spectrometers effectively reconciles the demands of intensive computational processes with the requirements of point-of-care testing (POCT). AI therefore provides a highly promising technological foundation for the real-time, non-destructive, and precise assessment of pathogen viability within complex food matrices.

CONCLUSION AND FUTURE PERSPECTIVES

Foodborne pathogen viability assessment has evolved from conventional culture-dependent analysis toward integrated molecular, metabolic, and functional detection strategies, particularly to address the limitations associated with VBNC cells. This review critically summarizes the major platforms and technologies for viability determination, including viability dye-coupled nucleic acid amplification, CRISPR-assisted systems, biosensors, bacteriophage-based assays, Raman spectroscopy, advanced microscopy, and AI-assisted analytical frameworks. Together, these technologies have shifted foodborne-pathogen detection from simple presence/absence analysis toward more biologically meaningful assessment of membrane integrity, metabolic activity, transcriptional signals, infectivity, and physiological functionality. Future advancements are anticipated to emerge from the convergence of multiple technological domains. The integration of biological recognition elements with nanomaterial-based amplification, microfluidic automation, and AI-driven data analytics represents a transformative pathway toward robust, field-

deployable systems. Intelligent algorithms capable of interpreting complex spectroscopic and cytometric data will enable precise differentiation of bacterial viability states. Equally important, the establishment of standardized protocols and certified VBNC reference materials will ensure reproducibility and comparability across laboratories and industrial applications. As the field progresses, the emphasis should shift toward multiplexed, real-time platforms capable of simultaneously identifying pathogens and assessing their viability in situ. Such next-generation systems will not only accelerate response times but also transition food-safety management from reactive detection to proactive prevention. Through interdisciplinary collaboration and continuous innovation, viability-based detection will evolve into a cornerstone of modern food surveillance, safeguarding public health within an increasingly complex global food network.

Funding This work was supported by the Shandong Province Key Research and Development Plan (Major Scientific and Technological Innovation Project) under the project titled "Key Technologies and Equipment Development and the Industrialization Application of Dairy Products Processing" (2023CXGC010711) and by the Beijing Medical Award Foundation (YXTL-2024-0002-0600).

Competing Interests The authors declare that there are no conflicts of interest regarding the submission of this manuscript. All of the authors have approved the manuscript for publication.

[&]These authors contributed equally to this work.

[#]Correspondence should be addressed to Hong Liu, PhD, E-mail: liuhongseminar@hotmail.com; Xinxin Shen, PhD, E-mail: shenxx@ivdc.chinacdc.cn

Biographical notes of the first authors: Zeyu Wang, Master's student, majoring in intelligent microbial detection, E-mail: Wangzeyu0701@hotmail.com; Anum Farid, PhD, majoring in intelligent microbial detection, E-mail: Anum.malik911@gmail.com

Received: March 4, 2026;

Accepted: August 8, 2026

REFERENCES

- Keddy KH, Hoffmann S, Founou LL, et al. Quantifying national burdens of foodborne disease—four imperatives for global impact. *PLoS Glob Public Health*, 2025; 5, e0004309.
- Haddad MA, Tarawneh H, Abu-Romman S, et al. Emerging analytical techniques for determination of foodborne pathogens and toxins: a review. *Discover Food*, 2025; 5, 370.
- Xiao LL, Li Z, Dou X, et al. Advancing microbial risk assessment: perspectives from the evolution of detection technologies. *npj Sci Food*, 2025; 9, 157.
- Wang J, Den Bakker HC, Denes TG. A critical review of metagenomic approaches for foodborne pathogen surveillance. *Crit Rev Food Sci Nutr*, 2025; 65, 8488–501.
- Darwesh OM, Mostafa A, El-Sayed HS, et al. Recent technologies for detection of milk contaminants: characterization and mechanism of action: a review article. *Discover Food*, 2025; 5, 140.
- Tchato P, Marion-Sanchez K, Lebielle T, et al. From dormancy to viability: the resuscitation processes of viable but non-culturable bacteria—a systematic review. *Microorganisms*, 2026; 14, 136.
- Khodaparast M, Sharley D, Marshall S, et al. Advances in point-of-care and molecular techniques to detect waterborne pathogens. *npj Clean Water*, 2024; 7, 74.
- Xu J, Akhtar M, Meng W, et al. Advances in pathogen detection: from traditional methods to nanotechnology, biosensing and AI integration. *WIREs Nanomed Nanobiotechnol*, 2025; 17, e70022.
- Guo LZ, Ze X, Jiao YX, et al. Development and validation of a PMA-qPCR method for accurate quantification of viable *Lactocaseibacillus paracasei* in probiotics. *Front Microbiol*, 2024; 15, 1456274.
- Tamura A, Azam AH, Nakamura T, et al. Synthetic phage-based approach for sensitive and specific detection of *Escherichia coli* O157. *Commun Biol*, 2024; 7, 535.
- Reynoso M, Chang AY, Wu Y, et al. 3D-printed, aptamer-based microneedle sensor arrays using magnetic placement on live rats for pharmacokinetic measurements in interstitial fluid. *Biosens Bioelectron*, 2024; 244, 115802.
- Park HW, Li ZG, Ma LY, et al. Deep learning enabled rapid detection of live bacteria in the presence of food debris. *npj Sci Food*, 2025; 9, 274.
- Banicod RJS, Tabassum N, Jo DM, et al. Integration of artificial intelligence in biosensors for enhanced detection of foodborne pathogens. *Biosensors*, 2025; 15, 690.
- Izgördü ÖK, Gurbanov R, Darcan C. Understanding the transition to viable but non-culturable state in *Escherichia coli* W3110: a comprehensive analysis of potential spectrochemical biomarkers. *World J Microbiol Biotechnol*, 2024; 40, 203.
- Arvaniti M, Vlachou E, Kourtelis M, et al. Monitoring sublethal injury in *Listeria monocytogenes* during heat treatment of pork frankfurter-type sausages: a single-cell vs. population level approach. *Foods*, 2025; 14, 3144.
- Mahdizade Ari M, Scholz KJ, Cieplik F, et al. Viable but non-cultivable state in oral microbiota: a critical review of an underexplored microbial survival strategy. *Front Cell Infect Microbiol*, 2025; 15, 1533768.
- Dinu LD, Al-Zaidi QJ, Matache AG, et al. Improving the efficiency of viability-qPCR with lactic acid enhancer for the selective detection of live pathogens in foods. *Foods*, 2024; 13, 1021.
- Balagurusamy R, Gopi L, Kumar DSS, et al. Significance of viable but non-culturable (VBNC) State in vibrios and other pathogenic bacteria: induction, detection and the role of resuscitation promoting factors (Rpf). *Curr Microbiol*, 2024; 81, 417.
- Zhang ML, Wang XS, Deng XF, et al. Viable but non-culturable state formation and resuscitation of different antibiotic-resistant *Escherichia coli* induced by UV/chlorine. *Water Res*, 2024; 261, 122011.
- Miyoshi SI, Ogasawara M, Niwaki S, et al. Efficient resuscitation of early-stage viable but non-culturable cells of *Vibrio cholerae* using treatment with proteolytic enzymes. *Appl Environ Microbiol*, 2026; 92, e0151325.

21. Wang W, Wang WS, Yi ZK, et al. An integrated transcriptome and proteomic investigation of the stress responses of viable but non-culturable (VBNC) state *Salmonella* Enteritidis triggered by sodium hypochlorite. *Food Sci Hum Wellness*, 2025; 14, 9250213.
22. Li XX, Ren Q, Sun ZB, et al. Resuscitation promotion factor: a pronounced bacterial cytokine in propelling bacterial resuscitation. *Microorganisms*, 2024; 12, 1528.
23. Li LY, Bae S. Quantitative detection and survival analysis of VBNC *Salmonella* Typhimurium in flour using droplet digital PCR and DNA-intercalating dyes. *Microbiol Spectr*, 2024; 12, e00249–24.
24. Shafi Z, Shahid M, Singh R. Food-borne bacterial pathogens: emerging approaches in detection and prevention. *Arch Microbiol*, 2026; 208, 109.
25. Kaur S, Bran L, Rudakov G, et al. Propidium monoazide is unreliable for quantitative live–dead molecular assays. *Anal Chem*, 2025; 97, 2914–21.
26. Fittipaldi M, Nocker A, Codony F. Progress in understanding preferential detection of live cells using viability dyes in combination with DNA amplification. *J Microbiol Methods*, 2012; 91, 276–89.
27. Thilakarathna SH, Stokowski T, Chui L. An improved real-time viability PCR assay to detect *Salmonella* in a culture-independent era. *Int J Mol Sci*, 2022; 23, 14708.
28. Qiao JL, Zhao ZY, Li YR, et al. Recent advances of food safety detection by nucleic acid isothermal amplification integrated with CRISPR/Cas. *Crit Rev Food Sci Nutr*, 2024; 64, 12061–82.
29. Lu YH, Yang H, Bai JR, et al. CRISPR-Cas based molecular diagnostics for foodborne pathogens. *Crit Rev Food Sci Nutr*, 2024; 64, 5269–89.
30. Feng W, Zhang HQ, Le XC. Signal amplification by the *trans*-cleavage activity of CRISPR-Cas systems: kinetics and performance. *Anal Chem*, 2023; 95, 206–17.
31. Jenniches L, Michaux C, Popella L, et al. Improved RNA stability estimation through Bayesian modeling reveals most *Salmonella* transcripts have subminute half-lives. *Proc Natl Acad Sci USA*, 2024; 121, e2308814121.
32. Zhang T, Zhou WH, Lin XY, et al. Light-up RNA aptamer signaling-CRISPR-Cas13a-based mix-and-read assays for profiling viable pathogenic bacteria. *Biosens Bioelectron*, 2021; 176, 112906.
33. Xue T, Lu Y, Yang H, et al. Isothermal RNA amplification for the detection of viable pathogenic bacteria to estimate the salmonella virulence for causing enteritis. *J Agric Food Chem*, 2022; 70, 1670–8.
34. Moon YJ, Lee SY, Oh SW. A review of isothermal amplification methods and food-origin inhibitors against detecting food-borne pathogens. *Foods*, 2022; 11, 322.
35. Gustafson GC, Geary B, Griffiths JS, et al. Feasibility of RNA-based pathogen detection in an infant formula matrix. *LWT*, 2026; 240, 118997.
36. Zhang LB, Zhu JB, Guo SJ, et al. Photoinduced electron transfer of DNA/Ag nanoclusters modulated by G-quadruplex/hemin complex for the construction of versatile biosensors. *J Am Chem Soc*, 2013; 135, 2403–6.
37. Fabiani L, Saroglia M, Galatà G, et al. Magnetic beads combined with carbon black-based screen-printed electrodes for COVID-19: a reliable and miniaturized electrochemical immunosensor for SARS-CoV-2 detection in saliva. *Biosens Bioelectron*, 2021; 171, 112686.
38. Belinato JR, Acquaro VR Jr, Dias Cappelini LT, et al. New prospects and problems in sample preparation methods for microbiome analysis. *TrAC Trends Anal Chem*, 2021; 143, 116356.
39. Shi DD, Li RW, Yu SN, et al. Ultrasensitive aptamer-based metal-organic framework-on-metal-organic framework platform for clinical detection of KPC-2 *Klebsiella pneumoniae* and multidrug-resistant *Acinetobacter baumannii*. *ACS Infect Dis*, 2025; 11, 2491–9.
40. Scoggin JL, Tan C, Nguyen NH, et al. An enzyme-based electrochemical biosensor probe with sensitivity to detect astrocytic versus glioma uptake of glutamate in real time in vitro. *Biosens Bioelectron*, 2019; 126, 751–7.
41. Li ZL, Liu Y, Chen XM, et al. Surface-modified mesoporous nanofibers for microfluidic immunosensor with an ultra-sensitivity and high signal-to-noise ratio. *Biosens Bioelectron*, 2020; 166, 112444.
42. Calabretta MM, Álvarez-Diduk R, Michelini E, et al. Nanolatern on paper for smartphone-based ATP detection. *Biosens Bioelectron*, 2020; 150, 111902.
43. Gan ZXE, Roslan MAM, Abd Shukor MY, et al. Advances in aptamer-based biosensors and cell-internalizing SELEX technology for diagnostic and therapeutic application. *Biosensors*, 2022; 12, 922.
44. Stangherlin S, Lui N, Lee JH, et al. Aptamer-based biosensors: from SELEX to biomedical diagnostics. *TrAC Trends Anal Chem*, 2025; 191, 118349.
45. Zhao LY, Yang HX, Zheng XK, et al. Dual signal amplification by polysaccharide and eATRP for ultrasensitive detection of CYFRA 21-1 DNA. *Biosens Bioelectron*, 2020; 150, 111895.
46. Yousefniyehjromi Z, Sembranti L, Vivaldi F, et al. Label-free impedance-based aptasensor for *Escherichia coli* detection in real matrices. *Microchem J*, 2025; 208, 112595.
47. Mishra A, Roy S, Israel Shaikh N, et al. Recent advances in multiplex aptasensor detection techniques for food-borne pathogens: a comprehensive review of novel approaches. *Biosens Bioelectron X*, 2024; 16, 100417.
48. Wang B, Wang H, Lu XB, et al. Recent advances in electrochemical biosensors for the detection of foodborne pathogens: current perspective and challenges. *Foods*, 2023; 12, 2795.
49. Bottari B, Santarelli M, Neviani E. Determination of microbial load for different beverages and foodstuff by assessment of intracellular ATP. *Trends Food Sci Technol*, 2015; 44, 36–48.
50. Parreiras PM, Vieira Nogueira JA, Da Cunha LR, et al. Effect of thermosonication on microorganisms, the antioxidant activity and the retinol level of human milk. *Food Control*, 2020; 113, 107172.
51. Franssen F, Gerard C, Cozma-Petrut A, et al. Inactivation of parasite transmission stages: efficacy of treatments on food of animal origin. *Trends Food Sci Technol*, 2019; 83, 114–28.
52. Martins RO, Assunção Magalhães MF, Valeria Perez Vite WE, et al. From concept to application: exploring trends and guiding developments in miniaturized sample preparation for food contaminant analysis. *TrAC Trends Anal Chem*, 2025; 191, 118376.
53. Klaiman K, Ortega DL, Garnache C. Perceived barriers to food packaging recycling: evidence from a choice experiment of US consumers. *Food Control*, 2017; 73, 291–9.
54. Al-Hindi RR, Teklemariam AD, Alharbi MG, et al. Bacteriophage-based biosensors: a platform for detection of foodborne bacterial pathogens from food and environment. *Biosensors*, 2022; 12, 905.
55. Meile S, Kilcher S, Loessner MJ, et al. Reporter phage-based detection of bacterial pathogens: design guidelines and recent developments. *Viruses*, 2020; 12, 944.
56. Schmelcher M, Loessner MJ. Application of bacteriophages for detection of foodborne pathogens. *Bacteriophage*, 2014; 4, e28137.
57. Calimci M, Tezcan T, Tayyarcan EK, et al. Bacteriophage-based live bacteria detection for rapid infection diagnosis.

- Talanta, 2025; 286, 127569.
58. Fernandes E, Martins VC, Nóbrega C, et al. A bacteriophage detection tool for viability assessment of *Salmonella* cells. *Biosens Bioelectron*, 2014; 52, 239–46.
 59. Li F, Jiang XX, Zhang YZ, et al. Phage-based biosensors: from recognition to signal transduction empowered by amplification strategies for detecting foodborne pathogens. *Trends Food Sci Technol*, 2026; 168, 105492.
 60. Awais R, Fukudomi H, Miyanaga K, et al. A recombinant bacteriophage-based assay for the discriminative detection of culturable and viable but nonculturable *Escherichia coli* O157: H7. *Biotechnol Prog*, 2006; 22, 853–9.
 61. Martino MM, Briquez PS, Maruyama K, et al. Extracellular matrix-inspired growth factor delivery systems for bone regeneration. *Adv Drug Deliv Rev*, 2015; 94, 41–52.
 62. Butler HJ, Ashton L, Bird B, et al. Using Raman spectroscopy to characterize biological materials. *Nat Protoc*, 2016; 11, 664–87.
 63. Yuan SY, Chen YW, Lin KC, et al. Single cell Raman spectroscopy deuterium isotope probing for rapid antimicrobial susceptibility test of *Elizabethkingia* spp. *Front Microbiol*, 2022; 13, 876925.
 64. Sun Y, Tang HB, Zou XB, et al. Raman spectroscopy for food quality assurance and safety monitoring: a review. *Curr Opin Food Sci*, 2022; 47, 100910.
 65. Ho CS, Jean N, Hogan CA, et al. Rapid identification of pathogenic bacteria using Raman spectroscopy and deep learning. *Nat Commun*, 2019; 10, 4927.
 66. Dufrêne YF, Ando T, Garcia R, et al. Imaging modes of atomic force microscopy for application in molecular and cell biology. *Nat Nanotechnol*, 2017; 12, 295–307.
 67. Cheng YF. Single-particle cryo-EM—How did it get here and where will it go. *Science*, 2018; 361, 876–80.
 68. Falk T, Mai D, Bensch R, et al. U-Net: deep learning for cell counting, detection, and morphometry. *Nat Methods*, 2019; 16, 67–70.
 69. Christiansen EM, Yang SJ, Ando DM, et al. *In Silico* labeling: predicting fluorescent labels in unlabeled images. *Cell*, 2018; 173, 792–803. e19.
 70. Eulenberg P, Köhler N, Blasi T, et al. Reconstructing cell cycle and disease progression using deep learning. *Nat Commun*, 2017; 8, 463.
 71. Rahman HU, Sikder R, Tripathi M, et al. Machine learning-assisted Raman spectroscopy and SERS for bacterial pathogen detection: clinical, food safety, and environmental applications. *Chemosensors*, 2024; 12, 140.
 72. Punjani A, Rubinstein JL, Fleet DJ, et al. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods*, 2017; 14, 290–6.
 73. Shwaiiki LN, Lynch KM, Arendt EK. Future of antimicrobial peptides derived from plants in food application – a focus on synthetic peptides. *Trends Food Sci Technol*, 2021; 112, 312–24.
 74. Hussain A, Abbas Q, Nadeem M, et al. Meat-borne bacterial pathogen detection: conventional, molecular and emerging AI-based strategies. *Diagnostics*, 2026; 16, 1360.
 75. Shewale M, Razban A, Deshmukh S, et al. Design, development and implementation of the position estimator algorithm for harmonic motion on the XY flexural mechanism for high precision positioning. *Sensors*, 2020; 20, 662.
 76. Yang MY, Liu XB, Luo YG, et al. Machine learning-enabled non-destructive paper chromogenic array detection of multiplexed viable pathogens on food. *Nat Food*, 2021; 2, 110–7.
 77. Jia Z, Luo YG, Wang DY, et al. Nondestructive multiplex detection of foodborne pathogens with background microflora and symbiosis using a paper chromogenic array and advanced neural network. *Biosens Bioelectron*, 2021; 183, 113209.
 78. Yang LY, Xu XY, Song Y, et al. Research progress of nanozymes in colorimetric biosensing: classification, activity and application. *Chem Eng J*, 2024; 487, 150612.
 79. Mazouz Z, Mokni M, Fourati N, et al. Computational approach and electrochemical measurements for protein detection with MIP-based sensor. *Biosens Bioelectron*, 2020; 151, 111978.
 80. Jiang F, Jin NN, Wang L, et al. A multimetallic nanozyme enhanced colorimetric biosensor for *Salmonella* detection on finger-actuated microfluidic chip. *Food Chem*, 2024; 460, 140488.
 81. Niu YD, Jiang JT, Zhi X, et al. A sliding microfluidic chip-integrated colorimetric biosensor using MnO₂ nanoflowers for rapid *Salmonella* detection. *Micromachines*, 2025; 16, 904.
 82. Cheng XD, Chen H, Li WJ, et al. Nanozyme-catalyzed colorimetric microfluidic immunosensor for the filtration enrichment and ultrasensitive detection of *Salmonella typhimurium* in food samples. *Anal Chem*, 2025; 97, 6454–63.
 83. Guk K, Yi S, Kim H, et al. PoreGlow: a split green fluorescent protein-based system for rapid detection of *Listeria monocytogenes*. *Food Chem*, 2024; 438, 138043.
 84. Gercke D, Lenz F, Jose J. Split-GFP complementation at the bacterial cell surface for antibody-free labeling and quantification of heterologous protein display. *Enzyme Microb Technol*, 2024; 174, 110391.
 85. Tamura R, Jiang FC, Xie J, et al. Multiplexed labeling of cellular proteins with split fluorescent protein tags. *Commun Biol*, 2021; 4, 257.
 86. Zhao JX, Jedlicka SS, Lannu JD, et al. Liposome-doped nanocomposites as artificial-cell-based biosensors: detection of Listeriolysin O. *Biotechnol Prog*, 2006; 22, 32–7.
 87. Jin NN, Yang FZ, Zhang XY, et al. Sensitive detection of multiplex bacteria based on finger driven microfluidics and recombinase aided amplification. *Biosens Bioelectron*, 2025; 287, 117750.
 88. Jin NN, Jiang F, Yang FZ, et al. Multiplex nanozymatic biosensing of *Salmonella* on a finger-actuated microfluidic chip. *Lab Chip*, 2024; 24, 2712–20.
 89. Zhao JP, Han MJ, Ma AM, et al. A machine vision-assisted Argonaute-mediated fluorescence biosensor for the detection of viable *Salmonella* in food without convoluted DNA extraction and amplification procedures. *J Hazard Mater*, 2024; 466, 133648.
 90. Wang XF, Hu XB, Pan SX, et al. DNA and ATP synergistically triggered Argonaute-mediated sensor for the ultrasensitive detection of viable *Salmonella* without DNA extraction and amplification. *Sens Actuators B Chem*, 2024; 408, 135543.
 91. Stadnicka D, Sep J, Amadio R, et al. Industrial needs in the fields of artificial intelligence, internet of things and edge computing. *Sensors*, 2022; 22, 4501.
 92. Li LT, Feng N, Chen YP. Prokaryotic Argonaute-based multiplexed detection assays for food safety: recent advances and perspectives. *Trends Food Sci Technol*, 2025; 164, 105243.
 93. Dietl S, Sobek H, Mizaikoff B. Epitope-imprinted polymers for biomacromolecules: recent strategies, future challenges and selected applications. *TrAC Trends Anal Chem*, 2021; 143, 116414.
 94. Yin WH, Hu K, Yang YX, et al. A propidium monoazide-assisted digital CRISPR/Cas12a assay for selective detection of live bacteria in sample. *Anal Chem*, 2024; 96, 17941–9.
 95. Shinoda H, Iida T, Makino A, et al. Automated amplification-

- free digital RNA detection platform for rapid and sensitive SARS-CoV-2 diagnosis. *Commun Biol*, 2022; 5, 473.
96. Shinoda H, Taguchi Y, Nakagawa R, et al. Amplification-free RNA detection with CRISPR-Cas13. *Commun Biol*, 2021; 4, 476.
 97. Zhang Y, Walker RSK, Sunna A, et al. Droplet digital CRISPR for nucleic acid detection. *Adv Sci*, 2026; 13, e17470.
 98. Politza AJ, Nouri R, Guan WH. Digital CRISPR systems for the next generation of nucleic acid quantification. *TrAC Trends Anal Chem*, 2023; 159, 116917.
 99. Cheng H, Chen L, McClements DJ, et al. Starch-based biodegradable packaging materials: a review of their preparation, characterization and diverse applications in the food industry. *Trends Food Sci Technol*, 2021; 114, 70–82.
 100. Dong X, Raghavan V. A comprehensive overview of emerging processing techniques and detection methods for seafood allergens. *Compr Rev Food Sci Food Saf*, 2022; 21, 3540–57.
 101. Mekki K, Bajic E, Chaxel F, et al. A comparative study of LPWAN technologies for large-scale IoT deployment. *ICT Express*, 2019; 5, 1–7.
 102. Cappelli A, Bettaccini L, Cini E. The kneading process: a systematic review of the effects on dough rheology and resulting bread characteristics, including improvement strategies. *Trends Food Sci Technol*, 2020; 104, 91–101.
 103. Sicari S, Rizzardi A, Grieco LA, et al. Security, privacy and trust in Internet of Things: the road ahead. *Comput Networks*, 2015; 76, 146–64.
 104. Dincer C, Bruch R, Costa-Rama E, et al. Disposable sensors in diagnostics, food, and environmental monitoring. *Adv Mater*, 2019; 31, 1806739.
 105. Rani P, Kotwal S, Manhas J, et al. Machine learning and deep learning based computational approaches in automatic microorganisms image recognition: methodologies, challenges, and developments. *Arch Comput Methods Eng*, 2022; 29, 1801–37.
 106. Azad R, Aghdam EK, Rauland A, et al. Medical image segmentation review: the success of U-Net. *IEEE Trans Pattern Anal Mach Intell*, 2024; 46, 10076–95.
 107. Park B, Shin T, Wang B, et al. Classification between live and dead foodborne bacteria with hyperspectral microscope imagery and machine learning. *J Microbiol Methods*, 2023; 209, 106739.
 108. Wang HD, Ceylan Koydemir H, Qiu YZ, et al. Early detection and classification of live bacteria using time-lapse coherent imaging and deep learning. *Light Sci Appl*, 2020; 9, 118.
 109. Rudin C. Stop explaining black box machine learning models for high stakes decisions and use interpretable models instead. *Nat Mach Intell*, 2019; 1, 206–15.
 110. Mairi A, Hamza L, Touati A. Artificial intelligence and its application in clinical microbiology. *Expert Rev Anti-Infect Ther*, 2025; 23, 469–90.
 111. Lundberg SM, Erion G, Chen H, et al. From local explanations to global understanding with explainable AI for trees. *Nat Mach Intell*, 2020; 2, 56–67.
 112. Chung CR, Wang HY, Yao CH, et al. Data-driven two-stage framework for identification and characterization of different antibiotic-resistant *Escherichia coli* isolates based on mass spectrometry data. *Microbiol Spectr*, 2023; 11, e0347922.
 113. Ben Atitallah S, Ben Rabah C, Driss M, et al. Self-supervised learning for graph-structured data in healthcare applications: a comprehensive review. *Comput Biol Med*, 2025; 188, 109874.
 114. Momeni A, Rahmani B, Scellier B, et al. Training of physical neural networks. *Nature*, 2025; 645, 53–61.